

PATENT APPLICATION TRANSMITTAL LETTER

PATENT GROUP
FOLEY, HOAG & ELIOT LLP
ONE POST OFFICE SQUARE
BOSTON, MA 02109-2170

Atty Docket No: BIV-052.02

A

jc490 U.S. PTO
09/435733
11/08/99

11/08/99

jc490 U.S. PTO
09/435733
11/08/99

the Assistant Commissioner for Patents:

Transmitted herewith for filing under 35 U.S.C. 111 and 37 CFR 1.53 is the patent application of:

Alphonse Galdes and Nagesh Mahanthappaentitled: Methods and Composition for Treating or Preventing Peripheral Neuropathies

Enclosed are:

- (X) 124 pages of written description, claims and abstract.
(X) 25 sheets of drawings.
() an assignment of the invention to _____
() executed declaration of the inventors.
() a certified copy of a _____ application.
() a verified statement to establish small entity status under 37 CFR 1.9 and 1.27.
(X) other: unexecuted Declaration, Petition and Power of Attorney.

CLAIMS AS FILED

	# FILED	# EXTRA	Rate	FEE	Rate (Sm. Entity)	FEE
BASIC FEE			\$760		\$380	
TOTAL CLAIMS	-20=		x \$18		x \$9	
INDEPENDENT CLAIMS	-3=		x \$78		x \$39	
MULTIPLE DEP. CLAIMS			\$260		\$130	
* NO. EXTRA MUST BE ZERO OR LARGER		TOTAL				\$

- () A check in the amount of \$ _____ to cover the filing fee is enclosed.
() The Commissioner is hereby authorized to charge and credit Deposit Account No. 06-1448 as described below. A duplicate copy of this sheet is enclosed.
() Charge the amount of \$ _____ as filing fee.
() Credit any overpayment.
() Charge any additional filing fees required under 37 CFR 1.16 and 1.17.

Certificate of Express Mail

Express Mail Label: EL354726015US

I hereby certify that this correspondence is being deposited with the United States Postal Service as Express Mail, postage prepaid, "Post Office to Addressee", in an envelope addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231 on the date specified below:

November 8, 1999

Date of Deposit

Carmen Parra

Carmen Parra

Matthew P. Vincent, Ph.D.
Reg. No. 36,709

Methods and Compositions for Treating or Preventing Peripheral Neuropathies

Reference to Related Application

The present application is a continuation-in-part of USSN 09/187,387, filed 6 November
5 1998, the specification of which is incorporated by reference herein.

Background of the Invention

Conditions that affect components of a motor unit (motor neuron cells of the spinal cord,
nerve, neuromuscular junction, and muscle fibers), sensory and autonomic nerves or their
supportive structures are included in the broad category of "neuromuscular disorders", and
10 include peripheral neuropathies.

Motor nerves are responsible for voluntary movement. Their cell bodies lie within the
spinal cord, and their processes transmit signals outward to specialized motor receptors on the
skeletal muscles. Sensory nerves allow the sensation of pain, vibrations or touch, and sense
where limbs are positioned in space. Their cell bodies are grouped in specialized structures called
5 sensory "ganglia" next to the spinal cord. And they transmit signals from sensory receptors in the
skin and other organs inward to the central nervous system (CNS). Autonomic nerves control
involuntary functions like breathing, heartbeat, blood pressure, digestion and sexual function.
Their cell bodies, clustered in autonomic ganglia, are spread throughout the body.

Neuropathy is a generic term used to describe diseases of the peripheral nervous system.
20 There are about 200 known different causes of peripheral neuropathies. Although most
neuropathies affect all three types of nerve fibers, to varying degrees, some diseases involve only
one or two, and are thus said to be purely or predominantly motor, sensory, or autonomic
neuropathies.

For instance, Guillain-Barré syndrome is an acute illness involving the peripheral
25 nervous system that usually occurs two to three weeks after a flu-like disease or other infections.
It is mostly a motor neuropathy, meaning that its symptoms are largely related to the
involvement of the motor nerves. Despite the primarily motor nature of the disease, the earliest
symptoms may be numbness and tingling felt in the lower extremities followed shortly by
weakness of the distal muscles of the lower extremities. The common early symptoms reported
30 by patients are those of tripping on the toes that later results in a footdrop. The weakness usually
ascends to involve the entire lower extremities and later the upper extremities. The danger occurs
when the weakness involves the muscles of respiration.

The diagnosis of Guillain-Barré syndrome is suggested when the patient presents with a history of ascending weakness and a physical examination consistent with a primarily motor neuropathy. The diagnosis is confirmed with the performance of a spinal tap, which usually shows elevation of the protein level in the spinal fluid without an increase in the number of white cells and by an electromyogram. All other conditions resembling Guillain-Barré syndrome must also be excluded.

Although Guillain-Barré syndrome is usually a self-limiting illness, intensive therapeutic intervention is often needed.

CIDP or chronic inflammatory demyelinating polyneuropathy is an immune-mediated neuropathy that affects the peripheral motor and sensory nerves. The symptoms are of a slowly progressive numbness and tingling that usually starts in the feet, but later spreads to the legs and hands. The patients also complain of some weakness, again usually starting in the lower extremities, but soon involving the upper extremities. With further involvement of the sensory system, other modalities of sensations, such as balance, are affected and the patients complain of inability to walk or maintain balance in the dark.

The diagnosis of CIDP is suspected with a history of progressive sensorimotor neuropathy. Physical examination consistent with distal sensory loss in the upper and lower extremities, in conjunction with motor weakness that can be more proximal than distal supports the clinical diagnosis. The spinal tap usually shows a significant rise in the protein level of the spinal fluid. Electromyography with nerve conduction studies also supports the diagnosis. Usually the main picture is that of slowing of the conduction velocities of the peripheral nerves. The final diagnostic step would be the performance of a nerve biopsy. Finding of inflammation on the nerve biopsy, although rare, definitely confirms the diagnosis. However, the absence of inflammation does not entirely rule it out. Findings of predominant demyelination on the nerve biopsy can be used in conjunction with the other studies and the clinical presentation to suggest a diagnosis of CIDP. Once the diagnosis is secured, treatment with immunosuppressive medications can be initiated. The first line of treatment remains high-dose steroids that are initiated orally every day and then slowly tapered over time depending on the patient's improved symptomatology. Steroid failure or intolerance to steroids necessitates the use of other immunosuppressing agents. However, better therapeutic intervention for CIDP is still a desired objective of the present invention.

Peripheral neuropathy is one of the many complications of long-standing diabetes. Usually neuropathy occurs about 8 to 10 years after the onset of diabetes. However, it is not uncommon to see patients presenting with neuropathic symptoms that have their diabetes diagnosed at that time or patients with 20 or more years of diabetes with little or no evidence of

neuropathy. The symptoms of diabetic neuropathy consist of a slow and insidious numbness and tingling of the lower extremities that can progress to become a painful neuropathy. The pain is usually described as a burning sensation in the feet. Occasionally, the pain is described as a sensation of sharp, electric jolts traveling down the lower extremities. As it worsens, the pain acquires a deep bony nature. It tends to be worse at night commonly preventing or awakening the patients from sleep. As the neuropathy worsens, it affects the upper extremities and may involve the motor nerves with the complaint of weakness in the distal muscles of the legs and arms. The neuropathy of diabetes can also involve the autonomic nervous system causing problems with sweating, blood pressure, and sexual function.

Diabetic neuropathy is suspected when the patient's history and physical examination are compatible with the clinical picture in a setting of diabetes. In the absence of the history of diabetes, diagnostic tests to rule out diabetes is required. The workup is completed by the performance of an electromyogram with nerve conduction studies to quantitate the extent of involvement of the peripheral nervous system.

Diabetic neuropathy, unfortunately, has no effective treatment at this point in the art. Adequate control of the patient's blood sugar, however, has been shown to slow the progression of the symptoms. Symptomatic treatment with various medications that suppress neuropathic pain, including Elavil, Tegretol and more recently Ultram, have been successful. Thus, a more effective treatment for diabetic neuropathy is an objective of the present invention.

Other common causes of neuropathy such include alcoholism or medication induced neuropathies, as well as inherited forms of such disorders.

Summary of the Invention

One aspect of the present application relates to a method for treating or alleviating all or a portion of the symptoms attendant neuromuscular disorders, and in particular, in the treatment of peripheral neuropathies. Briefly, the subject method comprises contacting the afflicted tissue with a hedgehog therapeutic or ptc therapeutic (defined *infra*) in an amount effective to alter the growth state of the treated cells, e.g., relative to the absence of administration of the hedgehog therapeutic or ptc therapeutic.

Wherein the subject method is carried out using a *hedgehog* therapeutic, the *hedgehog* therapeutic preferably a polypeptide including a *hedgehog* portion comprising at least a bioactive extracellular portion of a *hedgehog* protein, e.g., the *hedgehog* portion includes at least 50, 100 or 150 (contiguous) amino acid residues of an N-terminal half of a *hedgehog* protein. In preferred

embodiments, the *hedgehog* portion includes at least a portion of the *hedgehog* protein corresponding to a 19kd fragment of the extracellular domain of a *hedgehog* protein.

In preferred embodiments, the *hedgehog* portion has an amino acid sequence at least 60, 75, 85, or 95 percent identical with a hedgehog protein of any of SEQ ID Nos. 10-18 or 20, though sequences identical to those sequence listing entries are also contemplated as useful in the present method. The *hedgehog* portion can be encoded by a nucleic acid which hybridizes under stringent conditions to a nucleic acid sequence of any of SEQ ID Nos. 1-9 or 19, e.g., the *hedgehog* portion can be encoded by a vertebrate *hedgehog* gene, especially a human *hedgehog* gene.

In other embodiments, the subject method can be carried out by administering a gene activation construct, wherein the gene activation construct is designed to recombine with a genomic *hedgehog* gene of the patient to provide a heterologous transcriptional regulatory sequence operatively linked to a coding sequence of the *hedgehog* gene.


In still other embodiments, the subject method can be practiced with the administration of a gene therapy construct encoding a *hedgehog* polypeptide. For instance, the gene therapy construct can be provided in a composition selected from a group consisting of a recombinant viral particle, a liposome, and a poly-cationic nucleic acid binding agent,

In yet other embodiments, the subject method can be carried out using a *ptc* therapeutic. An exemplary *ptc* therapeutic is a small organic molecule which binds to a *patched* protein and derepresses *patched*-mediated inhibition of mitosis, e.g., a molecule which binds to *patched* and mimics *hedgehog*-mediated *patched* signal transduction, which binds to *patched* and regulates *patched*-dependent gene expression. For instance, the binding of the *ptc* therapeutic to *patched* may result in upregulation of *patched* and/or *gli* expression.

In a more generic sense, the *ptc* therapeutic can be a small organic molecule which interacts with MK cells to induce *hedgehog*-mediated *patched* signal transduction, such as by altering the localization, protein-protein binding and/or enzymatic activity of an intracellular protein involved in a *patched* signal pathway. For instance, the *ptc* therapeutic may alter the level of expression of a *hedgehog* protein, a *patched* protein or a protein involved in the intracellular signal transduction pathway of *patched*.

In certain embodiments, the *ptc* therapeutic is an antisense construct which inhibits the expression of a protein which is involved in the signal transduction pathway of *patched* and the expression of which antagonizes *hedgehog*-mediated signals. The antisense construct is preferably an oligonucleotide of about 20-30 nucleotides in length and having a GC content of at least 50 percent.

In other embodiments, the *ptc* therapeutic is an inhibitor of protein kinase A (PKA), such as a 5-isoquinolinesulfonamide. The PKA inhibitor can be a cyclic AMP analog. Exemplary PKA inhibitors include N-[2-((p-bromocinnamyl)amino)ethyl]-5-isoquinolinesulfonamide, 1-(5-isoquinoline-sulfonyl)-2-methylpiperazine, KT5720, 8-bromo-cAMP, dibutyryl-cAMP and PKA Heat Stable Inhibitor isoform α . Another exemplary PKA inhibitor is represented in the general formula:



wherein,

R_1 and R_2 each can independently represent hydrogen, and as valence and stability permit a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(CH_2)_m-R_8$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O-(CH_2)_m-R_8$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_8$, or

R_1 and R_2 taken together with N form a heterocycle (substituted or unsubstituted);

R_3 is absent or represents one or more substitutions to the isoquinoline ring such as a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, $-(CH_2)_m-R_8$, $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O-(CH_2)_m-R_8$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_8$;

R_8 represents a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle; and

n and m are independently for each occurrence zero or an integer in the range of 1 to 6.

Brief Description of the Figures

Figure 1. Variation of the weight of animals during the study in treated or control mice: control SHH=animals treated with 500 ug/kg SHH, without cisplatin; veh=vehicle group treated with cisplatin 2 mg/kg/day during 14 days; SHH500=animals treated with 500 ug/kg SHH and cisplatin; SHH50=animals treated with 50 ug/kg SHH and cisplatin. The compounds were administered 3 times per week subcutaneously. The weights are expressed in grams, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$; ***:significantly different at $p<0.001$.

Figure 2. Number of animals present throughout the study in treated or control mice. The number of animals in each group was compared by repeated Anova test and was not found to be significantly different between groups.

Figure 3. Time course of sensory nerve conduction velocity (SNCV) measured in treated or control mice. Results are expressed in m/sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$; ***:significantly different at $p<0.001$.

Figure 4. Tail flick latency measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$; ***:significantly different at $p<0.001$.

Figure 5. Latency to lick the paw measured in treated or control mice. Results are expressed in sec as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test.

Figure 6. Latency before first jump measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$.

Figure 7. Latency before adjusted jump measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test.

Figure 8. Ability to stay on rotarod measured in treated or control mice.

Figure 9. Duration of the walk on a rod needed to reach the platform, measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$; ***: significantly different at $p<0.001$.

Figures 10A and 10B. Ability to hold a weight with four limbs (10a) or only forelimbs (10b) measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$.

Figures 11A and 11B. Maximal strength exercised with four limbs (11a) or only forelimbs (11b) measured in treated or control mice. Results are expressed in sec, as means \pm SEM. Post-hoc comparison to vehicle group was performed with Fisher test; *:significantly different at $p<0.05$; **:significantly different at $p<0.01$; ***:significantly different at $p<0.001$.

Figure 12. Graph of motor neuron velocity in normal and Dhh^{-/-} mice

Figures 13A and 13B. Micrographs of peripheral nerve cells in normal and Dhh^{-/-} mice.

Figures 14A and 14B. Immunohistochemical stains of peripheral nerves using antibodies for neurofilament (an axonal marker) and Laminin (and ECM/connective tissue marker).

Figure 15. Effects of hedgehog on perineural cell proliferation.

Figure 16. Running time (walking test) in control and treated mice.

Figure 17. Time before falling from the rotarod in control and treated mice.

Figure 18. Histological study of SOD mice treated with 500 μ g/kg SHH. Motoneurons were counted in ventral horns of lumbar spinal cord sections originating from 100 day-old hSOD mice, after cresyl violet staining.

Figure 19. Histological study of SOD mice treated with 500 μ g/kg SHH (without Y0 littermate).

Figure 20. Histological study of male SOD mice treated with 500 μ g/kg SHH.

Figure 21. Histological study of female SOD mice treated with 500 μ g/kg SHH

Figure 22. Evaluating the effect of Hedgehog proteins on ability to grip following sciatic nerve crush injury.

Figure 23. Evaluating the effect of Hedgehog protein on sensory nerve conduction velocity in galactose intoxication-mediated neuropathies. CA= normal animal treated with control; CB= normal animal treated with Shh; GA= galactose intoxicated animal treated with vehicle; and GB= galactose intoxicated animal treated with Shh.

The Peripheral Nervous System is one of the two main divisions of the body's nervous system. The other is the Central Nervous System, which includes the brain and spinal cord. "Peripheral" means away from the center: and this system contains the nerves that connect the Central Nervous System to the muscles, skin and internal organs.

5 Peripheral Neuropathy is the term used to describe disorders resulting from injury (e.g., mechanical, chemical, viral, bacterial or genetic) to the peripheral nerves. It can be caused by diseases that affect only the peripheral nerves or by conditions that affect other parts of the body as well. Dymptoms almost always involve weakness, numbness or pain - usually in the arms and legs. It will be helpful for you to know a few basics of nerve biology to understand how
10 neuropathy gets started.

I. Overview

15 The present application is directed to the discovery that hedgehog gene products are able to protect peripheral nerve cells under conditions which otherwise result in peripheral neuropathy. Certain aspects of the invention are directed to preparations of hedgehog polypeptides, or other molecules which regulate *patched* or *smoothened* signalling, and their uses as protective agents against both acquired and hereditary neuropathies. As used herein, "peripheral neuropathy" refers to a disorder affecting a segment of the peripheral nervous system. For instance, the method of the present invention can be used as part of a treatment program in
20 the management of neuropathies associated with systemic disease, e.g., viral infections, diabetes, inflammation; as well as genetically acquired (hereditary) neuropathies, e.g., Charcot-Marie-Tooth disease; and neuropathies caused by a toxic agent, e.g., a chemotherapeutic agent such as vincristine.

25 To further illustrate, the subject method can be used in the treatment of such acquired neuropathies as diabetic neuropathies; immune-mediated neuropathies such as Guillain-Barre syndrome (GBS) and variants, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic polyneuropathies with antibodies to peripheral nerves, neuropathies associated with vasculitis or inflammation of the blood vessels in peripheral nerve, brachial or lumbosacral plexitis, and neuropathies associated with monoclonal gammopathies; neuropathies associated
30 with tumors or neoplasms such as sensory neuropathy associated with lung cancer, neuropathy associated with multiple myeloma, neuropathy associated with waldenstrom's macroglobulemia, chronic lymphocytic leukemia, or B-cell lymphoma; neuropathy associated with amyloidosis; neuropathies caused by infections; neuropathies caused by nutritional imbalance; neuropathy in kidney disease; hypothyroid neuropathy; neuropathy caused by alcohol and toxins; neuropathies

caused by drugs; neuropathy resulting from local irradiation; neuropathies caused by trauma or compression; idiopathic neuropathies

Likewise, the subject method can be used in the treatment of such hereditary neuropathies as Charcot-Marie Tooth Disease (CMT); Familial Amyloidotic Neuropathy and other Hereditary Neuropathies; and Hereditary Porphyria.

In another embodiment, the subject method can be used to inhibit or otherwise slow neurodegenerative events associated with age-related neuropathology.

As described in the appended examples, *hedgehog* proteins are neuroprotective under conditions which promote chemical lesioning of peripheral nerves. Indeed, *hedgehog* proteins showed a significant protective effect that was similar to the reported effect of NGF. Based upon its neurotrophic and neuroprotective activities, the administration of *hedgehog* or *ptc* therapeutics is suggested herein as a treatment for several types of neurodegenerative diseases including neuropathies. In general, the method of the present invention comprises administering to animal, or to cultured peripheral nerves in vitro, an amount of a *hedgehog* or *ptc* therapeutic (defined *infra*) which produces a non-toxic response by the cell of resistance to degeneration, e.g., marked by loss of differentiation, apoptosis and/or necrosis. The subject method can be carried out on cells which may be either dispersed in culture or a part of an intact tissue or organ. Moreover, the method can be performed on cells which are provided in culture (*in vitro*), or on cells in a whole animal (*in vivo*).

In one aspect, the present invention provides pharmaceutical preparations and methods for treating or preventing neuropathies utilizing, as an active ingredient, a *hedgehog* polypeptide or a mimetic thereof. The invention also relates to methods of controlling the functional performance of peripheral nerve cells by use of the pharmaceutical preparations of the invention.

The subject *hedgehog* treatments are effective on both human and animal subjects afflicted with these conditions. Animal subjects to which the invention is applicable extend to both domestic animals and livestock, raised either as pets or for commercial purposes. Examples are dogs, cats, cattle, horses, sheep, hogs and goats.

Without wishing to be bound by any particular theory, the neuroprotective effect of *hedgehog* treatment may be due at least in part to the ability of these proteins to antagonize (directly or indirectly) *patched*-mediated regulation of gene expression and other physiological effects mediated by that protein. The *patched* gene product, a cell surface protein, is understood to signal through a pathway which causes transcriptional repression of members of the Wnt and Dpp/BMP families of morphogens, proteins which impart positional information. In development of the CNS and patterning of limbs in vertebrates, the introduction of *hedgehog*

relieves (derepresses) this inhibition conferred by *patched*, allowing expression of particular gene programs.

Recently, it has been reported that mutations in the human version of *patched*, a gene first identified in a fruit fly developmental pathway, cause a hereditary skin cancer and may contribute to sporadic skin cancers. See, for example, Hahn et al. (1996) *Cell* 86:841-851; and Johnson et al. (1996) *Science* 272:1668-1671. The demonstration that nevoid basal-cell carcinoma (NBCC) results from mutations in the human *patched* gene provided an example of the roles *patched* plays in post-embryonic development. These observations have led the art to understand one activity of *patched* to be a tumor suppressor gene, which may act by inhibiting proliferative signals from *hedgehog*. Our observations set forth below reveal potential new roles for the *hedgehog/patched* pathway in maintenance of peripheral nerve cells. Accordingly, the present invention contemplates the use of other agents which are capable of mimicking the effect of the *hedgehog* protein on *patched* signalling, e.g., as may be identified from the drug screening assays described below.

In still other embodiments, antagonists of the *hedgehog* signaling can be used in the selective ablation of sensory neurons, for example, in the treatment of chronic pain syndromes.

II. Definitions

For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

The term "hedgehog therapeutic" refers to various forms of hedgehog polypeptides, as well as peptidomimetics, which can modulate the proliferation/differentiation state of peripheral nerve cells by, as will be clear from the context of individual examples, mimicking or potentiating (agonizing) or inhibiting (antagonizing) the effects of a naturally-occurring *hedgehog* protein. A *hedgehog* therapeutic which mimics or potentiates the activity of a wild-type hedgehog protein is a "hedgehog agonist". Conversely, a *hedgehog* therapeutic which inhibits the activity of a wild-type hedgehog protein is a "hedgehog antagonist".

In particular, the term "hedgehog polypeptide" encompasses preparations of *hedgehog* proteins and peptidyl fragments thereof, both agonist and antagonist forms as the specific context will make clear.

As used herein the term "bioactive fragment of a *hedgehog* protein" refers to a fragment of a full-length *hedgehog* polypeptide, wherein the fragment specifically agonizes or antagonizes inductive events mediated by wild-type *hedgehog* proteins. The *hedgehog* bioactive fragment

preferably is a soluble extracellular portion of a *hedgehog* protein, where solubility is with reference to physiologically compatible solutions. Exemplary bioactive fragments are described in PCT publications WO 95/18856 and WO 96/17924.

The term "ptc therapeutic" refers to agents which either (i) mimic the effect of *hedgehog* proteins on *patched* signalling, e.g., which antagonize the cell-cycle inhibitory activity of *patched*, or (ii) activate or potentiate *patched* signalling. In other embodiments, the ptc therapeutic can be a *hedgehog* antagonist. The ptc therapeutic can be, e.g., a peptide, a nucleic acid, a carbohydrate, a small organic molecule, or natural product extract (or fraction thereof).

An "effective amount" of, e.g., a *hedgehog* therapeutic, with respect to the subject method of treatment, refers to an amount of, e.g., a *hedgehog* polypeptide in a preparation which, when applied as part of a desired dosage regimen brings enhances the survival of peripheral nerves, relative to the absence of the *hedgehog* therapeutic, according to clinically acceptable standards for the disorder to be treated.

A "patient" or "subject" to be treated by the subject method can mean either a human or non-human animal.

The "growth state" of a cell refers to the rate of proliferation of the cell and the state of differentiation of the cell.

"Homology" and "identity" each refer to sequence similarity between two polypeptide sequences, with identity being a more strict comparison. Homology and identity can each be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same amino acid residue, then the polypeptides can be referred to as identical at that position; when the equivalent site is occupied by the same amino acid (e.g., identical) or a similar amino acid (e.g., similar in steric and/or electronic nature), then the molecules can be referred to as homologous at that position. A percentage of homology or identity between sequences is a function of the number of matching or homologous positions shared by the sequences. An "unrelated" or "non-homologous" sequence shares less than 40 percent identity, though preferably less than 25 percent identity, with an *hedgehog* sequence of the present invention.

The term "corresponds to", when referring to a particular polypeptide or nucleic acid sequence is meant to indicate that the sequence of interest is identical or homologous to the reference sequence to which it is said to correspond.

The terms "recombinant protein", "heterologous protein" and "exogenous protein" are used interchangeably throughout the specification and refer to a polypeptide which is produced by recombinant DNA techniques, wherein generally, DNA encoding the polypeptide is inserted

into a suitable expression construct which is in turn used to transform a host cell to produce the heterologous protein. That is, the polypeptide is expressed from a heterologous nucleic acid.

A "chimeric protein" or "fusion protein" is a fusion of a first amino acid sequence encoding a *hedgehog* polypeptide with a second amino acid sequence defining a domain foreign to and not substantially homologous with any domain of *hh* protein. A chimeric protein may present a foreign domain which is found (albeit in a different protein) in an organism which also expresses the first protein, or it may be an "interspecies", "intergenic", etc. fusion of protein structures expressed by different kinds of organisms. In general, a fusion protein can be represented by the general formula $(X)_n-(hh)_m-(Y)_n$, wherein *hh* represents all or a portion of the *hedgehog* protein, X and Y each independently represent an amino acid sequences which are not naturally found as a polypeptide chain contiguous with the *hedgehog* sequence, m is an integer greater than or equal to 1, and each occurrence of n is, independently, 0 or an integer greater than or equal to 1 (n and m are preferably no greater than 5 or 10).

III. Exemplary Applications of Method and Compositions

The subject method has wide applicability to the treatment or prophylaxis of disorders affecting the regulation of peripheral nerves, including peripheral ganglionic neurons, sympathetic, sensory neurons, and motor neurons. In general, the method can be characterized as including a step of administering to an animal an amount of a *ptc* or *hedgehog* therapeutic effective to alter the proliferative and/or differentiation state of treated peripheral nerve cells. Such therapeutic compositions may be useful in treatments designed to rescue, for example, retinal ganglia, inner ear and accoustical nerves, and motoneurons, from lesion-induced death as well as guiding reprojection of these neurons after such damage. Such diseases and conditions include, but are not limited to, chemical or mechanical trauma, infection (such as viral infection with varicella-zoster), metabolic disease such as diabetes, nutritional deficiency, toxic agents (such as cisplatin treatment). The goals of treatment in each case can be twofold: (1) to eliminate the cause of the disease and (2) to relieve its symptoms.

Peripheral neuropathy is a condition involving nerve-ending damage in the hands and feet. Peripheral neuropathy generally refers to a disorder that affects the peripheral nerves, most often manifested as one or a combination of motor, sensory, sensorimotor, or autonomic neural dysfunction. The wide variety of morphologies exhibited by peripheral neuropathies can each be uniquely attributed to an equally wide variety of causes. For instance, peripheral neuropathies can be genetically acquired, can result from a systemic disease, or can be induced by a toxic

agent. Some toxic agents that cause neurotoxicities are therapeutic drugs, antineoplastic agents, contaminants in foods or medicinals, and environmental and industrial pollutants.

In particular, chemotherapeutic agents known to cause sensory and/or motor neuropathies include vincristine, an antineoplastic drug used to treat haematological malignancies and sarcomas, as well as cisplatin, taxol and others. The neurotoxicity is dose-related, and exhibits as reduced intestinal motility and peripheral neuropathy, especially in the distal muscles of the hands and feet, postural hypotension, and atony of the urinary bladder. Similar problems have been documented with taxol and cisplatin (Mollman, J. E., 1990, New Eng Jour Med. 322:126-127), although cisplatin-related neurotoxicity can be alleviated with nerve growth factor (NGF) (Apfel, S. C. et al, 1992, Annals of Neurology 31:76-80). Although the neurotoxicity is sometimes reversible after removal of the neurotoxic agent, recovery can be a very slow process (Legha, S., 1986, Medical Toxicology 1:421-427; Olesen, et al., 1991, Drug Safety 6:302-314).

There are a number of inherited peripheral neuropathies, including: Refsum's disease, Abetalipoproteinemia, Tangier disease, Krabbe's disease, Metachromatic leukodystrophy, Fabry's disease, Dejerine-Sottas syndrome, and others. Of all the inherited neuropathies, the most common by far is Charcot-Marie-Tooth Disease.

Charcot-Marie-Tooth (CMT) Disease (also known as Peroneal Muscular Atrophy, or Hereditary Motor Sensory Neuropathy (HMSN)) is the most common hereditary neurological disorder. It is characterized by weakness and atrophy, primarily of the peroneal muscles, due to segmental demyelination of peripheral nerves and associated degeneration of axons and anterior horn cells. Autosomal dominant inheritance is usual, and associated degenerative CNS disorders, such as Friedreich's ataxia, are common.

In one aspect, the method of the present invention can be used in the treatment and maintenance of hereditary neuropathies. This group of neuropathies are now becoming increasingly recognized due to the dramatic advances in molecular genetics. The symptoms of the various hereditary neuropathies are wide ranging. A common denominator is usually the early onset of mild numbness and tingling in the feet that slowly progresses to involve the legs and the hands and later the rest of the upper extremities. Most of the hereditary neuropathies do have a motor component consisting of distal weakness in the lower and upper extremities. A majority of patients with hereditary neuropathies have high arches in their feet or other bony deformities. The symptoms are very slowly progressive and the majority of the patients are still walking two decades after the onset of their symptoms.

The diagnosis of a hereditary neuropathy is usually suggested with the early onset of neuropathic symptoms, especially when a positive family history is also present. Prior to the

recent genetic advances, the diagnosis was supported by typical findings of marked slowing of the nerve conduction studies on electromyography and a nerve biopsy. Typical findings on a nerve biopsy include the presence of so-called onion-bulbs, indicating a recurring demyelinating and remyelinating of the nerve fibers. With the most recent genetic advances, two major hereditary neuropathies known as "Charcot-Marie-Tooth disease" and "hereditary neuropathy with liability to pressure palsies" can be diagnosed with a simple blood test that identifies the different mutations responsible for these two entities.

Hereditary neuropathies are caused by genetic abnormalities which are transmitted from generation to generation. For several of these, the genetic defect is known, and tests are available for diagnosis and prenatal counseling.

As set forth above, the subject method can be used as part of a therapeutic regimen in the treatment of Charcot-Marie Tooth Disease (CMT). This is a general term given to the hereditary sensorimotor neuropathies. CMT type 1 (CMT 1) is associated with demyelination or breakdown of the myelin sheaths. Several different abnormalities have been identified. CMT Type 1A is most commonly caused by duplication of a gene encoding a myelin protein called PMP-22, and CMT type 1B is caused by a mutation in a myelin protein called the Po glycoprotein. CMTX is a hereditary sensorimotor neuropathy which affects only men. It is caused by a mutation in a gene encoding a protein called Connexin 32 on the X-chromosome.

In certain embodiments, the subject method can be used to treat, or at least reduce the severity of, Amyotrophic lateral sclerosis (ALS). According to the subject invention, a trophic amount of a *hedgehog* or *ptc* therapeutic can be administered to an animal suffering from, or at risk of developing, ALS.

In another embodiment, the subject method can be used in the treatment of Familial Amyloidotic Neuropathy and other related hereditary neuropathies. Amyloidotic neuropathy usually presents with pain, sensory loss and autonomic dysfunction. It is caused by a mutation in a protein called Transthyretin, resulting in deposition of the protein as amyloid in the peripheral nerves.

The subject method can be used in the treatment of hereditary porphyria, which can have components of peripheral neuropathy.

Still another hereditary neuropathy for which the subject methods can be used for treatment is hereditary sensory neuropathy Type II (HSN II).

The methods and compositions of the present invention can also be used in the treatment and maintenance of acquired neuropathies.

For example, hedgehog and ptc therapeutics can be used to prevent diabetic neuropathies. Diabetes is the most common known cause of neuropathy. It produces symptoms in approximately 10% of people with diabetes. In most cases, the neuropathy is predominantly sensory, with pain and sensory loss in the hands and feet. But some diabetics have mononeuritis or mononeuritis multiplex which causes weakness in one or more nerves, or lumbosacral plexopathy or amyotrophy which causes weakness in the legs.

The instant method can also be used in the treatment of immune-mediated neuropathies. The main function of the immune system is to protect the body against infectious organisms which enter from outside. In some cases, however the immune system turns against the body and causes autoimmune disease. The immune system consists of several types of white blood cells, including T-lymphocytes, which also regulate the immune response; and B-lymphocytes or plasma cells, which secrete specialized proteins called "antibodies" Sometimes, for unknown reasons, the immune system mistakenly attacks parts of the body such as the peripheral nerves. This is "autoimmune" Peripheral Neuropathy. There are several different types, depending on the part of the peripheral nerve which is attacked and the type of the immune reaction. The following are brief descriptions of the neuropathies which are mediated by the immune system.

For instance, a hedgehog or ptc therapeutic can be used to treat Guillain-Barre Syndrome (GBS). An acute neuropathy because it comes on suddenly or rapidly. Guillain-Barre Syndrome can progress to paralysis and respiratory failure within days or weeks after onset. The neuropathy is caused when the immune system destroys the myelin sheaths of the motor and sensory nerves. It is often preceded by infection, vaccination or trauma, and that is thought to be what triggers the autoimmune reaction. The disease is self-limiting, with spontaneous recovery within six to eight weeks. But the recovery is often incomplete.

Other neuropathies which begin acutely, and which can be treated by the method of the present invention, include Acute Motor Neuropathy, Acute Sensory Neuropathy, and Acute Autonomic Neuropathy, in which there is an immune attack against the motor, sensory or autonomic nerves, respectively. The Miller-Fisher Syndrome is another variant in which there is paralysis of eye gaze, incoordination, and unsteady gait .

Still another acquired neuropathy which is may be treated by the subject method is Chronic Inflammatory Demyelinating Polyneuropathy (CIDP). CIDP is thought to be a chronic and more indolent form of the Guillain-Barre Syndrome. The disease progresses either with repeated attacks, called relapses, or in a stepwise or steady fashion. As in GBS, there appears to be destruction of the myelin sheath by antibodies and T-lymphocytes. But since there is no specific test for CIDP, the diagnosis is based on the clinical and laboratory characteristics.

Chronic Polyneuropathies with antibodies to peripheral nerves is still another peripheral neuropathy for which the subject methods can be employed to treat or prevent. In some types of chronic neuropathies, antibodies to specific components of nerve have been identified. These include demyelinating neuropathy associated with antibodies to the Myelin Associated Glycoprotein (MAG), motor neuropathy associated with antibodies to the gangliosides GM1 or GD1a, and sensory neuropathy associated with anti-sulfatide or GD1b ganglioside antibodies. The antibodies in these cases bind to oligosaccharide or sugar like molecules, which are linked to proteins (glycoproteins) or lipids (glycolipids or gangliosides) in the nerves. It is suspected that these antibodies may be responsible for the neuropathies.

The subject method can also be used as part of a therapeutic plan for treating neuropathies associated with vasculitis or inflammation of the blood vessels in peripheral nerves. Neuropathy can also be caused by Vasculitis - an inflammation of the blood vessels in peripheral nerve. It produces small "strokes" along the course of the peripheral nerves, and may be restricted to the nerves or it may be generalized, include a skin rash, or involve other organs. Several rheumatological diseases like Rheumatoid Arthritis, Lupus, Periarteritis Nodosa, or Sjogren's Syndrome, are associated with generalized Vasculitis, which can also involve the peripheral nerves. Vasculitis can cause Polyneuritis, Mononeuritis, or Mononeuritis Multiplex, depending on the distribution and severity of the lesions.

In still another embodiment, the method of the present invention can be used for treatment of brachial or lumbosacral plexitis. The brachial plexus, which lies under the armpit, contains the nerves to the arm and hand. Brachial Plexitis is the result of inflammation of that nerve bundle, and produces weakness and pain in one or both arms. Lumbosacral Plexitis, which occurs in the pelvis, causes weakness and pain in the legs.

Hedgehog and ptc therapeutics mayu also be suitable for use in the treatment of neuropathies associated with monoclonal gammopathies. In Monoclonal Gammopathy, single clones of B-cells or plasma cells in the bone marrow or lymphoid organs expand to form benign or malignant tumors and secrete antibodies. "Monoclonal" is because there are single clones of antibodies. And "Gammopathy" stands for gammaglobulins, which is another name for antibodies. In some cases, the antibodies react with nerve components; in others, fragments of the antibodies form amyloid deposits.

Yet another aspect of the present invention relates to the use of the subject method in the treatment of neuropathies associated with tumors or neoplasms. Neuropathy can be due to direct infiltration of nerves by tumor cells or to indirect effect of the tumor. The latter is called Paraneoplastic Neuropathy. Several types have been described. For instance, the subject methods can be used to manage sensory neuropathy associated with lung cancer. This

neuropathy is associated with antibodies to a protein called Hu, which is present in the sensory neurons of the peripheral nerves. Likewise, the subject method can be used to treat neuropathies associated with multiple myeloma. Multiple myeloma is a bony tumor which is caused by antibody-secreting plasma cells in the bone marrow. The tumor is made up of a single clone of plasma cells, and the antibodies they produce are identical or monoclonal. Some people with multiple myeloma develop a Sensorimotor Polyneuropathy with degeneration of axons in the peripheral nerves. In other embodiments, the subject method can be used to treat neuropathies associated with Waldenstrom's Macroglobulemia, Chronic Lymphocytic Leukemia, or B-cell Lymphoma. These are tumors caused by antibody-secreting B-lymphocytes in the spleen, bone marrow or lymph nodes. These antibodies are monoclonal and frequently react with peripheral nerve components such as MAG, GM1, or sulfatide. In still other embodiments, the the hedgehog and ptc therapeutics of the present invention can be used as part of therapeutic protocol for the treatment of patients with cancers where neuropathy is a consequence of local irradiation or be caused by medications such as vincristine and cisplatinum.

The present invention also contemplates the use of hedgehog and ptc therapeutics for the treatment of neuropathies associated with amyloidosis. Amyloid is a substance which is deposited in the peripheral nerves and interferes with their operation: the disorder is Amyloidosis. There are two main types: Primary Amyloidosis, in which the deposits contain fragments of monoclonal antibodies (see the Monoclonal Gammopathy paragraph above); and Hereditary Amyloidosis in which the deposits contain a mutated protein called Transthyretin. Primary Amyloidosis is usually associated with Monoclonal Gammopathies or myeloma (See above.)

Still another aspect of the present invention provides the subject method as a means for treating neuropathies caused by infections. Peripheral neuropathies can be caused by infection of the peripheral nerves. Viruses that cause peripheral neuropathies include the AIDS virus, HIV-I, which causes slowly progressive sensory neuropathy, Cytomegalo virus which causes a rapidly progressive paralytic neuropathy, Herpes Zoster which cause Shingles, and Poliovirus which causes a motor neuropathy. Hepatitis B or C infections are sometimes associated with vasculitic neuropathy.

Bacterial infections that cause neuropathy include Leprosy which causes a patchy sensory neuropathy, and Diphtheria which can cause a rapidly progressive paralytic neuropathy. Other infectious diseases that cause neuropathy include Lyme disease which is caused by a spirochete, and Trypanosomiasis which is caused by a parasite. Both commonly present with a multifocal neuropathy

Neuropathies caused by nutritional imbalance are also candidate disorders for treatment by the subject method. Deficiencies of Vitamins B12, B1 (thiamine), B6 (pyridoxine), or E, for example, can produce polyneuropathies with degeneration of peripheral nerve axons. This can be due to poor diet, or inability to absorb the nutrients from the stomach or gut.

5 Moreover megadoses of Vitamin B6 can also cause a peripheral neuropathy, and the subject method can be used as part of a de-toxification program in such cases.

Yet another use of the subject method is in the treatment of neuropathies arising in kidney diseases. Chronic renal failure can cause a predominantly sensory peripheral neuropathy with degeneration of peripheral nerve axons.

10 Another aspect of the present invention provides a method for treating hypothyroid neuropathies. Hypothyroidism is sometimes associated with a painful sensory polyneuropathy with axonal degeneration. Mononeuropathy or Mononeuropathy Multiplex can also occur due to compression of the peripheral nerves by swollen tissues.

15 The subject method can also be used in the treatment of neuropathies caused by Alcohol and Toxins. Certain toxins can cause Peripheral Neuropathy. Lead toxicity is associated with a motor neuropathy; arsenic or mercury cause a sensory neuropathy, Thalium can cause a sensory and autonomic neuropathy. several of the organic solvents and insecticides can also cause polyneuropathy. Alcohol is directly toxic to nerves and alcohol abuse is a major cause of neuropathy. The subject method can be used, in certain embodiments, as part of a broader detoxification program.

20 In still another embodiment, the methods and compositions of the present invention can be used for the treatment of neuropathies caused by drugs. Several drugs are known to cause neuropathy. They include, among others, vincristine and cisplatin in cancer, nitrofurantoin, which is used in pyelonephritis, amiodarone in cardiac arrhythmias, disulfiram in alcoholism, 25 ddC and ddI in AIDS, and dapsone which is used to treat Leprosy. As above, the subject method can be used, in certain embodiments, as part of a broader detoxification program.

30 The method of the present invention can also be used in the treatment of neuropathies caused by trauma or compression. Localized neuropathies can result from compression of nerves by external pressure or overlying tendons and other tissues. The best known of these are the Carpal Tunnel Syndrome which results from compression at the wrist, and cervical or lumbar radiculopathies (Sciatica) which result from compression of nerve roots as they exit the spine. Other common areas of nerve compression include the elbows, armpits, and the back of the knees.

The subject method is also useful in variety of idiopathic neuropathies. The term "idiopathic" is used whenever the cause of the neuropathy cannot be found. In these cases, the neuropathy is classified according to its manifestations, i.e., sensory, motor, or sensorimotor idiopathic polyneuropathy.

Another aspect of the invention provides a conjoint therapy wherein one or more other therapeutic agents are administered with the hedgehog or ptc therapeutic agent. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment. For example, the subject method can be carried out conjointly with other neuroprotective agents. The dosages recited herein would be adjusted to compensate for such additional components in the therapeutic composition. Progress of the treated patient can be monitored by conventional methods.

IV. Exemplary hedgehog therapeutic compounds.

The *hedgehog* therapeutic compositions of the subject method can be generated by any of a variety of techniques, including purification of naturally occurring proteins, recombinantly produced proteins and synthetic chemistry. Polypeptide forms of the hedgehog therapeutics are preferably derived from vertebrate hedgehog proteins, e.g., have sequences corresponding to naturally occurring hedgehog proteins, or fragments thereof, from vertebrate organisms. However, it will be appreciated that the hedgehog polypeptide can correspond to a hedgehog protein (or fragment thereof) which occurs in any metazoan organism.

The various naturally-occurring *hedgehog* proteins from which the subject therapeutics can be derived are characterized by a signal peptide, a highly conserved N-terminal region, and a more divergent C-terminal domain. In addition to signal sequence cleavage in the secretory pathway (Lee, J.J. *et al.* (1992) *Cell* 71:33-50; Tabata, T. *et al.* (1992) *Genes Dev.* 2635-2645; Chang, D.E. *et al.* (1994) *Development* 120:3339-3353), *hedgehog* precursor proteins naturally undergo an internal autoproteolytic cleavage which depends on conserved sequences in the C-terminal portion (Lee *et al.* (1994) *Science* 266:1528-1537; Porter *et al.* (1995) *Nature* 374:363-366). This autocleavage leads to a 19 kD N-terminal peptide and a C-terminal peptide of 26-28 kD (Lee *et al.* (1992) *supra*; Tabata *et al.* (1992) *supra*; Chang *et al.* (1994) *supra*; Lee *et al.* (1994) *supra*; Bumcrot, D.A., *et al.* (1995) *Mol. Cell. Biol.* 15:2294-2303; Porter *et al.* (1995) *supra*; Ekker, S.C. *et al.* (1995) *Curr. Biol.* 5:944-955; Lai, C.J. *et al.* (1995) *Development* 121:2349-2360). The N-terminal peptide stays tightly associated with the surface of cells in which it was synthesized, while the C-terminal peptide is freely diffusible both *in vitro* and *in vivo* (Lee *et al.* (1994) *supra*; Bumcrot *et al.* (1995) *supra*; Mart', E. *et al.* (1995) *Development*

121:2537-2547; Roelink, H. *et al.* (1995) *Cell* 81:445-455). Cell surface retention of the N-terminal peptide is dependent on autocleavage, as a truncated form of *hedgehog* encoded by an RNA which terminates precisely at the normal position of internal cleavage is diffusible *in vitro* (Porter *et al.* (1995) *supra*) and *in vivo* (Porter, J.A. *et al.* (1996) *Cell* 86, 21-34). Biochemical studies have shown that the autoproteolytic cleavage of the *hedgehog* precursor protein proceeds through an internal thioester intermediate which subsequently is cleaved in a nucleophilic substitution. It is suggested that the nucleophile is a small lipophilic molecule, more particularly cholesterol, which becomes covalently bound to the C-terminal end of the N-peptide (Porter *et al.* (1996) *supra*), tethering it to the cell surface.

The vertebrate family of *hedgehog* genes includes at least four members, e.g., paralogs of the single drosophila *hedgehog* gene (SEQ ID No. 19). Three of these members, herein referred to as Desert *hedgehog* (*Dhh*), Sonic *hedgehog* (*Shh*) and Indian *hedgehog* (*Ihh*), apparently exist in all vertebrates, including fish, birds, and mammals. A fourth member, herein referred to as tiggie-winkle *hedgehog* (*Thh*), appears specific to fish. According to the appended sequence listing, (see also Table 1) a chicken *Shh* polypeptide is encoded by SEQ ID No:1; a mouse *Dhh* polypeptide is encoded by SEQ ID No:2; a mouse *Ihh* polypeptide is encoded by SEQ ID No:3; a mouse *Shh* polypeptide is encoded by SEQ ID No:4 a zebrafish *Shh* polypeptide is encoded by SEQ ID No:5; a human *Shh* polypeptide is encoded by SEQ ID No:6; a human *Ihh* polypeptide is encoded by SEQ ID No:7; a human *Dhh* polypeptide is encoded by SEQ ID No. 8; and a zebrafish *Thh* is encoded by SEQ ID No. 9.

Table 1
Guide to *hedgehog* sequences in Sequence Listing

	Nucleotide	Amino Acid
Chicken <i>Shh</i>	SEQ ID No. 1	SEQ ID No. 10
Mouse <i>Dhh</i>	SEQ ID No. 2	SEQ ID No. 11
Mouse <i>Ihh</i>	SEQ ID No. 3	SEQ ID No. 12
Mouse <i>Shh</i>	SEQ ID No. 4	SEQ ID No. 13
Zebrafish <i>Shh</i>	SEQ ID No. 5	SEQ ID No. 14
Human <i>Shh</i>	SEQ ID No. 6	SEQ ID No. 15
Human <i>Ihh</i>	SEQ ID No. 7	SEQ ID No. 16
Human <i>Dhh</i>	SEQ ID No. 8	SEQ ID No. 17
Zebrafish <i>Thh</i>	SEQ ID No. 9	SEQ ID No. 18
Drosophila <i>HH</i>	SEQ ID No. 19	SEQ ID No. 20

In addition to the sequence variation between the various *hedgehog* homologs, the *hedgehog* proteins are apparently present naturally in a number of different forms, including a

pro-form, a full-length mature form, and several processed fragments thereof. The pro-form includes an N-terminal signal peptide for directed secretion of the extracellular domain, while the full-length mature form lacks this signal sequence.

As described above, further processing of the mature form occurs in some instances to yield biologically active fragments of the protein. For instance, *sonic hedgehog* undergoes additional proteolytic processing to yield two peptides of approximately 19 kDa and 27 kDa, the 19kDa fragment corresponding to an proteolytic N-terminal portion of the mature protein.

In addition to proteolytic fragmentation, the vertebrate *hedgehog* proteins can also be modified post-translationally, such as by glycosylation and/or addition of lipophilic moieties, such as stents, fatty acids, etc., though bacterially produced (e.g. unmodified) forms of the proteins still maintain certain of the bioactivities of the native protein. Bioactive fragments of *hedgehog* polypeptides of the present invention have been generated and are described in great detail in, e.g., PCT publications WO 95/18856 and WO 96/17924.

There are a wide range of lipophilic moieties with which hedgehog polypeptides can be derivatived. The term "lipophilic group", in the context of being attached to a hedgehog polypeptide, refers to a group having high hydrocarbon content thereby giving the group high affinity to lipid phases. A lipophilic group can be, for example, a relatively long chain alkyl or cycloalkyl (preferably n-alkyl) group having approximately 7 to 30 carbons. The alkyl group may terminate with a hydroxy or primary amine "tail". To further illustrate, lipophilic molecules include naturally-occurring and synthetic aromatic and non-aromatic moieties such as fatty acids, sterols, esters and alcohols, other lipid molecules, cage structures such as adamantane and buckminsterfullerenes, and aromatic hydrocarbons such as benzene, perylene, phenanthrene, anthracene, naphthalene, pyrene, chrysene, and naphthacene.

In one embodiment, the hedgehog polypeptide is modified with one or more sterol moieties, such as cholesterol. See, for example, PCT publication WO 96/17924. In certain embodiments, the cholesterol is preferably added to the C-terminal glycine were the hedgehog polypeptide corresponds to the naturally-occurring N-terminal proteolytic fragment.

In another embodiment, the hedgehog polypeptide can be modified with a fatty acid moiety, such as a myristoyl, palmitoyl, stearyl, or arachidoyl moiety. See, e.g., Pepinsky et al. (1998) J Biol. Chem 273: 14037.

In addition to those effects seen by cholesterol-addition to the C-terminus or fatty acid addition to the N-terminus of extracellular fragments of the protein, at least certain of the biological activities of the hedgehog gene products are unexpectedly potentiated by derivatization of the protein with lipophilic moieties at other sites on the protein and/or by

moieties other than cholesterol or fatty acids. Certain aspects of the invention are directed to the use of preparations of hedgehog polypeptides which are modified at sites other than N-terminal or C-terminal residues of the natural processed form of the protein, and/or which are modified at such terminal residues with lipophilic moieties other than a sterol at the C-terminus or fatty acid at the N-terminus.

Particularly useful as lipophilic molecules are alicyclic hydrocarbons, saturated and unsaturated fatty acids and other lipid and phospholipid moieties, waxes, cholesterol, isoprenoids, terpenes and polyalicyclic hydrocarbons including adamantane and buckminsterfullerenes, vitamins, polyethylene glycol or oligoethylene glycol, (C1-C18)-alkyl phosphate diesters, -O-CH₂-CH(OH)-O-(C12-C18)-alkyl, and in particular conjugates with pyrene derivatives. The lipophilic moiety can be a lipophilic dye suitable for use in the invention include, but are not limited to, diphenylhexatriene, Nile Red, N-phenyl-1-naphthylamine, Prodan, Laurodan, Pyrene, Perylene, rhodamine, rhodamine B, tetramethylrhodamine, Texas Red, sulforhodamine, 1,1'-didodecyl-3,3',3'tetramethylindocarbocyanine perchlorate, octadecyl rhodamine B and the BODIPY dyes available from Molecular Probes Inc.

Other exemplary lipophilic moieties include aliphatic carbonyl radical groups include 1- or 2-adamantylacetyl, 3-methyladamant-1-ylacetyl, 3-methyl-3-bromo-1-adamantylacetyl, 1-decalinacetyl, camphoracetyl, camphaneacetyl, noradamantylacetyl, norbornaneacetyl, bicyclo[2.2.2.]-oct-5-eneacetyl, 1-methoxybicyclo[2.2.2.]-oct-5-ene-2-carbonyl, cis-5-norbornene-endo-2,3-dicarbonyl, 5-norbornen-2-ylacetyl, (1R)-(-)-myrtenaneacetyl, 2-norbornaneacetyl, anti-3-oxo-tricyclo[2.2.1.0<2,6>]-heptane-7-carbonyl, decanoyl, dodecanoyl, dodecenoyl, tetradecadienoyl, decynoyl or dodecynoyl.

The hedgehog polypeptide can be linked to the hydrophobic moiety in a number of ways including by chemical coupling means, or by genetic engineering.

There are a large number of chemical cross-linking agents that are known to those skilled in the art. For the present invention, the preferred cross-linking agents are heterobifunctional cross-linkers, which can be used to link the hedgehog polypeptide and hydrophobic moiety in a stepwise manner. Heterobifunctional cross-linkers provide the ability to design more specific coupling methods for conjugating to proteins, thereby reducing the occurrences of unwanted side reactions such as homo-protein polymers. A wide variety of heterobifunctional cross-linkers are known in the art. These include: succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), m-Maleimidobenzoyl-N-hydroxysuccinimide ester (MBS); N-succinimidyl (4-iodoacetyl) aminobenzoate (SIAB), succinimidyl 4-(p-maleimidophenyl) butyrate (SMPB), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC); 4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)-toluene (SMPT), N-succinimidyl 3-(2-

pyridyldithio) propionate (SPDP), succinimidyl 6-[3-(2-pyridyldithio) propionate] hexanoate (LC-SPDP). Those cross-linking agents having N-hydroxysuccinimide moieties can be obtained as the N-hydroxysulfosuccinimide analogs, which generally have greater water solubility. In addition, those cross-linking agents having disulfide bridges within the linking chain can be synthesized instead as the alkyl derivatives so as to reduce the amount of linker cleavage *in vivo*.

In addition to the heterobifunctional cross-linkers, there exists a number of other cross-linking agents including homobifunctional and photoreactive cross-linkers. Disuccinimidyl suberate (DSS), bismaleimido-hexane (BMH) and dimethylpimelimidate-2 HCl (DMP) are examples of useful homobifunctional cross-linking agents, and bis-[β -(4-azidosalicylamido)ethyl]disulfide (BASED) and N-succinimidyl-6(4'-azido-2'-nitrophenyl-amino)hexanoate (SANPAH) are examples of useful photoreactive cross-linkers for use in this invention. For a recent review of protein coupling techniques, see Means et al. (1990) *Bioconjugate Chemistry* 1:2-12, incorporated by reference herein.

One particularly useful class of heterobifunctional cross-linkers, included above, contain the primary amine reactive group, N-hydroxysuccinimide (NHS), or its water soluble analog N-hydroxysulfosuccinimide (sulfo-NHS). Primary amines (lysine epsilon groups) at alkaline pH's are unprotonated and react by nucleophilic attack on NHS or sulfo-NHS esters. This reaction results in the formation of an amide bond, and release of NHS or sulfo-NHS as a by-product.

Another reactive group useful as part of a heterobifunctional cross-linker is a thiol reactive group. Common thiol reactive groups include maleimides, halogens, and pyridyl disulfides. Maleimides react specifically with free sulfhydryls (cysteine residues) in minutes, under slightly acidic to neutral (pH 6.5-7.5) conditions. Halogens (iodoacetyl functions) react with -SH groups at physiological pH's. Both of these reactive groups result in the formation of stable thioether bonds.

The third component of the heterobifunctional cross-linker is the spacer arm or bridge. The bridge is the structure that connects the two reactive ends. The most apparent attribute of the bridge is its effect on steric hindrance. In some instances, a longer bridge can more easily span the distance necessary to link two complex biomolecules. For instance, SMPB has a span of 14.5 angstroms.

Preparing protein-protein conjugates using heterobifunctional reagents is a two-step process involving the amine reaction and the sulfhydryl reaction. For the first step, the amine reaction, the protein chosen should contain a primary amine. This can be lysine epsilon amines or a primary alpha amine found at the N-terminus of most proteins. The protein should not contain free sulfhydryl groups. In cases where both proteins to be conjugated contain free

sulfhydryl groups, one protein can be modified so that all sulfhydryls are blocked using for instance, N-ethylmaleimide (see Partis et al. (1983) J. Pro. Chem. 2:263, incorporated by reference herein). Ellman's Reagent can be used to calculate the quantity of sulfhydryls in a particular protein (see for example Ellman et al. (1958) Arch. Biochem. Biophys. 74:443 and Riddles et al. (1979) Anal. Biochem. 94:75, incorporated by reference herein).

The reaction buffer should be free of extraneous amines and sulfhydryls. The pH of the reaction buffer should be 7.0-7.5. This pH range prevents maleimide groups from reacting with amines, preserving the maleimide group for the second reaction with sulfhydryls.

The NHS-ester containing cross-linkers have limited water solubility. They should be dissolved in a minimal amount of organic solvent (DMF or DMSO) before introducing the cross-linker into the reaction mixture. The cross-linker/solvent forms an emulsion which will allow the reaction to occur.

The sulfo-NHS ester analogs are more water soluble, and can be added directly to the reaction buffer. Buffers of high ionic strength should be avoided, as they have a tendency to "salt out" the sulfo-NHS esters. To avoid loss of reactivity due to hydrolysis, the cross-linker is added to the reaction mixture immediately after dissolving the protein solution.

The reactions can be more efficient in concentrated protein solutions. The more alkaline the pH of the reaction mixture, the faster the rate of reaction. The rate of hydrolysis of the NHS and sulfo-NHS esters will also increase with increasing pH. Higher temperatures will increase the reaction rates for both hydrolysis and acylation.

Once the reaction is completed, the first protein is now activated, with a sulfhydryl reactive moiety. The activated protein may be isolated from the reaction mixture by simple gel filtration or dialysis. To carry out the second step of the cross-linking, the sulfhydryl reaction, the lipophilic group chosen for reaction with maleimides, activated halogens, or pyridyl disulfides must contain a free sulfhydryl. Alternatively, a primary amine may be modified with to add a sulfhydryl

In all cases, the buffer should be degassed to prevent oxidation of sulfhydryl groups. EDTA may be added to chelate any oxidizing metals that may be present in the buffer. Buffers should be free of any sulfhydryl containing compounds.

Maleimides react specifically with -SH groups at slightly acidic to neutral pH ranges (6.5-7.5). A neutral pH is sufficient for reactions involving halogens and pyridyl disulfides. Under these conditions, maleimides generally react with -SH groups within a matter of minutes. Longer reaction times are required for halogens and pyridyl disulfides.

The first sulfhydryl reactive-protein prepared in the amine reaction step is mixed with the sulfhydryl-containing lipophilic group under the appropriate buffer conditions. The conjugates can be isolated from the reaction mixture by methods such as gel filtration or by dialysis.

Exemplary activated lipophilic moieties for conjugation include: N-(1-pyrene)maleimide; 2,5-dimethoxystilbene-4'-maleimide, eosin-5-maleimide; fluorescein-5-maleimide; N-(4-(6-dimethylamino-2-benzofuranyl)phenyl)maleimide; benzophenone-4-maleimide; 4-dimethylaminophenylazophenyl-4'-maleimide (DABMI), tetramethylrhodamine-5-maleimide, tetramethylrhodamine-6-maleimide, Rhodamine Red™ C2 maleimide, N-(5-aminopentyl)maleimide, trifluoroacetic acid salt, N-(2-aminoethyl)maleimide, trifluoroacetic acid salt, Oregon Green™ 488 maleimide, N-(2-((2-(((4-azido-2,3,5,6-tetrafluoro)benzoyl)amino)ethyl)dithio)ethyl)maleimide (TFPAM-SS1), 2-(1-(3-dimethylaminopropyl)-indol-3-yl)-3-(indol-3-yl) maleimide (bisindolylmaleimide; GF 109203X), BODIPY® FL N-(2-aminoethyl)maleimide, N-(7-dimethylamino-4-methylcoumarin-3-yl)maleimide (DACM), Alexa™ 488 C5 maleimide, Alexa™ 594 C5 maleimide, sodium salt N-(1-pyrene)maleimide, 2,5-dimethoxystilbene-4'-maleimide, eosin-5-maleimide, fluorescein-5-maleimide, N-(4-(6-dimethylamino-2-benzofuranyl)phenyl)maleimide, benzophenone-4-maleimide, 4-dimethylaminophenylazophenyl-4'-maleimide, 1-(2-maleimidylethyl)-4-(5-(4-methoxyphenyl)oxazol-2-yl)pyridinium methanesulfonate, tetramethylrhodamine-5-maleimide, tetramethylrhodamine-6-maleimide, Rhodamine Red™ C2 maleimide, N-(5-aminopentyl)maleimide, N-(2-aminoethyl)maleimide, N-(2-((2-(((4-azido-2,3,5,6-tetrafluoro)benzoyl)amino)ethyl)dithio)ethyl)maleimide, 2-(1-(3-dimethylaminopropyl)-indol-3-yl)-3-(indol-3-yl) maleimide, N-(7-dimethylamino-4-methylcoumarin-3-yl)maleimide (DACM), 11H-Benzo[a]fluorene, Benzo[a]pyrene.

In one embodiment, the hedgehog polypeptide can be derivatized using pyrene maleimide, which can be purchased from Molecular Probes (Eugene, Oreg.), e.g., N-(1-pyrene)maleimide or 1-pyrenemethyl iodoacetate (PMIA ester).

For those embodiments wherein the hydrophobic moiety is a polypeptide, the modified hedgehog polypeptide of this invention can be constructed as a fusion protein, containing the hedgehog polypeptide and the hydrophobic moiety as one contiguous polypeptide chain.

In certain embodiments, the lipophilic moiety is an amphipathic polypeptide, such as magainin, cecropin, attacin, melittin, gramicidin S, alpha-toxin of Staph. aureus, alamethicin or a synthetic amphipathic polypeptide. Fusogenic coat proteins from viral particles can also be a convenient source of amphipathic sequences for the subject hedgehog proteins

Moreover, mutagenesis can be used to create modified *hh* polypeptides, e.g., for such purposes as enhancing therapeutic or prophylactic efficacy, or stability (e.g., *ex vivo* shelf life and resistance to proteolytic degradation *in vivo*). Such modified peptides can be produced, for instance, by amino acid substitution, deletion, or addition. Modified *hedgehog* polypeptides can also include those with altered post-translational processing relative to a naturally occurring *hedgehog* protein, e.g., altered glycosylation, cholesterolization, prenylation and the like.

In one embodiment, the *hedgehog* therapeutic is a polypeptide encodable by a nucleotide sequence that hybridizes under stringent conditions to a *hedgehog* coding sequence represented in one or more of SEQ ID Nos:1-7. Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C, are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50°C to a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C.

As described in the literature, genes for other *hedgehog* proteins, e.g., from other animals, can be obtained from mRNA or genomic DNA samples using techniques well known in the art. For example, a cDNA encoding a *hedgehog* protein can be obtained by isolating total mRNA from a cell, e.g. a mammalian cell, e.g. a human cell, including embryonic cells. Double stranded cDNAs can then be prepared from the total mRNA, and subsequently inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. The gene encoding a *hedgehog* protein can also be cloned using established polymerase chain reaction techniques.

Preferred nucleic acids encode a *hedgehog* polypeptide comprising an amino acid sequence at least 60% homologous or identical, more preferably 70% homologous or identical, and most preferably 80% homologous or identical with an amino acid sequence selected from the group consisting of SEQ ID Nos:8-14. Nucleic acids which encode polypeptides at least about 90%, more preferably at least about 95%, and most preferably at least about 98-99% homology or identity with an amino acid sequence represented in one of SEQ ID Nos:8-14 are also within the scope of the invention.

In addition to native *hedgehog* proteins, *hedgehog* polypeptides preferred by the present invention are at least 60% homologous or identical, more preferably 70% homologous or identical and most preferably 80% homologous or identical with an amino acid sequence represented by any of SEQ ID Nos:8-14. Polypeptides which are at least 90%, more preferably

at least 95%, and most preferably at least about 98-99% homologous or identical with a sequence selected from the group consisting of SEQ ID Nos:8-14 are also within the scope of the invention. The only prerequisite is that the *hedgehog* polypeptide is capable of modulating the growth state of peripheral nerve cells.

5 The term "recombinant protein" refers to a polypeptide of the present invention which is produced by recombinant DNA techniques, wherein generally, DNA encoding a *hedgehog* polypeptide is inserted into a suitable expression vector which is in turn used to transform a host cell to produce the heterologous protein. Moreover, the phrase "derived from", with respect to
10 those proteins having an amino acid sequence of a native *hedgehog* protein, or an amino acid sequence similar thereto which is generated by mutations including substitutions and deletions (including truncation) of a naturally occurring form of the protein.

The method of the present invention can also be carried out using variant forms of the naturally occurring *hedgehog* polypeptides, e.g., mutational variants.

15 As is known in the art, hedgehog polypeptides can be produced by standard biological techniques or by chemical synthesis. For example, a host cell transfected with a nucleic acid vector directing expression of a nucleotide sequence encoding the subject polypeptides can be cultured under appropriate conditions to allow expression of the peptide to occur. The polypeptide *hedgehog* may be secreted and isolated from a mixture of cells and medium containing the recombinant *hedgehog* polypeptide. Alternatively, the peptide may be retained
20 cytoplasmically by removing the signal peptide sequence from the recombinant *hedgehog* gene and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. The recombinant *hedgehog* polypeptide can be isolated from cell culture medium, host cells, or both using
25 techniques known in the art for purifying proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies specific for such peptide. In a preferred embodiment, the recombinant *hedgehog* polypeptide is a fusion protein containing a domain which facilitates its purification, such as an *hedgehog*/GST fusion protein. The host cell may be any prokaryotic or eukaryotic cell.

30 Recombinant *hedgehog* genes can be produced by ligating nucleic acid encoding an *hedgehog* protein, or a portion thereof, into a vector suitable for expression in either prokaryotic cells, eukaryotic cells, or both. Expression vectors for production of recombinant forms of the subject *hedgehog* polypeptides include plasmids and other vectors. For instance, suitable vectors for the expression of a *hedgehog* polypeptide include plasmids of the types: pBR322-derived

plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

A number of vectors exist for the expression of recombinant proteins in yeast. For instance, YEP24, YIP5, YEP51, YEP52, pYES2, and YRP17 are cloning and expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae* (see, for example, Broach *et al.* (1983) in *Experimental Manipulation of Gene Expression*, ed. M. Inouye Academic Press, p. 83, incorporated by reference herein). These vectors can replicate in *E. coli* due to the presence of the pBR322 ori, and in *S. cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. In addition, drug resistance markers such as ampicillin can be used. In an illustrative embodiment, an *hedgehog* polypeptide is produced recombinantly utilizing an expression vector generated by sub-cloning the coding sequence of one of the *hedgehog* genes represented in SEQ ID Nos:1-7.

The preferred mammalian expression vectors contain both prokaryotic sequences, to facilitate the propagation of the vector in bacteria, and one or more eukaryotic transcription units that are expressed in eukaryotic cells. The pcDNA1/amp, pcDNA1/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papillomavirus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells. The various methods employed in the preparation of the plasmids and transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989) Chapters 16 and 17.

In some instances, it may be desirable to express the recombinant *hedgehog* polypeptide by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the β -gal containing pBlueBac III).

When it is desirable to express only a portion of an *hedgehog* protein, such as a form lacking a portion of the N-terminus, i.e. a truncation mutant which lacks the signal peptide, it may be necessary to add a start codon (ATG) to the oligonucleotide fragment containing the desired sequence to be expressed. It is well known in the art that a methionine at the N-terminal

position can be enzymatically cleaved by the use of the enzyme methionine aminopeptidase (MAP). MAP has been cloned from *E. coli* (Ben-Bassat et al. (1987) *J. Bacteriol.* 169:751-757) and *Salmonella typhimurium* and its *in vitro* activity has been demonstrated on recombinant proteins (Miller et al. (1987) *PNAS* 84:2718-1722). Therefore, removal of an N-terminal methionine, if desired, can be achieved either *in vivo* by expressing *hedgehog*-derived polypeptides in a host which produces MAP (e.g., *E. coli* or CM89 or *S. cerevisiae*), or *in vitro* by use of purified MAP (e.g., procedure of Miller et al., *supra*).

Alternatively, the coding sequences for the polypeptide can be incorporated as a part of a fusion gene including a nucleotide sequence encoding a different polypeptide. It is widely appreciated that fusion proteins can also facilitate the expression of proteins, and accordingly, can be used in the expression of the *hedgehog* polypeptides of the present invention. For example, *hedgehog* polypeptides can be generated as glutathione-S-transferase (GST-fusion) proteins. Such GST-fusion proteins can enable easy purification of the *hedgehog* polypeptide, as for example by the use of glutathione-derivatized matrices (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. (N.Y.: John Wiley & Sons, 1991)). In another embodiment, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence, can be used to replace the signal sequence which naturally occurs at the N-terminus of the *hedgehog* protein (e.g. of the pro-form, in order to permit purification of the poly(His)-*hedgehog* protein by affinity chromatography using a Ni²⁺ metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase (e.g., see Hochuli et al. (1987) *J. Chromatography* 411:177; and Janknecht et al. *PNAS* 88:8972).

Techniques for making fusion genes are known to those skilled in the art. Essentially, the joining of various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992).

Hedgehog polypeptides may also be chemically modified to create *hedgehog* derivatives by forming covalent or aggregate conjugates with other chemical moieties, such as glycosyl

groups, cholesterol, isoprenoids, lipids, phosphate, acetyl groups and the like. Covalent derivatives of *hedgehog* proteins can be prepared by linking the chemical moieties to functional groups on amino acid sidechains of the protein or at the N-terminus or at the C-terminus of the polypeptide.

5 For instance, *hedgehog* proteins can be generated to include a moiety, other than sequence naturally associated with the protein, that binds a component of the extracellular matrix and enhances localization of the analog to cell surfaces. For example, sequences derived from the fibronectin "type-III repeat", such as a tetrapeptide sequence R-G-D-S (Pierschbacher et al. (1984) *Nature* 309:30-3; and Kornblihtt et al. (1985) *EMBO* 4:1755-9) can be added to the
10 *hedgehog* polypeptide to support attachment of the chimeric molecule to a cell through binding ECM components (Ruoslahti et al. (1987) *Science* 238:491-497; Pierschbacher et al. (1987) *J. Biol. Chem.* 262:17294-8.; Hynes (1987) *Cell* 48:549-54; and Hynes (1992) *Cell* 69:11-25).

In a preferred embodiment, the *hedgehog* polypeptide is isolated from, or is otherwise substantially free of, other cellular proteins, especially other extracellular or cell surface associated proteins which may normally be associated with the *hedgehog* polypeptide, unless provided in the form of fusion protein with the *hedgehog* polypeptide. The term "substantially free of other cellular or extracellular proteins" (also referred to herein as "contaminating proteins") or "substantially pure preparations" or "purified preparations" are defined as encompassing preparations of *hedgehog* polypeptides having less than 20% (by dry weight) contaminating protein, and preferably having less than 5% contaminating protein. By "purified",
15 it is meant that the indicated molecule is present in the substantial absence of other biological macromolecules, such as other proteins. The term "purified" as used herein preferably means at least 80% by dry weight, more preferably in the range of 95-99% by weight, and most preferably at least 99.8% by weight, of biological macromolecules of the same type present (but water,
20 buffers, and other small molecules, especially molecules having a molecular weight of less than 5000, can be present). The term "pure" as used herein preferably has the same numerical limits as "purified" immediately above.

As described above for recombinant polypeptides, isolated *hedgehog* polypeptides can include all or a portion of the amino acid sequences represented in any of SEQ ID Nos:10-18 or
30 20, or a homologous sequence thereto. Preferred fragments of the subject *hedgehog* proteins correspond to the N-terminal and C-terminal proteolytic fragments of the mature protein. Bioactive fragments of *hedgehog* polypeptides are described in great detail in PCT publications WO 95/18856 and WO 96/17924.

With respect to bioactive fragments of *hedgehog* polypeptide, preferred *hedgehog*
35 therapeutics include at least 50 (contiguous) amino acid residues of a *hedgehog* polypeptide,

more preferably at least 100 (contiguous), and even more preferably at least 150 (contiguous) residues.

Another preferred *hedgehog* polypeptide which can be included in the *hedgehog* therapeutic is an N-terminal fragment of the mature protein having a molecular weight of approximately 19 kDa.

Preferred human *hedgehog* proteins include N-terminal fragments corresponding approximately to residues 24-197 of SEQ ID No. 15, 28-202 of SEQ ID No. 16, and 23-198 of SEQ ID No. 17. By "corresponding approximately" it is meant that the sequence of interest is at most 20 amino acid residues different in length to the reference sequence, though more preferably at most 5, 10 or 15 amino acid different in length.

As described above for recombinant polypeptides, isolated *hedgehog* polypeptides can include all or a portion of the amino acid sequences represented in SEQ ID No:8, SEQ ID No:9, SEQ ID No:10, SEQ ID No:11, SEQ ID No:12, SEQ ID No:13 or SEQ ID No:14, or a homologous sequence thereto. Preferred fragments of the subject *hedgehog* proteins correspond to the N-terminal and C-terminal proteolytic fragments of the mature protein. Bioactive fragments of hedgehog polypeptides are described in great detail in PCT publications WO 95/18856 and WO 96/17924.

Still other preferred *hedgehog* polypeptides includes an amino acid sequence represented by the formula A-B wherein: (i) A represents all or the portion of the amino acid sequence designated by residues 1-168 of SEQ ID No:21; and B represents at least one amino acid residue of the amino acid sequence designated by residues 169-221 of SEQ ID No:21; (ii) A represents all or the portion of the amino acid sequence designated by residues 24-193 of SEQ ID No:15; and B represents at least one amino acid residue of the amino acid sequence designated by residues 194-250 of SEQ ID No:15; (iii) A represents all or the portion of the amino acid sequence designated by residues 25-193 of SEQ ID No:13; and B represents at least one amino acid residue of the amino acid sequence designated by residues 194-250 of SEQ ID No:13; (iv) A represents all or the portion of the amino acid sequence designated by residues 23-193 of SEQ ID No:11; and B represents at least one amino acid residue of the amino acid sequence designated by residues 194-250 of SEQ ID No:11; (v) A represents all or the portion of the amino acid sequence designated by residues 28-197 of SEQ ID No:12; and B represents at least one amino acid residue of the amino acid sequence designated by residues 198-250 of SEQ ID No:12; (vi) A represents all or the portion of the amino acid sequence designated by residues 29-197 of SEQ ID No:16; and B represents at least one amino acid residue of the amino acid sequence designated by residues 198-250 of SEQ ID No:16; or (vii) A represents all or the portion of the amino acid sequence designated by residues 23-193 of SEQ ID No. 17, and B

represents at least one amino acid residue of the amino acid sequence designated by residues 194-250 of SEQ ID No. 17. In certain preferred embodiments, A and B together represent a contiguous polypeptide sequence designated sequence, A represents at least 25, 50, 75, 100, 125 or 150 (contiguous) amino acids of the designated sequence, and B represents at least 5, 10, or 20 (contiguous) amino acid residues of the amino acid sequence designated by corresponding entry in the sequence listing, and A and B together preferably represent a contiguous sequence corresponding to the sequence listing entry. Similar fragments from other *hedgehog* also contemplated, e.g., fragments which correspond to the preferred fragments from the sequence listing entries which are enumerated above. In preferred embodiments, the *hedgehog* polypeptide includes a C-terminal glycine (or other appropriate residue) which is derivatized with a cholesterol.

Isolated peptidyl portions of *hedgehog* proteins can be obtained by screening peptides recombinantly produced from the corresponding fragment of the nucleic acid encoding such peptides. In addition, fragments can be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. For example, a *hedgehog* polypeptide of the present invention may be arbitrarily divided into fragments of desired length with no overlap of the fragments, or preferably divided into overlapping fragments of a desired length. The fragments can be produced (recombinantly or by chemical synthesis) and tested to identify those peptidyl fragments which can function as either agonists or antagonists of a wild-type (e.g., "authentic") *hedgehog* protein. For example, Román et al. (1994) *Eur J Biochem* 222:65-73 describe the use of competitive-binding assays using short, overlapping synthetic peptides from larger proteins to identify binding domains.

The recombinant *hedgehog* polypeptides of the present invention also include homologs of the authentic *hedgehog* proteins, such as versions of those protein which are resistant to proteolytic cleavage, as for example, due to mutations which alter potential cleavage sequences or which inactivate an enzymatic activity associated with the protein. *Hedgehog* homologs of the present invention also include proteins which have been post-translationally modified in a manner different than the authentic protein. Exemplary derivatives of *hedgehog* proteins include polypeptides which lack N-glycosylation sites (e.g. to produce an unglycosylated protein), which lack sites for cholesterolization, and/or which lack N-terminal and/or C-terminal sequences.

Modification of the structure of the subject *hedgehog* polypeptides can also be for such purposes as enhancing therapeutic or prophylactic efficacy, or stability (e.g., *ex vivo* shelf life and resistance to proteolytic degradation *in vivo*). Such modified peptides, when designed to retain at least one activity of the naturally-occurring form of the protein, are considered functional equivalents of the *hedgehog* polypeptides described in more detail herein. Such

modified peptides can be produced, for instance, by amino acid substitution, deletion, or addition.

It is well known in the art that one could reasonably expect that certain isolated replacements of amino acids, e.g., replacement of an amino acid residue with another related amino acid (i.e. isosteric and/or isoelectric mutations), can be carried out without major effect on the biological activity of the resulting molecule. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids can be divided into four families: (1) acidic = aspartate, glutamate; (2) basic = lysine, arginine, histidine; (3) nonpolar = alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar = glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids. In similar fashion, the amino acid repertoire can be grouped as (1) acidic = aspartate, glutamate; (2) basic = lysine, arginine, histidine, (3) aliphatic = glycine, alanine, valine, leucine, isoleucine, serine, threonine, with serine and threonine optionally be grouped separately as aliphatic-hydroxyl; (4) aromatic = phenylalanine, tyrosine, tryptophan; (5) amide = asparagine, glutamine; and (6) sulfur-containing = cysteine and methionine. (see, for example, *Biochemistry*, 2nd ed., Ed. by L. Stryer, WH Freeman and Co.: 1981). Whether a change in the amino acid sequence of a peptide results in a functional *hedgehog* homolog (e.g. functional in the sense that it acts to mimic or antagonize the wild-type form) can be readily determined by assessing the ability of the variant peptide to produce a response in cells in a fashion similar to the wild-type protein, or competitively inhibit such a response. Polypeptides in which more than one replacement has taken place can readily be tested in the same manner.

It is specifically contemplated that the methods of the present invention can be carried using homologs of naturally occurring hedgehog proteins. In one embodiment, the invention contemplates using hedgehog polypeptides generated by combinatorial mutagenesis. Such methods, as are known in the art, are convenient for generating both point and truncation mutants, and can be especially useful for identifying potential variant sequences (e.g. homologs) that are functional in binding to a receptor for *hedgehog* proteins. The purpose of screening such combinatorial libraries is to generate, for example, novel *hedgehog* homologs which can act as either agonists or antagonist. To illustrate, *hedgehog* homologs can be engineered by the present method to provide more efficient binding to a cognate receptor, such as *patched*, yet still retain at least a portion of an activity associated with *hedgehog*. Thus, combinatorially-derived homologs can be generated to have an increased potency relative to a naturally occurring form of the protein. Likewise, *hedgehog* homologs can be generated by the present combinatorial approach

to act as antagonists, in that they are able to mimic, for example, binding to other extracellular matrix components (such as receptors), yet not induce any biological response, thereby inhibiting the action of authentic *hedgehog* or *hedgehog* agonists. Moreover, manipulation of certain domains of *hedgehog* by the present method can provide domains more suitable for use in fusion proteins, such as one that incorporates portions of other proteins which are derived from the extracellular matrix and/or which bind extracellular matrix components.

To further illustrate the state of the art of combinatorial mutagenesis, it is noted that the review article of Gallop et al. (1994) *J Med Chem* 37:1233 describes the general state of the art of combinatorial libraries as of the earlier 1990's. In particular, Gallop et al state at page 1239 "[s]creening the analog libraries aids in determining the minimum size of the active sequence and in identifying those residues critical for binding and intolerant of substitution". In addition, the Ladner et al. PCT publication WO90/02809, the Goeddel et al. U.S. Patent 5,223,408, and the Markland et al. PCT publication WO92/15679 illustrate specific techniques which one skilled in the art could utilize to generate libraries of *hedgehog* variants which can be rapidly screened to identify variants/fragments which retained a particular activity of the *hedgehog* polypeptides. These techniques are exemplary of the art and demonstrate that large libraries of related variants/truncants can be generated and assayed to isolate particular variants without undue experimentation. Gustin et al. (1993) *Virology* 193:653, and Bass et al. (1990) *Proteins: Structure, Function and Genetics* 8:309-314 also describe other exemplary techniques from the art which can be adapted as means for generating mutagenic variants of *hedgehog* polypeptides.

Indeed, it is plain from the combinatorial mutagenesis art that large scale mutagenesis of *hedgehog* proteins, without any preconceived ideas of which residues were critical to the biological function, and generate wide arrays of variants having equivalent biological activity. Indeed, it is the ability of combinatorial techniques to screen billions of different variants by high throughput analysis that removes any requirement of *a priori* understanding or knowledge of critical residues.

To illustrate, the amino acid sequences for a population of *hedgehog* homologs or other related proteins are aligned, preferably to promote the highest homology possible. Such a population of variants can include, for example, *hedgehog* homologs from one or more species. Amino acids which appear at each position of the aligned sequences are selected to create a degenerate set of combinatorial sequences. In a preferred embodiment, the variegated library of *hedgehog* variants is generated by combinatorial mutagenesis at the nucleic acid level, and is encoded by a variegated gene library. For instance, a mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential *hedgehog*

sequences are expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g. for phage display) containing the set of *hedgehog* sequences therein.

As illustrated in PCT publication WO 95/18856, to analyze the sequences of a population of variants, the amino acid sequences of interest can be aligned relative to sequence homology.

- 5 The presence or absence of amino acids from an aligned sequence of a particular variant is relative to a chosen consensus length of a reference sequence, which can be real or artificial.

In an illustrative embodiment, alignment of exons 1, 2 and a portion of exon 3 encoded sequences (e.g. the N-terminal approximately 221 residues of the mature protein) of each of the *Shh* clones produces a degenerate set of *Shh* polypeptides represented by the general formula:

10 C-G-P-G-R-G-X(1)-G-X(2)-R-R-H-P-K-K-L-T-P-L-A-Y-K-Q-F-I-P-N-V-A-E-
K-T-L-G-A-S-G-R-Y-E-G-K-I-X(3)-R-N-S-E-R-F-K-E-L-T-P-N-Y-N-P-D-I-I-F-
K-D-E-E-N-T-G-A-D-R-L-M-T-Q-R-C-K-D-K-L-N-X(4)-L-A-I-S-V-M-N-X(5)-
W-P-G-V-X(6)-L-R-V-T-E-G-W-D-E-D-G-H-H-X(7)-E-E-S-L-H-Y-E-G-R-A-
V-D-I-T-T-S-D-R-D-X(8)-S-K-Y-G-X(9)-L-X(10)-R-L-A-V-E-A-G-F-D-W-V-
5 Y-Y-E-S-K-A-H-I-H-C-S-V-K-A-E-N-S-V-A-A-K-S-G-G-C-F-P-G-S-A-X(11)-
V-X(12)-L-X(13)-X(14)-G-G-X(15)-K-X-(16)-V-K-D-L-X(17)-P-G-D-X(18)-V-
L-A-A-D-X(19)-X(20)-G-X(21)-L-X(22)-X(23)-S-D-F-X(24)-X(25)-F-X(26)-D-
R (SEQ ID No: 21)

wherein each of the degenerate positions "X" can be an amino acid which occurs in that position in one of the human, mouse, chicken or zebrafish *Shh* clones, or, to expand the library, each X can also be selected from amongst amino acid residue which would be conservative substitutions for the amino acids which appear naturally in each of those positions. For instance, Xaa(1) represents Gly, Ala, Val, Leu, Ile, Phe, Tyr or Trp ; Xaa(2) represents Arg, His or Lys; Xaa(3) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(4) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(5) represents Lys, Arg, His, Asn or Gln; Xaa(6) represents Lys, Arg or His; Xaa(7) represents Ser, Thr, Tyr, Trp or Phe; Xaa(8) represents Lys, Arg or His; Xaa(9) represents Met, Cys, Ser or Thr; Xaa(10) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(11) represents Leu, Val, Met, Thr or Ser; Xaa(12) represents His, Phe, Tyr, Ser, Thr, Met or Cys; Xaa(13) represents Gln, Asn, Glu, or Asp; Xaa(14) represents His, Phe, Tyr, Thr, Gln, Asn, Glu or Asp; Xaa(15) represents Gln, Asn, Glu, Asp, Thr, Ser, Met or Cys; Xaa(16) represents Ala, Gly, Cys, Leu, Val or Met; Xaa(17) represents Arg, Lys, Met, Ile, Asn, Asp, Glu, Gln, Ser, Thr or Cys; Xaa(18) represents Arg, Lys, Met or Ile; Xaa(19) represents Ala, Gly, Cys, Asp, Glu, Gln, Asn, Ser, Thr or Met; Xaa(20) represents Ala, Gly, Cys, Asp, Asn, Glu or Gln; Xaa(21) represents Arg, Lys, Met, Ile, Asn, Asp, Glu or Gln; Xaa(22) represent Leu, Val, Met or Ile; Xaa(23) represents Phe, Tyr, Thr, His or Trp; Xaa(24) represents Ile, Val, Leu or Met; .Xaa(25) represents Met, Cys, Ile,

Leu, Val, Thr or Ser; Xaa(26) represents Leu, Val, Met, Thr or Ser. In an even more expansive library, each X can be selected from any amino acid.

In similar fashion, alignment of each of the human, mouse, chicken and zebrafish *hedgehog* clones, can provide a degenerate polypeptide sequence represented by the general formula:

C-G-P-G-R-G-X(1)-X(2)-X(3)-R-R-X(4)-X(5)-X(6)-P-K-X(7)-L-X(8)-P-L-X(9)-
Y-K-Q-F-X(10)-P-X(11)-X(12)-X(13)-E-X(14)-T-L-G-A-S-G-X(15)-X(16)-E-G-
X(17)-X(18)-X(19)-R-X(20)-S-E-R-F-X(21)-X(22)-L-T-P-N-Y-N-P-D-I-I-F-K-
D-E-E-N-X(23)-G-A-D-R-L-M-T-X(24)-R-C-K-X(25)-X(26)-X(27)-N-X(28)-L-
A-I-S-V-M-N-X(29)-W-P-G-V-X(30)-L-R-V-T-E-G-X(31)-D-E-D-G-H-H-
X(32)-X(33)-X(34)-S-L-H-Y-E-G-R-A-X(35)-D-I-T-T-S-D-R-D-X(36)-X(37)-K-
Y-G-X(38)-L-X(39)-R-L-A-V-E-A-G-F-D-W-V-Y-Y-E-S-X(40)-X(41)-H-X(42)-
H-X(43)-S-V-K-X(44)-X(45) (SEQ IDNo:22)

wherein, as above, each of the degenerate positions "X" can be an amino acid which occurs in a corresponding position in one of the wild-type clones, and may also include amino acid residue which would be conservative substitutions, or each X can be any amino acid residue. In an exemplary embodiment, Xaa(1) represents Gly, Ala, Val, Leu, Ile, Pro, Phe or Tyr; Xaa(2) represents Gly, Ala, Val, Leu or Ile; Xaa(3) represents Gly, Ala, Val, Leu, Ile, Lys, His or Arg; Xaa(4) represents Lys, Arg or His; Xaa(5) represents Phe, Trp, Tyr or an amino acid gap; Xaa(6) represents Gly, Ala, Val, Leu, Ile or an amino acid gap; Xaa(7) represents Asn, Gln, His, Arg or Lys; Xaa(8) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(9) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(10) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(11) represents Ser, Thr, Gln or Asn; Xaa(12) represents Met, Cys, Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(13) represents Gly, Ala, Val, Leu, Ile or Pro; Xaa(14) represents Arg, His or Lys; Xaa(15) represents Gly, Ala, Val, Leu, Ile, Pro, Arg, His or Lys; Xaa(16) represents Gly, Ala, Val, Leu, Ile, Phe or Tyr; Xaa(17) represents Arg, His or Lys; Xaa(18) represents Gly, Ala, Val, Leu, Ile, Ser or Thr; Xaa(19) represents Thr or Ser; Xaa(20) represents Gly, Ala, Val, Leu, Ile, Asn or Gln; Xaa(21) represents Arg, His or Lys; Xaa(22) represents Asp or Glu; Xaa(23) represents Ser or Thr; Xaa(24) represents Glu, Asp, Gln or Asn; Xaa(25) represents Glu or Asp; Xaa(26) represents Arg, His or Lys; Xaa(27) represents Gly, Ala, Val, Leu or Ile; Xaa(28) represents Gly, Ala, Val, Leu, Ile, Thr or Ser; Xaa(29) represents Met, Cys, Gln, Asn, Arg, Lys or His; Xaa(30) represents Arg, His or Lys; Xaa(31) represents Trp, Phe, Tyr, Arg, His or Lys; Xaa(32) represents Gly, Ala, Val, Leu, Ile, Ser, Thr, Tyr or Phe; Xaa(33) represents Gln, Asn, Asp or Glu; Xaa(34) represents Asp or Glu; Xaa(35) represents Gly, Ala, Val, Leu, or Ile; Xaa(36) represents Arg, His or Lys; Xaa(37) represents Asn, Gln, Thr or Ser; Xaa(38) represents Gly, Ala, Val, Leu, Ile, Ser, Thr, Met or Cys; Xaa(39) represents Gly, Ala, Val, Leu, Ile, Thr or Ser; Xaa(40) represents Arg, His

or Lys; Xaa(41) represents Asn, Gln, Gly, Ala, Val, Leu or Ile; Xaa(42) represents Gly, Ala, Val, Leu or Ile; Xaa(43) represents Gly, Ala, Val, Leu, Ile, Ser, Thr or Cys; Xaa(44) represents Gly, Ala, Val, Leu, Ile, Thr or Ser; and Xaa(45) represents Asp or Glu.

There are many ways by which the library of potential *hedgehog* homologs can be generated from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be carried out in an automatic DNA synthesizer, and the synthetic genes then ligated into an appropriate expression vector. The purpose of a degenerate set of genes is to provide, in one mixture, all of the sequences encoding the desired set of potential *hedgehog* sequences. The synthesis of degenerate oligonucleotides is well known in the art (see for example, Narang, SA (1983) *Tetrahedron* 39:3; Itakura et al. (1981) *Recombinant DNA, Proc 3rd Cleveland Sympos. Macromolecules*, ed. AG Walton, Amsterdam: Elsevier pp273-289; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477. Such techniques have been employed in the directed evolution of other proteins (see, for example, Scott et al. (1990) *Science* 249:386-390; Roberts et al. (1992) *PNAS* 89:2429-2433; Devlin et al. (1990) *Science* 249: 404-406; Cwirla et al. (1990) *PNAS* 87: 6378-6382; as well as U.S. Patents Nos. 5,223,409, 5,198,346, and 5,096,815).

A wide range of techniques are known in the art for screening gene products of combinatorial libraries made by point mutations, and for screening cDNA libraries for gene products having a certain property. Such techniques will be generally adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of *hedgehog* homologs. The most widely used techniques for screening large gene libraries typically comprises cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates relatively easy isolation of the vector encoding the gene whose product was detected. Each of the illustrative assays described below are amenable to high through-put analysis as necessary to screen large numbers of degenerate *hedgehog* sequences created by combinatorial mutagenesis techniques.

In one embodiment, the combinatorial library is designed to be secreted (e.g. the polypeptides of the library all include a signal sequence but no transmembrane or cytoplasmic domains), and is used to transfect a eukaryotic cell that can be co-cultured with peripheal nerve cells. A functional *hedgehog* protein secreted by the cells expressing the combinatorial library will diffuse to neighboring peripheral nerve cells and induce a particular biological response, such as proliferation or differentiation. The pattern of detection of such a change in phenotype will resemble a gradient function, and will allow the isolation (generally after several repetitive rounds of selection) of cells producing *hedgehog* homologs active as neurotrophic agents.

Likewise, *hedgehog* antagonists can be selected in similar fashion by the ability of the cell producing a functional antagonist to protect neighboring cells (e.g., to inhibit proliferation) from the effect of wild-type *hedgehog* added to the culture media.

To illustrate, target peripheral nerve cells are cultured in 24-well microtitre plates. Other eukaryotic cells are transfected with the combinatorial *hedgehog* gene library and cultured in cell culture inserts (e.g. Collaborative Biomedical Products, Catalog #40446) that are able to fit into the wells of the microtitre plate. The cell culture inserts are placed in the wells such that recombinant *hedgehog* homologs secreted by the cells in the insert can diffuse through the porous bottom of the insert and contact the target cells in the microtitre plate wells. After a period of time sufficient for functional forms of a *hedgehog* protein to produce a measurable response in the target cells, such as growth state, the inserts are removed and the effect of the variant *hedgehog* proteins on the target cells determined. Cells from the inserts corresponding to wells which score positive for activity can be split and re-cultured on several inserts, the process being repeated until the active clones are identified.

In yet another screening assay, the candidate *hedgehog* gene products are displayed on the surface of a cell or viral particle, and the ability of particular cells or viral particles to associate with a *hedgehog*-binding moiety (such as the *patched* protein or other *hedgehog* receptor) via this gene product is detected in a "panning assay". Such panning steps can be carried out on cells cultured from embryos. For instance, the gene library can be cloned into the gene for a surface membrane protein of a bacterial cell, and the resulting fusion protein detected by panning (Ladner et al., WO 88/06630; Fuchs et al. (1991) *Bio/Technology* 9:1370-1371; and Goward et al. (1992) *TIBS* 18:136-140). In a similar fashion, fluorescently labeled molecules which bind *hedgehog* can be used to score for potentially functional *hedgehog* homologs. Cells can be visually inspected and separated under a fluorescence microscope, or, where the morphology of the cell permits, separated by a fluorescence-activated cell sorter.

In an alternate embodiment, the gene library is expressed as a fusion protein on the surface of a viral particle. For instance, in the filamentous phage system, foreign peptide sequences can be expressed on the surface of infectious phage, thereby conferring two significant benefits. First, since these phage can be applied to affinity matrices at very high concentrations, large number of phage can be screened at one time. Second, since each infectious phage displays the combinatorial gene product on its surface, if a particular phage is recovered from an affinity matrix in low yield, the phage can be amplified by another round of infection. The group of almost identical *E.coli* filamentous phages M13, fd, and f1 are most often used in phage display libraries, as either of the phage gIII or gVIII coat proteins can be used to generate fusion proteins without disrupting the ultimate packaging of the viral particle (Ladner et al. PCT publication WO

90/02909; Garrard et al., PCT publication WO 92/09690; Marks et al. (1992) J. Biol. Chem. 267:16007-16010; Griffiths et al. (1993) EMBO J 12:725-734; Clackson et al. (1991) Nature 352:624-628; and Barbas et al. (1992) PNAS 89:4457-4461).

In an illustrative embodiment, the recombinant phage antibody system (RPAS, Pharmacia Catalog number 27-9400-01) can be easily modified for use in expressing and screening *hedgehog* combinatorial libraries. For instance, the pCANTAB 5 phagemid of the RPAS kit contains the gene which encodes the phage gIII coat protein. The *hedgehog* combinatorial gene library can be cloned into the phagemid adjacent to the gIII signal sequence such that it will be expressed as a gIII fusion protein. After ligation, the phagemid is used to transform competent *E. coli* TG1 cells. Transformed cells are subsequently infected with M13KO7 helper phage to rescue the phagemid and its candidate *hedgehog* gene insert. The resulting recombinant phage contain phagemid DNA encoding a specific candidate *hedgehog*, and display one or more copies of the corresponding fusion coat protein. The phage-displayed candidate *hedgehog* proteins which are capable of binding an *hedgehog* receptor are selected or enriched by panning. For instance, the phage library can be applied to cells which express the *patched* protein and unbound phage washed away from the cells. The bound phage is then isolated, and if the recombinant phage express at least one copy of the wild type gIII coat protein, they will retain their ability to infect *E. coli*. Thus, successive rounds of reinfection of *E. coli*, and panning will greatly enrich for *hedgehog* homologs, which can then be screened for further biological activities in order to differentiate agonists and antagonists.

Combinatorial mutagenesis has a potential to generate very large libraries of mutant proteins, e.g., in the order of 10^{26} molecules. Combinatorial libraries of this size may be technically challenging to screen even with high throughput screening assays such as phage display. To overcome this problem, a new technique has been developed recently, recursive ensemble mutagenesis (REM), which allows one to avoid the very high proportion of non-functional proteins in a random library and simply enhances the frequency of functional proteins, thus decreasing the complexity required to achieve a useful sampling of sequence space. REM is an algorithm which enhances the frequency of functional mutants in a library when an appropriate selection or screening method is employed (Arkin and Yourvan, 1992, *PNAS USA* 89:7811-7815; Yourvan et al., 1992, *Parallel Problem Solving from Nature*, 2., In Maenner and Manderick, eds., Elsevir Publishing Co., Amsterdam, pp. 401-410; Delgrave et al., 1993, *Protein Engineering* 6(3):327-331).

The invention also provides for reduction of the *hedgehog* protein to generate mimetics, e.g. peptide or non-peptide agents, which are able to disrupt binding of a *hedgehog* polypeptide of the present invention with an *hedgehog* receptor. Thus, such mutagenic techniques as

described above are also useful to map the determinants of the *hedgehog* proteins which participate in protein-protein interactions involved in, for example, binding of the subject *hedgehog* polypeptide to other extracellular matrix components. To illustrate, the critical residues of a subject *hedgehog* polypeptide which are involved in molecular recognition of an *hedgehog* receptor such as *patched* can be determined and used to generate *hedgehog*-derived peptidomimetics which competitively inhibit binding of the authentic *hedgehog* protein with that moiety. By employing, for example, scanning mutagenesis to map the amino acid residues of each of the subject *hedgehog* proteins which are involved in binding other extracellular proteins, peptidomimetic compounds can be generated which mimic those residues of the *hedgehog* protein which facilitate the interaction. Such mimetics may then be used to interfere with the normal function of a *hedgehog* protein. For instance, non-hydrolyzable peptide analogs of such residues can be generated using benzodiazepine (e.g., see Freidinger et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), azepine (e.g., see Huffman et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), substituted gamma lactam rings (Garvey et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), keto-methylene pseudopeptides (Ewenson et al. (1986) *J Med Chem* 29:295; and Ewenson et al. in *Peptides: Structure and Function* (Proceedings of the 9th American Peptide Symposium) Pierce Chemical Co. Rockland, IL, 1985), β -turn dipeptide cores (Nagai et al. (1985) *Tetrahedron Lett* 26:647; and Sato et al. (1986) *J Chem Soc Perkin Trans* 1:1231), and β -aminoalcohols (Gordon et al. (1985) *Biochem Biophys Res Commun* 126:419; and Dann et al. (1986) *Biochem Biophys Res Commun* 134:71).

Recombinantly produced forms of the *hedgehog* proteins can be produced using, e.g., expression vectors containing a nucleic acid encoding a *hedgehog* polypeptide, operably linked to at least one transcriptional regulatory sequence. Operably linked is intended to mean that the nucleotide sequence is linked to a regulatory sequence in a manner which allows expression of the nucleotide sequence. Regulatory sequences are art-recognized and are selected to direct expression of a *hedgehog* polypeptide. Accordingly, the term transcriptional regulatory sequence includes promoters, enhancers and other expression control elements. Such regulatory sequences are described in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). For instance, any of a wide variety of expression control sequences, sequences that control the expression of a DNA sequence when operatively linked to it, may be used in these vectors to express DNA sequences encoding *hedgehog* polypeptide. Such useful expression control sequences, include, for example, a viral LTR, such as the LTR of the Moloney murine leukemia virus, the early and late promoters of SV40, adenovirus or cytomegalovirus immediate early promoter, the lac system, the trp system, the

TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage λ , the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α -mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. Moreover, the vector's copy number, the ability to control that copy number and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered.

In addition to providing a ready source of hedgehog polypeptides for purification, the gene constructs of the present invention can also be used as a part of a gene therapy protocol to deliver nucleic acids encoding either an agonistic or antagonistic form of a *hedgehog* polypeptide. Thus, another aspect of the invention features expression vectors for *in vivo* transfection of a *hedgehog* polypeptide in particular cell types so as cause ectopic expression of a *hedgehog* polypeptide in an peripheral neurons or other cells associated therewith.

Formulations of such expression constructs may be administered in any biologically effective carrier, e.g. any formulation or composition capable of effectively delivering the recombinant gene to cells *in vivo*. Approaches include insertion of the hedgehog coding sequence in viral vectors including recombinant retroviruses, adenovirus, adeno-associated virus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors transfect cells directly; plasmid DNA can be delivered with the help of, for example, cationic liposomes (lipofectin) or derivatized (e.g. antibody conjugated), polylysine conjugates, gramicidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or CaPO_4 precipitation carried out *in vivo*. It will be appreciated that because transduction of appropriate target cells represents the critical first step in gene therapy, choice of the particular gene delivery system will depend on such factors as the phenotype of the intended target and the route of administration, e.g. locally or systemically. Furthermore, it will be recognized that the particular gene construct provided for *in vivo* transduction of *hedgehog* expression are also useful for *in vitro* transduction of cells, such as for use in the *ex vivo* tissue culture systems described below.

A preferred approach for *in vivo* introduction of nucleic acid into a cell is by use of a viral vector containing nucleic acid, e.g. a cDNA, encoding the particular form of the *hedgehog* polypeptide desired. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid. Additionally, molecules encoded

within the viral vector, e.g., by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid.

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery system of choice for the transfer of exogenous genes *in vivo*, particularly into humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. A major prerequisite for the use of retroviruses is to ensure the safety of their use, particularly with regard to the possibility of the spread of wild-type virus in the cell population. The development of specialized cell lines (termed "packaging cells") which produce only replication-defective retroviruses has increased the utility of retroviruses for gene therapy, and defective retroviruses are well characterized for use in gene transfer for gene therapy purposes (for a review see Miller, A.D. (1990) *Blood* 76:271). Thus, recombinant retrovirus can be constructed in which part of the retroviral coding sequence (*gag*, *pol*, *env*) has been replaced by nucleic acid encoding a hedgehog polypeptide and renders the retrovirus replication defective. The replication defective retrovirus is then packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F.M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include Crip, Cre, 2 and Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including neuronal cells, *in vitro* and/or *in vivo* (see for example Eglitis, et al. (1985) *Science* 230:1395-1398; Danos and Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:6460-6464; Wilson et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6141-6145; Huber et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Chowdhury et al. (1991) *Science* 254:1802-1805; van Beusechem et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:7640-7644; Kay et al. (1992) *Human Gene Therapy* 3:641-647; Dai et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu et al. (1993) *J. Immunol.* 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

Furthermore, it has been shown that it is possible to limit the infection spectrum of retroviruses and consequently of retroviral-based vectors, by modifying the viral packaging proteins on the surface of the viral particle (see, for example PCT publications WO93/25234 and

WO94/06920). For instance, strategies for the modification of the infection spectrum of retroviral vectors include: coupling antibodies specific for cell surface antigens to the viral *env* protein (Roux et al. (1989) *PNAS* 86:9079-9083; Julan et al. (1992) *J. Gen Virol* 73:3251-3255; and Goud et al. (1983) *Virology* 163:251-254); or coupling cell surface receptor ligands to the viral *env* proteins (Neda et al. (1991) *J Biol Chem* 266:14143-14146). Coupling can be in the form of the chemical cross-linking with a protein or other variety (e.g. lactose to convert the *env* protein to an asialoglycoprotein), as well as by generating fusion proteins (e.g. single-chain antibody/*env* fusion proteins). This technique, while useful to limit or otherwise direct the infection to certain tissue types, can also be used to convert an ecotropic vector in to an amphotropic vector.

Moreover, use of retroviral gene delivery can be further enhanced by the use of tissue- or cell-specific transcriptional regulatory sequences which control expression of the *hedgehog* gene of the retroviral vector.

Another viral gene delivery system useful in the present method utilizes adenovirus-derived vectors. The genome of an adenovirus can be manipulated such that it encodes and expresses a gene product of interest but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See for example Berkner et al. (1988) *BioTechniques* 6:616; Rosenfeld et al. (1991) *Science* 252:431-434; and Rosenfeld et al. (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (e.g., Ad2, Ad3, Ad7 etc.) are well known to those skilled in the art. Recombinant adenoviruses can be advantageous in certain circumstances in that they can be used to infect a wide variety of cell types, including peripheral nerve cells. Furthermore, the virus particle is relatively stable and amenable to purification and concentration, and as above, can be modified so as to affect the spectrum of infectivity. Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis in situations where introduced DNA becomes integrated into the host genome (e.g., retroviral DNA). Moreover, the carrying capacity of the adenoviral genome for foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner et al. cited *supra*; Haj-Ahmand and Graham (1986) *J. Virol.* 57:267). Most replication-defective adenoviral vectors currently in use and therefore favored by the present invention are deleted for all or parts of the viral E1 and E3 genes but retain as much as 80% of the adenoviral genetic material (see, e.g., Jones et al. (1979) *Cell* 16:683; Berkner et al., *supra*; and Graham et al. in Methods in Molecular Biology, E.J. Murray, Ed. (Humana, Clifton, NJ, 1991) vol. 7. pp. 109-127). Expression of the inserted *hedgehog* gene can be under control of, for example, the E1A promoter, the major late promoter

(MLP) and associated leader sequences, the E3 promoter, or exogenously added promoter sequences.

In addition to viral transfer methods, such as those illustrated above, non-viral methods can also be employed to cause expression of a *hedgehog* polypeptide in the tissue of an animal. Most nonviral methods of gene transfer rely on normal mechanisms used by mammalian cells for the uptake and intracellular transport of macromolecules. In preferred embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the *hedgehog* polypeptide gene by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, poly-lysine conjugates, and artificial viral envelopes.

In clinical settings, the gene delivery systems for the therapeutic *hedgehog* gene can be introduced into a patient by any of a number of methods, each of which is familiar in the art. For instance, a pharmaceutical preparation of the gene delivery system can be introduced systemically, e.g. by intravenous injection, and specific transduction of the protein in the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the receptor gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter (see U.S. Patent 5,328,470) or by stereotactic injection (e.g. Chen et al. (1994) *PNAS* 91: 3054-3057). A *hedgehog* expression construct can be delivered in a gene therapy construct to dermal cells by, e.g., electroporation using techniques described, for example, by Dev et al. ((1994) *Cancer Treat Rev* 20:105-115).

The pharmaceutical preparation of the gene therapy construct can consist essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery system can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can comprise one or more cells which produce the gene delivery system.

In yet another embodiment, the *hedgehog* or *ptc* therapeutic can be a "gene activation" construct which, by homologous recombination with a genomic DNA, alters the transcriptional regulatory sequences of an endogenous gene. For instance, the gene activation construct can replace the endogenous promoter of a *hedgehog* gene with a heterologous promoter, e.g., one which causes constitutive expression of the *hedgehog* gene or which causes inducible expression of the gene under conditions different from the normal expression pattern of the gene. Other genes in the *patched* signaling pathway can be similarly targeted. A variety of different formats for the gene activation constructs are available. See, for example, the Transkaryotic Therapies,

Inc PCT publications WO93/09222, WO95/31560, WO96/29411, WO95/31560 and WO94/12650.

In preferred embodiments, the nucleotide sequence used as the gene activation construct can be comprised of (1) DNA from some portion of the endogenous *hedgehog* gene (exon sequence, intron sequence, promoter sequences, etc.) which direct recombination and (2) heterologous transcriptional regulatory sequence(s) which is to be operably linked to the coding sequence for the genomic *hedgehog* gene upon recombination of the gene activation construct. For use in generating cultures of *hedgehog* producing cells, the construct may further include a reporter gene to detect the presence of the knockout construct in the cell.

The gene activation construct is inserted into a cell, and integrates with the genomic DNA of the cell in such a position so as to provide the heterologous regulatory sequences in operative association with the native *hedgehog* gene. Such insertion occurs by homologous recombination, i.e., recombination regions of the activation construct that are homologous to the endogenous *hedgehog* gene sequence hybridize to the genomic DNA and recombine with the genomic sequences so that the construct is incorporated into the corresponding position of the genomic DNA.

The terms "recombination region" or "targeting sequence" refer to a segment (i.e., a portion) of a gene activation construct having a sequence that is substantially identical to or substantially complementary to a genomic gene sequence, e.g., including 5' flanking sequences of the genomic gene, and can facilitate homologous recombination between the genomic sequence and the targeting transgene construct.

As used herein, the term "replacement region" refers to a portion of a activation construct which becomes integrated into an endogenous chromosomal location following homologous recombination between a recombination region and a genomic sequence.

The heterologous regulatory sequences, e.g., which are provided in the replacement region, can include one or more of a variety elements, including: promoters (such as constitutive or inducible promoters), enhancers, negative regulatory elements, locus control regions, transcription factor binding sites, or combinations thereof. Promoters/enhancers which may be used to control the expression of the targeted gene *in vivo* include, but are not limited to, the cytomegalovirus (CMV) promoter/enhancer (Karasuyama et al., 1989, *J. Exp. Med.*, 169:13), the human β -actin promoter (Gunning et al. (1987) *PNAS* 84:4831-4835), the glucocorticoid-inducible promoter present in the mouse mammary tumor virus long terminal repeat (MMTV LTR) (Klessig et al. (1984) *Mol. Cell Biol.* 4:1354-1362), the long terminal repeat sequences of Moloney murine leukemia virus (MuLV LTR) (Weiss et al. (1985) *RNA Tumor Viruses*, Cold

Spring Harbor Laboratory, Cold Spring Harbor, New York), the SV40 early or late region promoter (Bernoist et al. (1981) *Nature* 290:304-310; Templeton et al. (1984) *Mol. Cell Biol.*, 4:817; and Sprague et al. (1983) *J. Virol.*, 45:773), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (RSV) (Yamamoto et al., 1980, *Cell*, 22:787-797), the herpes simplex virus (HSV) thymidine kinase promoter/enhancer (Wagner et al. (1981) *PNAS* 82:3567-71), and the herpes simplex virus LAT promoter (Wolfe et al. (1992) *Nature Genetics*, 1:379-384).

In an exemplary embodiment, portions of the 5' flanking region of the human Shh gene are amplified using primers which add restriction sites, to generate the following fragments

5' -gcgcgccttcgaaGCGAGGCAGCCAGCGAGGGAGAGAGCGAGCGGGCGAGCCGGAGC-
GAGGAAatcgatgcgcgc (primer 1)

5' -gcgcgcagatctGGGAAAGCGCAAGAGAGAGCGCACACGCACACACCCGCCGCGCG-
CACTCGgatccgcgcgc (primer 2)

As illustrated, primer 1 includes a 5' non-coding region of the human Shh gene and is flanked by an AsuII and ClaI restriction sites. Primer 2 includes a portion of the 5' non-coding region immediately 3' to that present in primer 1. The hedgehog gene sequence is flanked by XhoII and BamHI restriction sites. The purified amplimers are cut with each of the enzymes as appropriate.

The vector pCDNA1.1 (Invitrogen) includes a CMV promoter. The plasmid is cut with with AsuII, which cleaves just 3' to the CMV promoter sequence. The AsuII/ClaI fragment of primer 1 is ligated to the AsuII cleavage site of the pcDNA vector. The ClaI/AsuII ligation destroys the AsuII site at the 3' end of a properly inserted primer 1.

The vector is then cut with BamHI, and an XhoII/BamHI fragment of primer 2 is ligated to the BamHI cleavage site. As above, the BamHI/XhoII ligation destroys the BamHI site at the 5' end of a properly inserted primer 2.

Individual colonies are selected, cut with AsuII and BamHI, and the size of the AsuII/BamHI fragment determined. Colonies in which both the primer 1 and primer 2 sequences are correctly inserted are further amplified, an cut with AsuII and BamHI to produce the gene activation construct

cgaagcgaggcagccagcgagggagagagcgagcgggagcgagccgagcgaggaaATCGAAGGTTC
GAATCCTTCCCCACCACCATCACTTTCAAAGTCCGAAAGAATCTGCTCCCTGCTTGTGTGTTG
GAGGTCGCTGAGTAGTGCGCGAGTAAAATTTAAGCTACAACAAGGCAAGGCTTGACCGACAATTG

CATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTACGGGGCCAGATATACGC
GTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCA
TATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCC
CCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCATTGAC
5 GTCAATGGGTGGACTATTTACGGTAAACTGCCCCTTGGCAGTACATCAAGTGTATCATATGCCA
AGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCAGTACATGAC
CTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGC
GGTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCGAAGTCTCCAC
CCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAATCAACGGGACTTTCGAAATGTGCTAA
10 CAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAG
CTCTCTGGCTAACTAGAGAACCCACTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAGG
GAGACCCAAGCTTGGTACCGAGCTCGGATCgatctgggaaagcgcaagagagagcgcacacgcac
acacccgccgcgcgcactcgg

In this construct, the flanking primer 1 and primer 2 sequences provide the recombination region
which permits the insertion of the CMV promoter in front of the coding sequence for the human
Shh gene. Other heterologous promoters (or other transcriptional regulatory sequences) can be
inserted in a genomic *hedgehog* gene by a similar method.

In still other embodiments, the replacement region merely deletes a negative
transcriptional control element of the native gene, e.g., to activate expression, or ablates a
positive control element, e.g., to inhibit expression of the targeted gene.

V. Exemplary *ptc* therapeutic compounds.

In another embodiment, the subject method is carried out using a *ptc* therapeutic
composition. Such compositions can be generated with, for example, compounds which bind to
patched and alter its signal transduction activity, compounds which alter the binding and/or
enzymatic activity of a protein (e.g., intracellular) involved in patched signal pathway, and
compounds which alter the level of expression of a hedgehog protein, a patched protein or a
protein involved in the intracellular signal transduction pathway of patched.

The availability of purified and recombinant *hedgehog* polypeptides facilitates the
generation of assay systems which can be used to screen for drugs, such as small organic
molecules, which are either agonists or antagonists of the normal cellular function of a *hedgehog*
and/or patched protein, particularly their role in the pathogenesis of peripheral nerve proliferation
and/or differentiation. In one embodiment, the assay evaluates the ability of a compound to
modulate binding between a *hedgehog* polypeptide and a *hedgehog* receptor such as *patched*. In

other embodiments, the assay merely scores for the ability of a test compound to alter the signal transduction activity of the *patched* protein. In this manner, a variety of *hedgehog* and/or *ptc* therapeutics, both proliferative and anti-proliferative in activity, can be identified. A variety of assay formats will suffice and, in light of the present disclosure, will be comprehended by skilled artisan.

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with receptor proteins.

Accordingly, in an exemplary screening assay for *ptc* therapeutics, the compound of interest is contacted with a mixture including a *hedgehog* receptor protein (e.g., a cell expressing the *patched* receptor) and a *hedgehog* protein under conditions in which it is ordinarily capable of binding the *hedgehog* protein. To the mixture is then added a composition containing a test compound. Detection and quantification of receptor/*hedgehog* complexes provides a means for determining the test compound's efficacy at inhibiting (or potentiating) complex formation between the receptor protein and the *hedgehog* polypeptide. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control assay can also be performed to provide a baseline for comparison. In the control assay, isolated and purified *hedgehog* polypeptide is added to the receptor protein, and the formation of receptor/*hedgehog* complex is quantitated in the absence of the test compound.

In other embodiments, a *ptc* therapeutic of the present invention is one which disrupts the association of *patched* with *smoothened*.

Agonist and antagonists of peripheral nerve maintenance can be distinguished, and the efficacy of the compound can be assessed, by subsequent testing with peripheral nerve cells, e.g., in culture.

In an illustrative embodiment, the polypeptide utilized as a *hedgehog* receptor can be generated from the *patched* protein. Accordingly, an exemplary screening assay includes all or a suitable portion of the *patched* protein which can be obtained from, for example, the human

patched gene (GenBank U43148) or other vertebrate sources (see GenBank Accession numbers U40074 for chicken *patched* and U46155 for mouse *patched*), as well as from drosophila (GenBank Accession number M28999) or other invertebrate sources. The *patched* protein can be provided in the screening assay as a whole protein (preferably expressed on the surface of a cell), or alternatively as a fragment of the full length protein which binds to *hedgehog* polypeptides, e.g., as one or both of the substantial extracellular domains (e.g. corresponding to residues Asn120-Ser438 and/or Arg770-Trp1027 of the human *patched* protein - which are also potential antagonists of *hedgehog*-dependent signal transduction). For instance, the *patched* protein can be provided in soluble form, as for example a preparation of one of the extracellular domains, or a preparation of both of the extracellular domains which are covalently connected by an unstructured linker (see, for example, Huston et al. (1988) PNAS 85:4879; and U.S. Patent No. 5,091,513). In other embodiments, the protein can be provided as part of a liposomal preparation or expressed on the surface of a cell. The *patched* protein can be derived from a recombinant gene, e.g., being ectopically expressed in a heterologous cell. For instance, the protein can be expressed on oocytes, mammalian cells (e.g., COS, CHO, 3T3 or the like), or yeast cell by standard recombinant DNA techniques. These recombinant cells can be used for receptor binding, signal transduction or gene expression assays. Marigo et al. (1996) *Development* 122:1225-1233 illustrates a binding assay of human *hedgehog* to chick *patched* protein ectopically expressed in *Xenopus laevis* oocytes. The assay system of Marigo et al. can be adapted to the present drug screening assays. As illustrated in that reference, *Shh* binds to the *patched* protein in a selective, saturable, dose-dependent manner, thus demonstrating that *patched* is a receptor for *Shh*.

Complex formation between the *hedgehog* polypeptide and a *hedgehog* receptor may be detected by a variety of techniques. For instance, modulation of the formation of complexes can be quantitated using, for example, detectably labelled proteins such as radiolabelled, fluorescently labelled, or enzymatically labelled *hedgehog* polypeptides, by immunoassay, or by chromatographic detection.

Typically, for cell-free assays, it will be desirable to immobilize either the *hedgehog* receptor or the *hedgehog* polypeptide to facilitate separation of receptor/*hedgehog* complexes from uncomplexed forms of one of the proteins, as well as to accommodate automation of the assay. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase/receptor (GST/receptor) fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the *hedgehog* polypeptide, e.g. an ³⁵S-labeled *hedgehog* polypeptide, and the test compound and incubated

under conditions conducive to complex formation, e.g. at physiological conditions for salt and pH, though slightly more stringent conditions may be desired. Following incubation, the beads are washed to remove any unbound *hedgehog* polypeptide, and the matrix bead-bound radiolabel determined directly (e.g. beads placed in scintillant), or in the supernatant after the receptor/*hedgehog* complexes are dissociated. Alternatively, the complexes can be dissociated from the bead, separated by SDS-PAGE gel, and the level of *hedgehog* polypeptide found in the bead fraction quantitated from the gel using standard electrophoretic techniques.

Other techniques for immobilizing proteins on matrices are also available for use in the subject assay. For instance, soluble portions of the *hedgehog* receptor protein can be immobilized utilizing conjugation of biotin and streptavidin. For instance, biotinylated receptor molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with the *hedgehog* receptor but which do not interfere with *hedgehog* binding can be derivatized to the wells of the plate, and the receptor trapped in the wells by antibody conjugation. As above, preparations of a *hedgehog* polypeptide and a test compound are incubated in the receptor-presenting wells of the plate, and the amount of receptor/*hedgehog* complex trapped in the well can be quantitated. Exemplary methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the *hedgehog* polypeptide, or which are reactive with the receptor protein and compete for binding with the *hedgehog* polypeptide; as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the *hedgehog* polypeptide. In the instance of the latter, the enzyme can be chemically conjugated or provided as a fusion protein with the *hedgehog* polypeptide. To illustrate, the *hedgehog* polypeptide can be chemically cross-linked or genetically fused with alkaline phosphatase, and the amount of *hedgehog* polypeptide trapped in the complex can be assessed with a chromogenic substrate of the enzyme, e.g. paranitrophenylphosphate. Likewise, a fusion protein comprising the *hedgehog* polypeptide and glutathione-S-transferase can be provided, and complex formation quantitated by detecting the GST activity using 1-chloro-2,4-dinitrobenzene (Habig et al (1974) *J Biol Chem* 249:7130).

For processes which rely on immunodetection for quantitating one of the proteins trapped in the complex, antibodies against the protein, such as the anti-*hedgehog* antibodies described herein, can be used. Alternatively, the protein to be detected in the complex can be "epitope tagged" in the form of a fusion protein which includes, in addition to the *hedgehog* polypeptide or *hedgehog* receptor sequence, a second polypeptide for which antibodies are readily available

(e.g. from commercial sources). For instance, the GST fusion proteins described above can also be used for quantification of binding using antibodies against the GST moiety. Other useful epitope tags include myc-epitopes (e.g., see Ellison et al. (1991) *J Biol Chem* 266:21150-21157) which includes a 10-residue sequence from c-myc, as well as the pFLAG system (International Biotechnologies, Inc.) or the pEZZ-protein A system (Pharmacia, NJ).

Where the desired portion of the *hedgehog* receptor (or other *hedgehog* binding molecule) cannot be provided in soluble form, liposomal vesicles can be used to provide manipulatable and isolatable sources of the receptor. For example, both authentic and recombinant forms of the *patched* protein can be reconstituted in artificial lipid vesicles (e.g. phosphatidylcholine liposomes) or in cell membrane-derived vesicles (see, for example, Bear et al. (1992) *Cell* 68:809-818; Newton et al. (1983) *Biochemistry* 22:6110-6117; and Reber et al. (1987) *J Biol Chem* 262:11369-11374).

In addition to cell-free assays, such as described above, the readily available source of *hedgehog* proteins provided by the art also facilitates the generation of cell-based assays for identifying small molecule agonists/antagonists and the like. Analogous to the cell-based assays described above for screening combinatorial libraries, cells which are sensitive to *hedgehog* induction, e.g. *patched*-expressing cells or other myoblast-derived cells sensitive to *hedgehog* induction, can be contacted with a *hedgehog* protein and a test agent of interest, with the assay scoring for anything from simple binding to the cell to modulation in *hedgehog* inductive responses by the target cell in the presence and absence of the test agent. As with the cell-free assays, agents which produce a statistically significant change in *hedgehog* activities (either inhibition or potentiation) can be identified.

In other embodiments, the cell-based assay scores for agents which disrupt association of *patched* and *smoothed* proteins, e.g., in the cell surface membrane or liposomal preparation.

In addition to characterizing cells that naturally express the *patched* protein, cells which have been genetically engineered to ectopically express *patched* can be utilized for drug screening assays. As an example, cells which either express low levels or lack expression of the *patched* protein, e.g. *Xenopus laevis* oocytes, COS cells or yeast cells, can be genetically modified using standard techniques to ectopically express the *patched* protein. (see Marigo et al., *supra*).

The resulting recombinant cells, e.g., which express a functional *patched* receptor, can be utilized in receptor binding assays to identify agonist or antagonist of *hedgehog* binding. Binding assays can be performed using whole cells. Furthermore, the recombinant cells of the present invention can be engineered to include other heterologous genes encoding proteins

involved in *hedgehog*-dependent signal pathways. For example, the gene products of one or more of *smoothened*, *costal-2* and/or *fused* can be co-expressed with *patched* in the reagent cell, with assays being sensitive to the functional reconstitution of the *hedgehog* signal transduction cascade.

Alternatively, liposomal preparations using reconstituted *patched* protein can be utilized. *Patched* protein purified from detergent extracts from both authentic and recombinant origins can be reconstituted in artificial lipid vesicles (e.g. phosphatidylcholine liposomes) or in cell membrane-derived vesicles (see, for example, Bear et al. (1992) *Cell* 68:809-818; Newton et al. (1983) *Biochemistry* 22:6110-6117; and Reber et al. (1987) *J Biol Chem* 262:11369-11374). The lamellar structure and size of the resulting liposomes can be characterized using electron microscopy. External orientation of the *patched* protein in the reconstituted membranes can be demonstrated, for example, by immunoelectron microscopy. The *hedgehog* protein binding activity of liposomes containing *patched* and liposomes without the protein in the presence of candidate agents can be compared in order to identify potential modulators of the *hedgehog-patched* interaction.

The *hedgehog* protein used in these cell-based assays can be provided as a purified source (natural or recombinant in origin), or in the form of cells/tissue which express the protein and which are co-cultured with the target cells. As in the cell-free assays, where simple binding (rather than induction) is the *hedgehog* activity scored for in the assay, the protein can be labelled by any of the above-mentioned techniques, e.g., fluorescently, enzymatically or radioactively, or detected by immunoassay.

In addition to binding studies, functional assays can be used to identify modulators, i.e., agonists or antagonists, of *hedgehog* or *patched* activities. By detecting changes in intracellular signals, such as alterations in second messengers or gene expression, in *patched*-expressing cells contacted with a test agent, candidate agonists and antagonists to *patched* signaling can be identified.

A number of gene products have been implicated in *patched*-mediated signal transduction, including *patched*, the transcription factor *cubitus interruptus* (ci), the serine/threonine kinase *fused* (fu) and the gene products of *costal-2*, *smoothened* and *suppressor of fused*.

The interaction of a *hedgehog* protein with *patched* sets in motion a cascade involving the activation and inhibition of downstream effectors, the ultimate consequence of which is, in some instances, a detectable change in the transcription or translation of a gene. Potential transcriptional targets of *patched* signaling are the *patched* gene itself (Hidalgo and Ingham, 1990 *Development* 110, 291-301; Marigo et al., 1996) and the vertebrate homologs of the

drosophila cubitus interruptus gene, the *GLI* genes (Hui et al. (1994) *Dev Biol* 162:402-413). *Patched* gene expression has been shown to be induced in cells of the limb bud and the neural plate that are responsive to *Shh*. (Marigo et al. (1996) *PNAS*, in press; Marigo et al. (1996) *Development* 122:1225-1233). The *GLI* genes encode putative transcription factors having zinc
5 finger DNA binding domains (Orenic et al. (1990) *Genes & Dev* 4:1053-1067; Kinzler et al. (1990) *Mol Cell Biol* 10:634-642). Transcription of the *GLI* gene has been reported to be upregulated in response to *hedgehog* in limb buds, while transcription of the *GLI3* gene is downregulated in response to *hedgehog* induction (Marigo et al. (1996) *Development* 122:1225-1233). By selecting transcriptional regulatory sequences from such target genes, e.g. from
10 *patched* or *GLI* genes, that are responsible for the up- or down regulation of these genes in response to *patched* signalling, and operatively linking such promoters to a reporter gene, one can derive a transcription based assay which is sensitive to the ability of a specific test compound to modify *patched* signalling pathways. Expression of the reporter gene, thus, provides a valuable screening tool for the development of compounds that act as agonists or antagonists of
15 *ptc* induction of differentiation/quiescence.

Reporter gene based assays of this invention measure the end stage of the above described cascade of events, e.g., transcriptional modulation. Accordingly, in practicing one embodiment of the assay, a reporter gene construct is inserted into the reagent cell in order to generate a detection signal dependent on *ptc* signaling. To identify potential regulatory elements
20 responsive to *ptc* signaling present in the transcriptional regulatory sequence of a target gene, nested deletions of genomic clones of the target gene can be constructed using standard techniques. See, for example, Current Protocols in Molecular Biology, Ausubel, F.M. et al. (eds.) Greene Publishing Associates, (1989); U.S. Patent 5,266,488; Sato et al. (1995) *J Biol Chem* 270:10314-10322; and Kube et al. (1995) *Cytokine* 7:1-7. A nested set of DNA fragments
25 from the gene's 5'-flanking region are placed upstream of a reporter gene, such as the luciferase gene, and assayed for their ability to direct reporter gene expression in *patched* expressing cells. Host cells transiently transfected with reporter gene constructs can be scored for the induction of expression of the reporter gene in the presence and absence of *hedgehog* to determine regulatory sequences which are response to *patched*-dependent signalling.

30 In practicing one embodiment of the assay, a reporter gene construct is inserted into the reagent cell in order to generate a detection signal dependent on second messengers generated by induction with *hedgehog* protein. Typically, the reporter gene construct will include a reporter gene in operative linkage with one or more transcriptional regulatory elements responsive to the *hedgehog* activity, with the level of expression of the reporter gene providing the *hedgehog*-
35 dependent detection signal. The amount of transcription from the reporter gene may be

measured using any method known to those of skill in the art to be suitable. For example, mRNA expression from the reporter gene may be detected using RNase protection or RNA-based PCR, or the protein product of the reporter gene may be identified by a characteristic stain or an intrinsic activity. The amount of expression from the reporter gene is then compared to the amount of expression in either the same cell in the absence of the test compound (or *hedgehog*) or it may be compared with the amount of transcription in a substantially identical cell that lacks the target receptor protein. Any statistically or otherwise significant difference in the amount of transcription indicates that the test compound has in some manner altered the signal transduction of the *patched* protein, e.g., the test compound is a potential ptc therapeutic.


As described in further detail below, in preferred embodiments the gene product of the reporter is detected by an intrinsic activity associated with that product. For instance, the reporter gene may encode a gene product that, by enzymatic activity, gives rise to a detection signal based on color, fluorescence, or luminescence. In other preferred embodiments, the reporter or marker gene provides a selective growth advantage, e.g., the reporter gene may enhance cell viability, relieve a cell nutritional requirement, and/or provide resistance to a drug.

Preferred reporter genes are those that are readily detectable. The reporter gene may also be included in the construct in the form of a fusion gene with a gene that includes desired transcriptional regulatory sequences or exhibits other desirable properties. Examples of reporter genes include, but are not limited to CAT (chloramphenicol acetyl transferase) (Alton and Vapnek (1979), Nature 282: 864-869) luciferase, and other enzyme detection systems, such as beta-galactosidase; firefly luciferase (deWet et al. (1987), Mol. Cell. Biol. 7:725-737); bacterial luciferase (Engebrecht and Silverman (1984), PNAS 1: 4154-4158; Baldwin et al. (1984), Biochemistry 23: 3663-3667); alkaline phosphatase (Toh et al. (1989) Eur. J. Biochem. 182: 231-238, Hall et al. (1983) J. Mol. Appl. Gen. 2: 101), human placental secreted alkaline phosphatase (Cullen and Malim (1992) Methods in Enzymol. 216:362-368).

Transcriptional control elements which may be included in a reporter gene construct include, but are not limited to, promoters, enhancers, and repressor and activator binding sites. Suitable transcriptional regulatory elements may be derived from the transcriptional regulatory regions of genes whose expression is induced after modulation of a *patched* signal transduction pathway. The characteristics of preferred genes from which the transcriptional control elements are derived include, but are not limited to, low or undetectable expression in quiescent cells, rapid induction at the transcriptional level within minutes of extracellular stimulation, induction that is transient and independent of new protein synthesis, subsequent shut-off of transcription requires new protein synthesis, and mRNAs transcribed from these genes have a short half-life. It is not necessary for all of these properties to be present.

In yet other embodiments, second messenger generation can be measured directly in the detection step, such as mobilization of intracellular calcium, phospholipid metabolism or adenylate cyclase activity are quantitated, for instance, the products of phospholipid hydrolysis IP₃, DAG or cAMP could be measured. For example, recent studies have implicated protein kinase A (PKA) as a possible component of *hedgehog/patched* signaling (Hammerschmidt et al. (1996) *Genes & Dev* 10:647). High PKA activity has been shown to antagonize *hedgehog* signaling in these systems. Although it is unclear whether PKA acts directly downstream or in parallel with *hedgehog* signaling, it is possible that *hedgehog* signalling occurs via inhibition of PKA activity. Thus, detection of PKA activity provides a potential readout for the instant assays.

In a preferred embodiment, the *ptc* therapeutic is a PKA inhibitor. A variety of PKA inhibitors are known in the art, including both peptidyl and organic compounds. For instance, the *ptc* therapeutic can be a 5-isoquinolinesulfonamide, such as represented in the general formula:



wherein,

R₁ and R₂ each can independently represent hydrogen, and as valence and stability permit a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-R₈, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₈, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₈, or

R₁ and R₂ taken together with N form a heterocycle (substituted or unsubstituted);


R₃ is absent or represents one or more substitutions to the isoquinoline ring such as a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-

R_8 , $-(CH_2)_m-OH$, $-(CH_2)_m-O$ -lower alkyl, $-(CH_2)_m-O$ -lower alkenyl, $-(CH_2)_n-O-(CH_2)_m-R_8$, $-(CH_2)_m-SH$, $-(CH_2)_m-S$ -lower alkyl, $-(CH_2)_m-S$ -lower alkenyl, $-(CH_2)_n-S-(CH_2)_m-R_8$;


R_8 represents a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle; and

5 n and m are independently for each occurrence zero or an integer in the range of 1 to 6.


In a preferred embodiment, the PKA inhibitor is N-[2-((p-bromocinnamyl)amino)ethyl]-5-isoquinolinesulfonamide (H-89; Calbiochem Cat. No. 371963), e.g., having the formula:





In another embodiment, the PKA inhibitor is 1-(5-isoquinolinesulfonyl)-2-methylpiperazine (H-7; Calbiochem Cat. No. 371955), e.g., having the formula:



In still other embodiments, the PKA inhibitor is KT5720 (Calbiochem Cat. No. 420315), having the structure



A variety of nucleoside analogs are also useful as PKA inhibitors. For example, the subject method can be carried out cyclic AMP analogs which inhibit the kinase activity of PKA, as for example, 8-bromo-cAMP or dibutyryl-cAMP



Exemplary peptidyl inhibitors of PKA activity include the PKA Heat Stable Inhibitor (isoform α ; see, for example, Calbiochem Cat. No. 539488, and Wen et al. (1995) *J Biol Chem* 270:2041).

Certain *hedehog* receptors may stimulate the activity of phospholipases. Inositol lipids can be extracted and analyzed using standard lipid extraction techniques. Water soluble derivatives of all three inositol lipids (IP₁, IP₂, IP₃) can also be quantitated using radiolabelling techniques or HPLC.

The mobilization of intracellular calcium or the influx of calcium from outside the cell may be a response to *hedehog* stimulation or lack thereof. Calcium flux in the reagent cell can be measured using standard techniques. The choice of the appropriate calcium indicator, fluorescent, bioluminescent, metallochromic, or Ca⁺⁺-sensitive microelectrodes depends on the cell type and the magnitude and time constant of the event under study (Borle (1990) *Environ Health Perspect* 84:45-56). As an exemplary method of Ca⁺⁺ detection, cells could be loaded

with the Ca^{++} sensitive fluorescent dye fura-2 or indo-1, using standard methods, and any change in Ca^{++} measured using a fluorometer.

In certain embodiments of the assay, it may be desirable to screen for changes in cellular phosphorylation. As an example, the drosophila gene *fused* (*fu*) which encodes a serine/threonine kinase has been identified as a potential downstream target in *hedgehog* signaling. (Preat et al., 1990 *Nature* 347, 87-89; Therond et al. 1993, *Mech. Dev.* 44. 65-80). The ability of compounds to modulate serine/threonine kinase activation could be screened using colony immunoblotting (Lyons and Nelson (1984) *Proc. Natl. Acad. Sci. USA* 81:7426-7430) using antibodies against phosphorylated serine or threonine residues. Reagents for performing such assays are commercially available, for example, phosphoserine and phosphothreonine specific antibodies which measure increases in phosphorylation of those residues can be purchased from commercial sources.

In yet another embodiment, the *ptc* therapeutic is an antisense molecule which inhibits expression of a protein involved in a *patched*-mediated signal transduction pathway. To illustrate, by inhibiting the expression of a protein which are involved in *patched* signals, such as *fused*, *costal-2*, *smoothed* and/or *Gli* genes, the ability of the *patched* signal pathway(s) to inhibit proliferation of a cell can be altered, e.g., potentiated or repressed.

As used herein, "antisense" therapy refers to administration or *in situ* generation of oligonucleotide probes or their derivatives which specifically hybridize (e.g. bind) under cellular conditions with cellular mRNA and/or genomic DNA encoding a *hedgehog* protein, *patched*, or a protein involved in *patched*-mediated signal transduction. The hybridization should inhibit expression of that protein, e.g. by inhibiting transcription and/or translation. The binding may be by conventional base pair complementarity, or, for example, in the case of binding to DNA duplexes, through specific interactions in the major groove of the double helix. In general, "antisense" therapy refers to the range of techniques generally employed in the art, and includes any therapy which relies on specific binding to oligonucleotide sequences.

An antisense construct of the present invention can be delivered, for example, as an expression plasmid which, when transcribed in the cell, produces RNA which is complementary to at least a unique portion of the target cellular mRNA. Alternatively, the antisense construct is an oligonucleotide probe which is generated *ex vivo* and which, when introduced into the cell causes inhibition of expression by hybridizing with the mRNA and/or genomic sequences of a target gene. Such oligonucleotide probes are preferably modified oligonucleotide which are resistant to endogenous nucleases, e.g. exonucleases and/or endonucleases, and is therefore stable *in vivo*. Exemplary nucleic acid molecules for use as antisense oligonucleotides are phosphoramidate, phosphothioate and methylphosphonate analogs of DNA (see also U.S. Patents

5,176,996; 5,264,564; and 5,256,775). Additionally, general approaches to constructing oligomers useful in antisense therapy have been reviewed, for example, by Van der Krol et al. (1988) *Biotechniques* 6:958-976; and Stein et al. (1988) *Cancer Res* 48:2659-2668.

Several considerations should be taken into account when constructing antisense oligonucleotides for the use in the methods of the invention: (1) oligos should have a GC content of 50% or more; (2) avoid sequences with stretches of 3 or more G's; and (3) oligonucleotides should not be longer than 25-26 mers. When testing an antisense oligonucleotide, a mismatched control can be constructed. The controls can be generated by reversing the sequence order of the corresponding antisense oligonucleotide in order to conserve the same ratio of bases.

In an illustrative embodiment, the *ptc* therapeutic can be an antisense construct for inhibiting the expression of *patched*, e.g., to mimic the inhibition of *patched* by *hedgehog*. Exemplary antisense constructs include:

5'-GTCCTGGCGCCGCCGCCGCCGTCGCC

5'-TTCCGATGACCGGCCTTTCGCGGTGA

5'-GTGCACGGAAAGGTGCAGGCCACACT

VI. Exemplary pharmaceutical preparations of *hedgehog* and *ptc* therapeutics

The source of the *hedgehog* and *ptc* therapeutics to be formulated will depend on the particular form of the agent. Small organic molecules and peptidyl fragments can be chemically synthesized and provided in a pure form suitable for pharmaceutical/cosmetic usage. Products of natural extracts can be purified according to techniques known in the art. For example, the Cox et al. U.S. Patent 5,286,654 describes a method for purifying naturally occurring forms of a secreted protein and can be adapted for purification of *hedgehog* polypeptides. Recombinant sources of *hedgehog* polypeptides are also available. For example, the gene encoding *hedgehog* polypeptides, are known, *inter alia*, from PCT publications WO 95/18856 and WO 96/17924.

Those of skill in treating peripheral neuropathies can determine the effective amount of an *hedgehog* or *ptc* therapeutic to be formulated in a pharmaceutical or cosmetic preparation.

The *hedgehog* or *ptc* therapeutic formulations used in the method of the invention are most preferably applied in the form of appropriate compositions. As appropriate compositions there may be cited all compositions usually employed for systemically or topically administering drugs. The pharmaceutically acceptable carrier should be substantially inert, so as not to act with

the active component. Suitable inert carriers include water, alcohol polyethylene glycol, mineral oil or petroleum gel, propylene glycol and the like.

To prepare the pharmaceutical compositions of this invention, an effective amount of the particular hedgehog or ptc therapeutic as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represents the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

In addition to the direct topical application of the preparations they can be topically administered by other methods, for example, encapsulated in a temperature and/or pressure sensitive matrix or in film or solid carrier which is soluble in body fluids and the like for subsequent release, preferably sustained-release of the active component.

As appropriate compositions for topical application there may be cited all compositions usually employed for topically administering therapeutics, e.g., creams, gellies, dressings, shampoos, tinctures, pastes, ointments, salves, powders, liquid or semiliquid formulation and the like. Application of said compositions may be by aerosol e.g. with a propellant such as nitrogen carbon dioxide, a freon, or without a propellant such as a pump spray, drops, lotions, or a semisolid such as a thickened composition which can be applied by a swab. In particular

compositions, semisolid compositions such as salves, creams, pastes, gellies, ointments and the like will conveniently be used.

It is especially advantageous to formulate the subject compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powders packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The pharmaceutical preparations of the present invention can be used, as stated above, for the many applications which can be considered cosmetic uses. Cosmetic compositions known in the art, preferably hypoallergic and pH controlled are especially preferred, and include toilet waters, packs, lotions, skin milks or milky lotions. The preparations contain, besides the hedgehog or ptc therapeutic, components usually employed in such preparations. Examples of such components are oils, fats, waxes, surfactants, humectants, thickening agents, antioxidants, viscosity stabilizers, chelating agents, buffers, preservatives, perfumes, dyestuffs, lower alkanols, and the like. If desired, further ingredients may be incorporated in the compositions, e.g. antiinflammatory agents, antibacterials, antifungals, disinfectants, vitamins, sunscreens, antibiotics, or other anti-acne agents.

Examples of oils comprise fats and oils such as olive oil and hydrogenated oils; waxes such as beeswax and lanolin; hydrocarbons such as liquid paraffin, ceresin, and squalane; fatty acids such as stearic acid and oleic acid; alcohols such as cetyl alcohol, stearyl alcohol, lanolin alcohol, and hexadecanol; and esters such as isopropyl myristate, isopropyl palmitate and butyl stearate. As examples of surfactants there may be cited anionic surfactants such as sodium stearate, sodium cetylsulfate, polyoxyethylene laurylether phosphate, sodium N-acyl glutamate; cationic surfactants such as stearyldimethylbenzylammonium chloride and stearyltrimethylammonium chloride; ampholytic surfactants such as alkylaminoethylglycine hydrochloride solutions and lecithin; and nonionic surfactants such as glycerin monostearate, sorbitan monostearate, sucrose fatty acid esters, propylene glycol monostearate, polyoxyethylene oleylether, polyethylene glycol monostearate, polyoxyethylene sorbitan monopalmitate, polyoxyethylene coconut fatty acid monoethanolamide, polyoxypropylene glycol (e.g. the materials sold under the trademark "Pluronic"), polyoxyethylene castor oil, and polyoxyethylene lanolin. Examples of humectants include glycerin, 1,3-butylene glycol, and propylene glycol; examples of lower alcohols include ethanol and isopropanol; examples of thickening agents

include xanthan gum, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, polyethylene glycol and sodium carboxymethyl cellulose; examples of antioxidants comprise butylated hydroxytoluene, butylated hydroxyanisole, propyl gallate, citric acid and ethoxyquin; examples of chelating agents include disodium edetate and ethanedihydroxy diphosphate; examples of buffers comprise citric acid, sodium citrate, boric acid, borax, and disodium hydrogen phosphate; and examples of preservatives are methyl parahydroxybenzoate, ethyl parahydroxybenzoate, dehydroacetic acid, salicylic acid and benzoic acid.

For preparing ointments, creams, toilet waters, skin milks, and the like, typically from 0.01 to 10% in particular from 0.1 to 5% and more in particular from 0.2 to 2.5% of the active ingredient, e.g., of the hedgehog or ptc therapeutic, will be incorporated in the compositions. In ointments or creams, the carrier for example consists of 1 to 20%, in particular 5 to 15% of a humectant, 0.1 to 10% in particular from 0.5 to 5% of a thickener and water; or said carrier may consist of 70 to 99%, in particular 20 to 95% of a surfactant, and 0 to 20%, in particular 2.5 to 15% of a fat; or 80 to 99.9% in particular 90 to 99% of a thickener; or 5 to 15% of a surfactant, 2-15% of a humectant, 0 to 80% of an oil, very small (< 2%) amounts of preservative, coloring agent and/or perfume, and water. In a toilet water, the carrier for example consists of 2 to 10% of a lower alcohol, 0.1 to 10% or in particular 0.5 to 1% of a surfactant, 1 to 20%, in particular 3 to 7% of a humectant, 0 to 5% of a buffer, water and small amounts (< 2%) of preservative, dyestuff and/or perfume. In a skin milk, the carrier typically consists of 10-50% of oil, 1 to 10% of surfactant, 50-80% of water and 0 to 3% of preservative and/or perfume. In the aforementioned preparations, all % symbols refer to weight by weight percentage.

Particular compositions for use in the method of the present invention are those wherein the hedgehog or ptc therapeutic is formulated in liposome-containing compositions. Liposomes are artificial vesicles formed by amphiphatic molecules such as polar lipids, for example, phosphatidyl cholines, ethanolamines and serines, sphingomyelins, cardiolipins, plasmalogens, phosphatidic acids and cerebiosides. Liposomes are formed when suitable amphiphatic molecules are allowed to swell in water or aqueous solutions to form liquid crystals usually of multilayer structure comprised of many bilayers separated from each other by aqueous material (also referred to as coarse liposomes). Another type of liposome known to be consisting of a single bilayer encapsulating aqueous material is referred to as a unilamellar vesicle. If water-soluble materials are included in the aqueous phase during the swelling of the lipids they become entrapped in the aqueous layer between the lipid bilayers.

Water-soluble active ingredients such as, for example, various salt forms of a hedgehog polypeptide, are encapsulated in the aqueous spaces between the molecular layers. The lipid soluble active ingredient of hedgehog or ptc therapeutic, such as an organic mimetic, is

predominantly incorporated into the lipid layers, although polar head groups may protude from the layer into the aqueous space. The encapsulation of these compounds can be achieved by a number of methods. The method most commonly used involves casting a thin film of phospholipid onto the walls of a flask by evaporation from an organic solvent. When this film is dispersed in a suitable aqueous medium, multilamellar liposomes are formed. Upon suitable sonication, the coarse liposomes form smaller similarly closed vesicles.

Water-soluble active ingredients are usually incorporated by dispersing the cast film with an aqueous solution of the compound. The unencapsulated compound is then removed by centrifugation, chromatography, dialysis or other art-known suitable procedures. The lipid-soluble active ingredient is usually incorporated by dissolving it in the organic solvent with the phospholipid prior to casting the film. If the solubility of the material in the lipid phase is not exceeded or the amount present is not in excess of that which can be bound to the lipid, liposomes prepared by the above method usually contain most of the material bound in the lipid bilayers; separation of the liposomes from unencapsulated material is not required.

A particularly convenient method for preparing liposome formulated forms of hedgehog and ptc therapeutics is the method described in EP-A-253,619, incorporated herein by reference. In this method, single bilayered liposomes containing encapsulated active ingredients are prepared by dissolving the lipid component in an organic medium, injecting the organic solution of the lipid component under pressure into an aqueous component while simultaneously mixing the organic and aqueous components with a high speed homogenizer or mixing means, whereupon the liposomes are formed spontaneously.

The single bilayered liposomes containing the encapsulated hedgehog or ptc therapeutic can be employed directly or they can be employed in a suitable pharmaceutically acceptable carrier for topical administration. The viscosity of the liposomes can be increased by the addition of one or more suitable thickening agents such as, for example xanthan gum, hydroxypropyl cellulose, hydroxypropyl methylcellulose and mixtures thereof. The aqueous component may consist of water alone or it may contain electrolytes, buffered systems and other ingredients, such as, for example, preservatives. Suitable electrolytes which can be employed include metal salts such as alkali metal and alkaline earth metal salts. The preferred metal salts are calcium chloride, sodium chloride and potassium chloride. The concentration of the electrolyte may vary from zero to 260 mM, preferably from 5 mM to 160 mM. The aqueous component is placed in a suitable vessel which can be adapted to effect homogenization by effecting great turbulence during the injection of the organic component. Homogenization of the two components can be accomplished within the vessel, or, alternatively, the aqueous and organic components may be injected separately into a mixing means which is located outside the vessel. In the latter case, the

liposomes are formed in the mixing means and then transferred to another vessel for collection purpose.

The organic component consists of a suitable non-toxic, pharmaceutically acceptable solvent such as, for example ethanol, glycerol, propylene glycol and polyethylene glycol, and a suitable phospholipid which is soluble in the solvent. Suitable phospholipids which can be employed include lecithin, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, phosphatidylinositol, lysophosphatidylcholine and phosphatidyl glycerol, for example. Other lipophilic additives may be employed in order to selectively modify the characteristics of the liposomes. Examples of such other additives include stearylamine, phosphatidic acid, tocopherol, cholesterol and lanolin extracts.

In addition, other ingredients which can prevent oxidation of the phospholipids may be added to the organic component. Examples of such other ingredients include tocopherol, butylated hydroxyanisole, butylated hydroxytoluene, ascorbyl palmitate and ascorbyl oleate. Preservatives such as benzoic acid, methyl paraben and propyl paraben may also be added.

Apart from the above-described compositions, use may be made of covers, e.g. plasters, bandages, dressings, gauze pads and the like, containing an appropriate amount of a hedgehog or ptc therapeutic. In some cases use may be made of plasters, bandages, dressings, gauze pads and the like which have been impregnated with a topical formulation containing the therapeutic formulation.

Exemplification

The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1: Evaluation of the neuroprotective action sonic hedge hog in a cisplatin-induced neuropathy

The use of antiviral or anticancer chemotherapy may induce a severe neuropathy, that implies a reduction of the dosage used and hence a risk of unsucccess of the treatment. For example cisplatin is largely used for the treatment of tumors of the bladder, testis or ovary; however the dosage is limited because of the appearance of a partially irreversible toxic neuropathy, with a preference for the sensory fibers of large diameter that modifies the

proprioceptive sensitivity (Mollman, 1990). However there is presently no real treatment to cure or prevent such neurotoxicity.

It should be noted that NGF has been shown to be able to limit the importance of neuropathies induced by such chemotherapeutic agents (Apfel et al, 1991, Apfel et al, 1992). Two other peptides (NT3 and an ACTH analog) have also been tested in a similar model (Gao et al, 1995; Hamers et al, 1993). sonic hedge hog has been implicated in antero-posterior patterning of the developing chick limb (Riddle et al, 1993) and in motor neurons differentiation (Roelink et al, 1995). The present study was performed in order to measure the effect of Sonic Hedge Hog (SHH) as protective with regard to cisplatin-induced neuropathy. Behavioral and EMG measurements showed that SHH efficiently protected peripheral neurons against neuropathy, particularly at the highest concentration tested (500 ug/kg).

1) Materials and Methods

1.1) Animals housing and treatment

Thirty nine mice were included in this study and divided into 4 groups of 9-10 mice 38-40 g at onset; one group was treated with SHH (50 ug/kg, s.c.) 3 times per week; the second group received a dosage of 500 ug/kg; a third group was a vehicle group. These three groups were also treated with cisplatin (as described below). A fourth group was a control group without cisplatin administration but treated with 500 ug/kg SHH (control 500). Stock solution SHH (2.8mg/ml) was stored frozen at -70°C; on the day of use a vial was diluted to 0.2mg/ml with PBS and protein was mixed gently by pipetting. The animals were housed in plastic cages at room temperature in a 12: 12 h light-dark cycle. The mice had free access to food and water.

Animals were weighted once weekly and checked for their general behavior walking attitude and general outlook. Electromyographical and behavioral tests were also performed once weekly.

1.2) Cisplatin administration

Cisplatin was administered as an aqueous solution (1 mg/ml) at a dosage of 2 mg/kg i.p once daily during 14 consecutive days (cumulative dose). In order to avoid an important loss of weight of the animals, a Ringer-lactate solution was administered daily (0.4 ml/day i.p).

1.3) Behavioral testing

1.3.1) Pain threshold measurement

1.3.1.1) Tail flick test

The tail of the mouse was placed under a shutter-controlled lamp as a heat source. The latency before the mouse flicked its tail from the heat was recorded. A sensory alteration would increase the latency to flick.

1.3.1.2) Hot plate test

The animal was placed inside a glass cylinder of 17 cm height and 9 cm diameter on a hot plate at 52°C. The animal's behavior was observed, particularly the licking of a foot, the jump in the cylinder and the adjusted leap. The latency before licking its foot or before jumping to escape the heat was recorded. If the thermal sensitivity was altered, the time needed to feel the pain would be increased.

1.3.2) Motor coordination measurement

1.3.2.1) Rotarod test

The ability of an animal to stay on a rotating dowel (rotarod) is a good mean to measure the motor coordination and the proprioceptive sensitivity. The apparatus consisted of a rod, 1 cm in diameter, which turned at 12 rpm. The mice were tested for their ability to balance on the rotating bar during 180 sec maximum time (Tilson and Mitchell, 1984).

1.3.2.2) Walk on a rod

The animals were placed on a rod 1.5 cm in diameter and 40 cm long, that was situated horizontally at 50 cm over the floor; they were placed at one extremity and tended to reach the other end, that consisted of a wooden platform. The time needed to reach the platform was related to the motor coordination: the longest it was, the most important the motor deficit.

1.3.3) Muscle performance measurement

1.3.3.1) Muscular endurance

The muscular strength was evaluated by measuring the ability of an animal to hold a weight of 32 grams when it was lifted by the tail. The animal was allowed to use either two or the four legs. The time during which it held the weight was recorded, with a maximum of 60 sec, and reflected the muscular endurance.

1.3.3.2) Maximal strength

The maximal muscle strength was measured with an isometric transducer attached to a piece of wire. When the animal held the wire with either two or the four legs, it was slowly

moved backwards until it released the wire. The transducer measured the maximal strength; results are given in newton.

1.4) Electrophysiological measurement

Sensitive evoked response: Sensory nerve conduction velocity (SNCV).

5 Animals were anaesthetized with ketamine chlorhydrate (Ketalar) and diazepam (Valium) (1 ml/kg of a solution containing 11.25 mg ketalar and 0.375 mg of valium; i.p). Electrophysiological recordings were performed using a Neuromatic electromyogram (EMG) apparatus (Dantec, Les Ulis, France). Mice were deeply anaesthetized and normal body temperature maintained with a heating lamp.

10 The sensitive evoked response was measured in the caudal nerve. Stimulation of the caudal nerve was performed at the base of the tail, with two electrodes (one active, one reference) separated by 3mm; a unipolar recording needle was placed in a proximal site at approximately 40 mm. Sensory nerve velocity was recorded according to orthodromic conduction (from the tip of tail to the base). A ground needle electrode was inserted between the stimulating and recording electrode needles. The SNCV was calculated according to the distance between the two active electrodes.

1.5) Statistical studies.

20 The Electrophysiological and behavioral data were statistically analyzed by an analysis of variance with repeated measures (ANOVA). Following these analysis, a Scheffe's post hoc test was used to check for differences between individual groups.

2) Results

2.1) General survey

25 General behavior of animals was normal during the initial 2 weeks of study; however locomotor activity decreased while neuropathy was progressing, hair color changed and finally animals were almost immobile in their cages. Weight decreased strikingly after 2 weeks and remained low in vehicle group until 5 weeks. (Fig. 1; difference between treatments significant at $p<0.0001$; correlation between treatment effect and time changes significant at $p<0.0001$). However weight of animals treated with SHH (at both concentrations) increased immediately after the end of cisplatin administration and was almost normal at the end of study. In vehicle group, weight only started to increase at 5 weeks and was significantly below normal value at the

30 end of study.

As a consequence of cisplatin toxicity, some animals died during the study, starting at 3 weeks. However number of surviving animals was higher in SHH treated group, compare to

vehicle (Fig. 2). On the other hand, 3 controlSHH animals died during anaesthesia at 1 and 5 weeks.

2.2) EMG: Sensory nerve conduction velocity (SNCV)

According to EMG measurements, the neuropathy was found to appear after 1 week of cisplatin administration, was maximal at 3 weeks (delayed effect) and recover period went up to 8 weeks.

In standard conditions SNCV varied between 47 and 51 m/s for mice of 8 weeks of age. After cisplatin administration, SNCV decreased significantly in vehicle and SHH50 groups (Fig. 3; difference between treatments significant at $p < 0.0001$; correlation between treatment effect and time changes significant at $p < 0.0001$); recovery started immediately after end of cisplatin administration in SHH50 group, but was delayed one week later in vehicle group. Normal SNCV values were recovered after 8 weeks. However no significant decrease was found in SHH500 or control500 groups.

2.3) Behavioral testing

2.3.1) Pain threshold measurement

2.3.1.1) Tail flick test

Latency to flick the tail was increased after cisplatin administration in vehicle group, with a maximum at 4 weeks (Fig. 4; difference between treatments significant at $p < 0.0001$; correlation between treatment effect and time changes significant at $p < 0.0002$). A similar tendency was found in SHH50 group, but the curve was always below vehicle, i.e pain threshold defect was less important. In SHH50 group, latency increase was only transiently measured at 3 weeks.

2.3.1.2) Hot plate test

The latency before licking the paw did not vary much during the study, except a transient increase in vehicle group at 6 weeks (Fig. 5; difference between treatments not significant; correlation between treatment effect and time changes not significant). It should be noted that a great variation was found at that time and no significant difference was seen.

When pain was more important, mice tried to escape by jumping; the latency before first jump was recorded. It was found to be increased in vehicle group until 7 weeks and in SHH50 until 2 weeks (Fig. 6); the difference between treatments was only statistically significant at 6

weeks because of large variations in vehicle group (time course significant at $p<0.0001$; correlation between treatment effect and time changes not significant). A minor increase in SHH500 group was also measured until 3 weeks; values returned to normal thereafter and they were significantly lower than vehicle at 5 weeks.

- 5 After prolonged exposure to heat, mice escaped by jumping onto the rim of cylinder; some increase of the latency to escape was found at 2 weeks (particularly in SHH50 group) without reaching significance (Fig. 7). A greater increase was transiently found in vehicle group after 5 weeks and difference was statistically significant when compared to SHH treated groups (time course significant at $p<0.0001$; correlation between treatment effect and time changes
10 significant at $p<0.0001$).

2.3.2) Motor coordination measurement

2.3.2.1) Rotarod test

15 The ability of an animal to stay on a rotating rod was found to be significantly decreased in vehicle group, with a minimum performance at 3 weeks (Fig. 8). No decrease was measured in control500 or SHH500 groups and only a transient decrease at 2 weeks in SHH50 group (difference between treatments significant at $p<0.0001$; correlation between treatment effect and time changes significant at $p<0.0072$).

2.3.2.2) Walk on a rod

20 The time needed to walk on the rod in order to reach the platform significantly increased in vehicle group at 2 and 5 weeks, but only at 2 weeks in SHH50 group (Fig. 9; difference between treatments significant at $p<0.0015$; correlation between treatment effect and time changes significant at $p<0.0001$). No increase was found in SHH500 group, except at 3 weeks.

25 2.3.3) Muscle performance measurement

2.3.3.1) Muscular endurance

 When mice were allowed to use all 4 limbs to pull the wire, no decrease of muscular endurance was measured, except in vehicle group at 5 weeks (Fig. 10a; difference between treatments not significant; correlation between treatment effect and time changes not significant).

- 30 When mice were allowed to use only forelimbs to pull the wire, some decrease in muscular endurance was measured in vehicle group, but not in SHH50 or SHH500 groups (Fig. 10b; difference between treatments not significant; correlation between treatment effect and time

changes not significant). It should be noted that some decrease was also transiently found in control500 at 4 and 5 weeks.

2.3.3.2) Maximal strength

5 The maximal muscle strength exerted by the 4 limbs was decreased after 1-2 weeks in all cisplatin-treated groups (Fig. 11a; time course significant at $p<0.019$; correlation between treatment effect and time changes not significant). Recovery occurred at 5 weeks in SHH50 and SHH500 groups, but only at 7 weeks in vehicle group. No decrease was found in control.

10 The maximal muscle strength exerted by the forelimbs progressively decreased in vehicle group, with a minimum value at 6 weeks and recovery at 7 weeks (Fig. 11b; difference between treatments significant at $p<0.014$; correlation between treatment effect and time changes significant at $p<0.005$). A transient (and not significant decrease) was found in SHH50 at 2 weeks and no decrease was measured in SHH500 or control500 groups.

3) Discussion

5 The results obtained in the present study show that SHH was able to protect peripheral nerve against neuropathy induced by cisplatin, particularly at the highest concentration. The most striking effect was observed on SNCV, where no decrease was noticed in SHH500 group. In SHH50 group, SNCV decrease similar to vehicle was measured at 2 weeks; however recovery already began at 3 weeks, i.e one week earlier than in vehicle group. Similarly sensory defect is shown with tail flick test in vehicle group that lasted throughout the study while it was only transient in SHH500 (at 3 weeks). Sensory defect measured on the hot plate (first jump) was found until week 5 in vehicle group and week 2 in SHH50. No significant defect was measured in SHH500 group. Proprioceptive defect is also suggested by rotarod data in vehicle group until week 7 and transiently in SHH50 at week 2. No defect was found in SHH500 group. However
25 these changes may also be related to alteration of motor coordination.

Initial sensory neuropathy is known to extend towards motor impairment in patients treated with cisplatin. Similarly in the present study, muscle performance was impaired in the forelimbs endurance test in vehicle group, but not in any SHH group. Maximal muscle strength exerted by the 5 limbs was decreased in vehicle and both SHH groups, but recovery of function
30 occurred earlier in SHH groups. No such decrease was found in the forelimbs maximal strength test in SHH500 group.

Weight variation is a good indicator of general metabolism of the animals. It decreased strikingly at 2 weeks following cisplatin administration and lasted until week 5 in vehicle group;

in both SHH groups recovery occurred immediately after the end of cisplatin administration. Similarly animal survival was improved by SHH treatment.

It is concluded that SHH treatment with 500 ug/kg avoids neuropathy impairment in most tests or accelerates recovery when some defect is measured. Treatment with 50 ug/kg does not protect to the same extent, but also improves recovery (SNCV, jump, rotarod, muscle strength). Difference in time course of recovery is 2 weeks or more, when compared to vehicle group. These effects are similar to those observed with NGF or ACTH analog treatment in a similar paradigm (Apfel et al, 1992; Hamers et al, 1993); recovery of weight loss and SNCV decrease were also observed after end of cisplatin treatment. Dosage of ACTH was similar (75 ug/kg s.c every 48h), while amount of NGF was 10 times higher (5 mg/kg 3 times per week) and 1 mg/kg had no effect.

It should be noted that naive animals treated with 500 ug/kg SHH (but without cisplatin) did not show any impairment, except in forelimbs endurance. However as mentioned 3 animals of this group died during anaesthesia, at 1 and 5 weeks. Taken together with the absence of other signs of impairment in this group, it is most improbable that this occurrence may be due to toxicity of prolonged administration of the compound. However a similar study with lower dosage (100 or 200 ug/kg SHH) may be useful.

5) References for Example 1

Apfel S.C, Arezzo J.C, Lipson L.A and Kessler J.A, NGF prevents experimental cisplatin neuropathy, *Ann Neurol* (1992) 31, 76-80

Apfel S.C Lipton R.B, Arezzo J.C and Kessler J.A, NGF prevents toxic neuropathy in mice, *Ann Neurol* (1991) 29, 87-90

Gao WQ, Dybdal N, Shinsky N et al, *Ann Neurol* (1995) 38, 30-37

Hamers FPT, Pette C, Bravenboer B, Vecht CJ, Neujt JO and Gispen WH, *Cancer Chemother Pharmacol* (1993) 32, 162-166

Lipton R.B Apfel S.C and Dutcher J.P, *Neurology* (1989) 39, 368-373

Mollman J.E, *N.England J. Med* (1990) 322, 126-127

Riddle RD, Johnson RL, Laufer E and Tabin C, Sonic hedgehog mediates the polarizing activity of the ZPA, *Cell* 75 (1993) 1401-16.

Roelink H, Porter JA et al, Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis, *Cell* 81 (1995) 445-55.

Tilson H.A and Mitchell C.L, Neurobehavioral techniques of chemicals on the nervous system, Ann Rev Pharm Toxicol (1984) 24, 425-450.

Example 2: Evaluation of peripherhal nerves in normal and transgenic Dhh knockout mice.

5 We also undertook a comparison of the electrophysiology and morphology of peripheral nerve cells and bundles in normal mice and in transgenic mice in which the Dhh gene has been disrupted (the "Dhh^{-/-}" phenotype).

10 Adult mice were anesthetized with 0.5cc of ketamine/xylazine (diluted 1:10 with sterile saline) delivered by i.p. injection. The hair over the hindlimbs was shaved and the legs were taped in an extended position. Their core temperature was maintained at 38oC with an infrared lamp. A pair of surface recording electrodes were placed on the bottom of each foot; one over the intrinsic plantar muscles, the other more distally. The sciatic nerve was stimulated both proximally (at the level of the L5 vertebrae) and distally (the tibial nerve was stimulated at the ankle) with a pair of subcutaneous electrodes using a Dantec Neuromatic 2000. The stimulus strength was gradually increased until a maximal compound muscle action potential was obtained. The distance between the proximal and distal stimulation sites was measured and used to calculate the motor nerve conduction velocity.

15 Figure 12 illustrates that motor neuron conductance velocities are diminished in the Dhh^{-/-} mice, e.g., showing a functional deficit in peripheral nerve of Dhh^{-/-} mice.

20 The morphology of the peripheral nerve bundles in these mice were also observed (compare Figure 13A with 13B, and 14A with 14B). The integrity of the epineurial and perineurial sheath was altered in the Dhh^{-/-} mice. In another line of experiments, we tested the ability of Shh and Dhh to alter the proliferation of perineurial cells. Based on BrdU incorporation, both hedgehog proteins were able to increase proliferation of perineurial cells, but Dhh was dramatically more effective.

25 In addition to suggesting a role for hedgehog gene products in peripheral neuropathies, the observation that hedgehog proteins can induce proliferation of perineurial cells suggests that antagonists of hedgehog activity may be useful in disorders marked by unwanted proliferation of perineurial cells. For instance, localized hypertrophic mononeuropathy (LHM) is a rare focal neuropathy associated with perineurial cell proliferation due to an undefined stimulus. Perineuriomas. Likewise, in leprous neuropathy, proliferation of perineurial cells can be implicated in the abnormal multilayered appearance of the perineurium. Antagonists of hedgehog signalling may therefor be useful to inhibit proliferation of perineurial cells in the treatment of such disorders.

30

Example 3: Evaluation of the neuroprotective action sonic hedgehog in a taxol-induced neuropathy

The use of antiviral or anticancer chemotherapy may induce a severe neuropathy, that implies reduction of the dosage used and enhances the risk of unsuccess of the treatment. For example, taxol is used for the treatment of ovarian cancer or melanoma ; however the dosage is limited because of the appearance of a sensory toxic neuropathy (Lipton et al. 1989). It should be noted that NGF has been shown to limit the importance of neuropathies induced by such chemotherapeutic agents. The present study was designed to investigate the potency of Shh to protect against taxol-induced neuropathy. As shown in Figures 16 and 17, Shh has positive effects on taxol-treated mice, e.g., enhancing their ability to walk the length of a long suspended rod and to stay on a rotating drum (the so-called rotorod). Both are measures of motor ability and coordination.

1) Animals

Sixty four 22-24 g male Swiss mice (IFFA-CREDO, L'Arbresle, France) were used in this study. They were housed in collective cages (4-5 per cage) and maintained in a room with controlled temperature (21-22°C) and light under a reversed 12-12 light-dark cycle (light on at 7 p.m.), with food and water available ad libitum. All experiments were carried out in accordance with institutional guidelines.

2) Pharmacological treatment

Taxol (Sigma, l'Isle d'Abeau, France) was diluted in saline using cremophor 10 % V/v (Sigma) (20 mg taxol, 1 ml cremophor, 9 ml saline), and administered intraperitoneally (IP) as a volume of 10 ml/kg at the dose of 20 mg/kg once daily during 7 consecutive days. Shh was supplied by Biogen (Cambridge, MA, USA). Stock solutions Shh (2 mg/ml and 0.2 mg/ml) were stored at -70°C. Shh and vehicle solutions were labeled as A, B or C in order to perform a double-blind study. On the day of use, vials containing Shh or vehicle (A, B or Q were diluted to 1/40 in saline (200 µl sample + 7.8 ml saline) and injected as a volume of 10 ml/kg. Shh (50 or 500 µg/kg) or saline was administered subcutaneously (SC) 3 times per week (n=16 mice per group). These 3 groups were also treated with taxol. A fourth group consisting of a control group received cremophor IP and saline SC (n = 16). Shh treatment started from the first day of taxol administration on and lasted for 2 weeks.

3) Behavioral testing

Sensorimotor tests were performed once a week for 3 weeks. These tests were always done one day before electrophysiological (EPG) recordings. Each group was divided in two subgroups (series 1 and 2). Series 1 was tested on Mondays for behavioral tests and on Tuesdays for EPG test, Series 2 was tested on Wednesdays for behavioral tests and on Thursdays for EPG analysis. Behavioral testing was performed on day 0 (baseline, before taxol intoxication), day 7 (after 6 days of taxol injection), and day 14 (6 days after taxol discontinuation). EPG measurements, were performed before taxol intoxication (day 1), on day 8 (one day after the last injection of taxol), and on day 15 (7 days after discontinuing taxol). The first injection of taxol was performed on day 1, immediately after EPG recording.

3.1) Motor coordination measurements

Walking test: The apparatus used was a rod of 1.5 cm diameter and 80 cm long, maintained horizontally 40 cm above a table. The rod was graduated starting in the middle (0 cm) towards the two ends (40 cm) allowing to measure the distance walked by the animal.

Animals were tested once each week. Three consecutive trials were performed. For each trial (60 s maximum), each mouse was placed in the middle of the rod and the time needed to walk the 40 cm distance was recorded. Should the animal fall down or be unable to walk the 40 cm distance, 60 s were credited. For each animal, the mean time of the 3 trials was calculated. This time reflects the motor coordination performance.

Rotarod test: The ability of an animal to remain on a rotating rod (rotarod) reflects motor coordination and proprioceptive sensitivity. The apparatus used was a 3 cm diameter automated rod (Bloseb, Paris, France) with 12 rotations per min.

Animals were tested once each week. The mouse was placed on the rotating rod, and the time it remained on rod was recorded (300 s maximum). If the animal falls before 300 s, an additional trial is performed (3 trials maximum).

3.2) Muscular power

Maximal strength: The maximal muscle strength was measured with an isometric dynamometer connected to a grid. Once the animal was holding the grid with either two or the four paws, it was slowly moved backwards until it released it. The dynamometer measured the

maximal strength developed ; results are given in N. Two trials per session were performed. The mean of both trials was calculated for each animal.

Muscular endurance: The muscular endurance was evaluated by measuring the time (maximum of 60 s) during which an animal, lifted by the tail, was able to hold a weight of 38 g.

5 The animal was allowed to use either two or the four paws. Two consecutive trials were performed. The mean of both trials was calculated.

3.3) Sensitivity tests

10 Tail flick test: The apparatus consisted of a shutter-controlled lamp as a heat source (Bioseb). Each weekly session consisted of two consecutive trials with an interval of about 1 min and the mean was calculated.

Example 4: Evaluation of the neuroprotective action sonic hedge hog on spinal motor neurons

15 Amyotrophic lateral sclerosis (ALS) is a progressive neurodegenerative disorder primarily involving motoneurons (Ripps and al., 1995). Overexpression of a mutated human superoxide dismutase gene in mice causes a progressive paralytic disease as result of motoneuron loss in lumbar spinal cord (Mohajeri and al., 1998). The SOD1-G93A transgenic mouse model, used for preclinical drug studies in ALS (Gurney, 1997 ; Morrison and al., 1996),
20 is a good model to explore etiological mechanisms and to screen potential therapeutics. The present experiment, the results of which are illustrated in Figures 18-21, demonstrate the positive effects of hedgehog treatment on the survival of spinal motoneurons in SOD transgenic mice, a mouse model of ALS.

25 With intent to complete a study which analyses the effect of SHH by electromyographical and sensorimotor tests on progressive motoneurons degeneration in transgenic mice overexpressing mutant human superoxide dismutase, nervous tissue was harvested and histological studies performed on lumbar spinal cord sections.

1) Animals and treatment

30 SOD mice were genotyped by polymerized chain reaction (PCR) amplification of DNA extracted from the tail, 30 days after birth. DNA (10 ng) was added to 50 µl of mix reaction containing MgCl₂ and deoxynucleotide triphosphate mixture. The reaction uses primer

sequences set b for exon 4 described by Rosen and al. (1993) that hybridize to opposite strands and flank the target DNA sequence that is to be amplified using a GeneAmp PCR 2400 thermal cyclor (Perkin-Elmer, USA). The elongation of the primers is catalyzed by Tag DNA Polymerase (Appligene Oncor, France), a heat-stable DNA polymerase. A repetitive series of 30 cycles involving template denaturation (20 seconds at 92°C, primer annealing (20 seconds at 60°C) and extension of the annealed primers (20 seconds at 72°C) by Tag DNA Polymerase results in exponential accumulation of a specific DNA fragment. The resulting PCR products were electrophoresed on an 2% agarose gel and visualized with ethidium bromide (Sigma, L'Isle d'Abeau, France).

Twelve transgenic G93A heterozygotes mice (6 males and 6 females) were included in the study and were divided into 2 groups of 6 mice. One group was treated with vehicle and the olier with SHH at 500 µg/kg of body weight. They were housed in plastic cages and had free access to food and water. The local was maintained at a constant temperature of 22°C and humidity of 55% under conventional conditions and on a 12h light / 12h dark photocycle (light on 7 p.m.).

SHH was administered subcutaneously (SC) 3 times per week starting at 60 days of age, until 100 days.

2) Tissue harvesting and staining

Mice at 100 days of age were anaesthetized with 60 mg/kg ketamine hydrochloride (Ketalar) and 2 mg/kg diazepam. (Valium). They were perfused transcardially with phosphate-buffered saline (PBS) containing 0.1% heparin (Sigma, L'Isle d'Abeau, France). Then, animals were perfused with 4% paraformaldehyde in PBS until they became rigid. Spinal, cord was harvested and postfixed overnight. Tissue was then placed in 30% sucrose (Sigma, L'Isle d'Abeau, France) at 4°C until use.

Spinal cord was frozen in cold isopentane (Prolabo, Fontenay-sous-bois, France), embedded with Tissue-tek O.C.T. compound (Miles, USA) and sections (thickness: 30 gm) were made with a cryostat (Leica Jung CM 1800, Rueil-Malmaison, France). The sections were stained with a 0. 1% aqueous solution of cresyl violet (Sigma, L'Isle d'Abeau, France) for 30 to 45 seconds and then dehydrated and mounted in Eukitt (0. Kindler GmbH and Co., Freiburg). Only sections from lumbar segment were examined and to avoid the possibility of a given neuron being counted twice in two contiguous sections, only series of one section out of two were collected. Twenty seven to thirty one sections were obtained from a given lumbar segment.

Sections were observed using an optical microscope (Nikon, Japan). Results are expressed as the mean number of cells per animal counted in ventral horns on both sides.

3) Statistical analysis

5 Values are given as mean \pm s.e.mean. Differences between control group and SHH 500 group were evaluated by one factor ANOVA test using Statview Student vl.O VF software.

4) Results

10 Figure 18 shows that the group treated with SHH at the dose of 500 μ g/kg of body weight displayed a greater number of motoneurons than the control group, but difference was not significant [$F(1,10)=1.3$; N.S.]. It should be noted that in each group, the number of cells counted in the lumbar segment of the spinal cord of 1 mouse was much lower than the others (2Y0 for control group and 1Y0 for SHH 500 group, Table 1) and these mice were from the same littermate. It was therefore suggested to exclude these mice from the analysis. Figure 19 shows that without Y0 littermate, the number of cells counted was significantly different between control group and SHH 500 group, and that s.e.m. was much smaller. The number of cells in SHH 500 group was 15% higher than in the control group [$F(1,8)=13.7$; $p < 0.01$].

Table 1: Number of cells counted in each group (individual values)

	<u>Identification</u>	<u>Sex</u>	<u>Number of cells</u>
Control group	2W2.....	Female.....	874
	2Y0.....	Female.....	678
	1Z0.....	Male.....	920
	1Z2.....	Male.....	932
	2Z0.....	Female.....	835
	1A0.....	Male.....	851
	2T20.....	Female.....	1055
SHH 500 group	2T3.....	Female.....	928
	2T1.....	Female.....	1007
	1U1.....	Male.....	1111
	1U2.....	Male.....	985
	1Y0.....	Male.....	589

35 In order to further analyze data, motoneurons numbers measured in males and females were analyzed separately. Figure 20 shows that in males there was no statistical difference

between control group and SHH 500 group [$F(1,4)=0.0014$; N.S.]. However in females, the number of cells counted in SHH 500 group was significantly higher than in the control group [$F(1,4)=8.1$; $p < 0.05$] as shown in figure 21. These data suggest that SHH compound significantly improved motoneurons survival particularly in females mice.

- 5 The observation of individual data in control group shows that the number of cells counted in females, even not significantly, was lower than in males (795.7 ± 59.9 vs 901.0 ± 25.2). This difference may be explained by an earlier start of disease in females than in males. It may be interesting to measure the effects of SHH on motoneurons survival at later age and also to check if hormonal treatment may be able to synergies with SHH administration. In addition, it
10 may be important to begin SHH treatment earlier, as data suggest that neuromuscular impairments may already be present at 60 days.

5) References

- 15 GURNEY M.E. (1997). J Neurol Sci, 152 Suppl 1, S67-73.
 MORRISON et al. (1996) J Comp Neurol, 373:619-631.
 MOHAJERI et al. (1998) Exp Neurol, 150:329-336
 RIPPS et al. (1995) Proc Natl Acad Sci USA, 92:689-693.
 ROELINK et al. (1995) Cell 81:445-455.
20 ROSEN et la. (1993) Nature, 362:59-62.
 TANABE et al. (1995) Curr. Biol 5: 651-658.

Example 5: Evaluating actions of hedgehog proteins on galactose intoxication-mediated neuropathies

- 25 Galactose intoxication is a mean of inducing neuropathy and disrupting neurotrophic support to peripheral nerve cells in rats. Feeding rats diets high in galactose causes morphologic abnormalities in, e.g., Schwann cells and muscle that are accompanied by a neuropathy characterized by axonal atrophy and slowing nerve conduction velocities.

- Adapting a methodology set forth in Mizisin et al. (1997) J. Neuropath Exp Neurol 56:
30 1290-1301, the effects of hedgehog treatment on functional and structural disorders in nerves of galactosemic rats can be assessed.

As illustrated in Figure 23, treatment with Shh can improve nerve conductance in the galactose intoxicated animal.

Example 6: Evaluating the ability of treatment with hedgehog proteins to protect against diabetic neuropathy

In rats, i.p. injections of streptozotocine (STZ) can be used to generate an animal model of diabetic neuropathy. Utilizing such procedures as described in Garrett et al. (1997) Neurosci. Lett 222:191-194 the ability of hedgehog treatment to protect STZ-induced neuropathies can be assessed.

Example 7: Evaluating the effect of Hedgehog treatment on nerve crush injury

Hedgehog proteins improve functional recovery following sciatic nerve crush injury. Male CD-1 mice (25-30 g) were given a bilateral sciatic nerve crush and monitored daily for functional recovery by assessing their ability to grip a wire mesh with each hindfoot. See Figure 22. The data are expressed the average number of grip failures for the right and left foot in 10 trials. Mice were treated every other day beginning on the day of nerve crush with either vehicle (control group), pegylated isoleucine-isoleucine sonic hedgehog (Shh-PEG) at a dose of 1 mg/kg s.c. or isoleucine-isoleucine sonic hedgehog murine Ig fusion protein (Shh-Ig) at doses of 1 or 5 mg/kg s.c. The values represent the mean \pm S.E.M. for 14 mice per group. * $P < 0.05$ for all Shh groups compared to vehicle-treated control, Student-Newman-Keuls test.

All of the above-cited references and publications are hereby incorporated by reference.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific polypeptides, nucleic acids, methods, assays and reagents described herein. Such equivalents are considered to be within the scope of this invention.

SEQUENCE LISTING

5 (2) INFORMATION FOR SEQ ID NO:1:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1277 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1275

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GTC GAA ATG CTG CTG TTG ACA AGA ATT CTC TTG GTG GGC TTC ATC	48
Met Val Glu Met Leu Leu Leu Thr Arg Ile Leu Leu Val Gly Phe Ile	
1 5 10 15	
TGC GCT CTT TTA GTC TCC TCT GGG CTG ACT TGT GGA CCA GGC AGG GGC	96
Cys Ala Leu Leu Val Ser Ser Gly Leu Thr Cys Gly Pro Gly Arg Gly	
20 25 30	
ATT GGA AAA AGG AGG CAC CCC AAA AAG CTG ACC CCG TTA GCC TAT AAG	144
Ile Gly Lys Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys	
35 40 45	
CAG TTT ATT CCC AAT GTG GCA GAG AAG ACC CTA GGG GCC AGT GGA AGA	192
Gln Phe Ile Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Arg	
50 55 60	
TAT GAA GGG AAG ATC ACA AGA AAC TCC GAG AGA TTT AAA GAA CTA ACC	240
Tyr Glu Gly Lys Ile Thr Arg Asn Ser Glu Arg Phe Lys Glu Leu Thr	
65 70 75 80	
CCA AAT TAC AAC CCT GAC ATT ATT TTT AAG GAT GAA GAG AAC ACG GGA	288
Pro Asn Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Thr Gly	
85 90 95	
GCT GAC AGA CTG ATG ACT CAG CGC TGC AAG GAC AAG CTG AAT GCC CTG	336
Ala Asp Arg Leu Met Thr Gln Arg Cys Lys Asp Lys Leu Asn Ala Leu	
100 105 110	
GCG ATC TCG GTG ATG AAC CAG TGG CCC GGG GTG AAG CTG CGG GTG ACC	384
Ala Ile Ser Val Met Asn Gln Trp Pro Gly Val Lys Leu Arg Val Thr	
115 120 125	
GAG GGC TGG GAC GAG GAT GGC CAT CAC TCC GAG GAA TCG CTG CAC TAC	432
Glu Gly Trp Asp Glu Asp Gly His His Ser Glu Glu Ser Leu His Tyr	
130 135 140	
GAG GGT CGC GCC GTG GAC ATC ACC ACG TCG GAT CGG GAC CGC AGC AAG	480
Glu Gly Arg Ala Val Asp Ile Thr Thr Ser Asp Arg Asp Arg Ser Lys	
145 150 155 160	

		TAC	GGA	ATG	CTG	GCC	CGC	CTC	GCC	GTC	GAG	GCC	GGC	TTC	GAC	TGG	GTC	528
		Tyr	Gly	Met	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	
					165						170					175		
5		TAC	TAC	GAG	TCC	AAG	GCG	CAC	ATC	CAC	TGC	TCC	GTC	AAA	GCA	GAA	AAC	576
		Tyr	Tyr	Glu	Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn	
					180					185					190			
10		TCA	GTG	GCA	GCG	AAA	TCA	GGA	GGC	TGC	TTC	CCT	GGC	TCA	GCC	ACA	GTG	624
		Ser	Val	Ala	Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Thr	Val	
				195					200					205				
15		CAC	CTG	GAG	CAT	GGA	GGC	ACC	AAG	CTG	GTG	AAG	GAC	CTG	AGC	CCT	GGG	672
		His	Leu	Glu	His	Gly	Gly	Thr	Lys	Leu	Val	Lys	Asp	Leu	Ser	Pro	Gly	
			210					215					220					
20		GAC	CGC	GTG	CTG	GCT	GCT	GAC	GCG	GAC	GGC	CGG	CTG	CTC	TAC	AGT	GAC	720
		Asp	Arg	Val	Leu	Ala	Ala	Asp	Ala	Asp	Gly	Arg	Leu	Leu	Tyr	Ser	Asp	
			225			230					235						240	
		TTC	CTC	ACC	TTC	CTC	GAC	CGG	ATG	GAC	AGC	TCC	CGA	AAG	CTC	TTC	TAC	768
		Phe	Leu	Thr	Phe	Leu	Asp	Arg	Met	Asp	Ser	Ser	Arg	Lys	Leu	Phe	Tyr	
					245						250					255		
25		GTC	ATC	GAG	ACG	CGG	CAG	CCC	CGG	GCC	CGG	CTG	CTA	CTG	ACG	GCG	GCC	816
		Val	Ile	Glu	Thr	Arg	Gln	Pro	Arg	Ala	Arg	Leu	Leu	Leu	Thr	Ala	Ala	
					260					265					270			
30		CAC	CTG	CTC	TTT	GTG	GCC	CCC	CAG	CAC	AAC	CAG	TCG	GAG	GCC	ACA	GGG	864
		His	Leu	Leu	Phe	Val	Ala	Pro	Gln	His	Asn	Gln	Ser	Glu	Ala	Thr	Gly	
					275				280					285				
35		TCC	ACC	AGT	GGC	CAG	GCG	CTC	TTC	GCC	AGC	AAC	GTG	AAG	CCT	GGC	CAA	912
		Ser	Thr	Ser	Gly	Gln	Ala	Leu	Phe	Ala	Ser	Asn	Val	Lys	Pro	Gly	Gln	
			290					295					300					
40		CGT	GTC	TAT	GTG	CTG	GGC	GAG	GGC	GGG	CAG	CAG	CTG	CTG	CCG	GCG	TCT	960
		Arg	Val	Tyr	Val	Leu	Gly	Glu	Gly	Gly	Gln	Gln	Leu	Leu	Pro	Ala	Ser	
			305			310					315						320	
45		GTC	CAC	AGC	GTC	TCA	TTG	CGG	GAG	GAG	GCG	TCC	GGA	GCC	TAC	GCC	CCA	1008
		Val	His	Ser	Val	Ser	Leu	Arg	Glu	Glu	Ala	Ser	Gly	Ala	Tyr	Ala	Pro	
					325						330					335		
50		CTC	ACC	GCC	CAG	GGC	ACC	ATC	CTC	ATC	AAC	CGG	GTG	TTG	GCC	TCC	TGC	1056
		Leu	Thr	Ala	Gln	Gly	Thr	Ile	Leu	Ile	Asn	Arg	Val	Leu	Ala	Ser	Cys	
					340					345					350			
55		TAC	GCC	GTC	ATC	GAG	GAG	CAC	AGT	TGG	GCC	CAT	TGG	GCC	TTC	GCA	CCA	1104
		Tyr	Ala	Val	Ile	Glu	Glu	His	Ser	Trp	Ala	His	Trp	Ala	Phe	Ala	Pro	
				355					360					365				
60		TTC	CGC	TTG	GCT	CAG	GGG	CTG	CTG	GCC	GCC	CTC	TGC	CCA	GAT	GGG	GCC	1152
		Phe	Arg	Leu	Ala	Gln	Gly	Leu	Leu	Ala	Ala	Leu	Cys	Pro	Asp	Gly	Ala	
			370					375					380					
65		ATC	CCT	ACT	GCC	GCC	ACC	ACC	ACC	ACT	GGC	ATC	CAT	TGG	TAC	TCA	CGG	1200
		Ile	Pro	Thr	Ala	Ala	Thr	Thr	Thr	Thr	Gly	Ile	His	Trp	Tyr	Ser	Arg	
			385			390					395						400	
70		CTC	CTC	TAC	CGC	ATC	GGC	AGC	TGG	GTG	CTG	GAT	GGT	GAC	GCG	CTG	CAT	1248
		Leu	Leu	Tyr	Arg	Ile	Gly	Ser	Trp	Val	Leu	Asp	Gly	Asp	Ala	Leu	His	

	405	410	415	
	CCG CTG GGC ATG GTG GCA CCG GCC AGC TG			1277
5	Pro Leu Gly Met Val Ala Pro Ala Ser			
	420	425		

(2) INFORMATION FOR SEQ ID NO:2:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1190 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

20 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1191

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	ATG GCT CTG CCG GCC AGT CTG TTG CCC CTG TGC TGC TTG GCA CTC TTG	48
	Met Ala Leu Pro Ala Ser Leu Leu Pro Leu Cys Cys Leu Ala Leu Leu	
	1 5 10 15	
30	GCA CTA TCT GCC CAG AGC TGC GGG CCG GGC CGA GGA CCG GTT GGC CGG	96
	Ala Leu Ser Ala Gln Ser Cys Gly Pro Gly Arg Gly Pro Val Gly Arg	
	20 25 30	
35	CGG CGT TAT GTG CGC AAG CAA CTT GTG CCT CTG CTA TAC AAG CAG TTT	144
	Arg Arg Tyr Val Arg Lys Gln Leu Val Pro Leu Leu Tyr Lys Gln Phe	
	35 40 45	
40	GTG CCC AGT ATG CCC GAG CGG ACC CTG GGC GCG AGT GGG CCA GCG GAG	192
	Val Pro Ser Met Pro Glu Arg Thr Leu Gly Ala Ser Gly Pro Ala Glu	
	50 55 60	
45	GGG AGG GTA ACA AGG GGG TCG GAG CGC TTC CGG GAC CTC GTA CCC AAC	240
	Gly Arg Val Thr Arg Gly Ser Glu Arg Phe Arg Asp Leu Val Pro Asn	
	65 70 75 80	
	TAC AAC CCC GAC ATA ATC TTC AAG GAT GAG GAG AAC AGC GGC GCA GAC	288
	Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Ser Gly Ala Asp	
	85 90 95	
50	CGC CTG ATG ACA GAG CGT TGC AAA GAG CGG GTG AAC GCT CTA GCC ATC	336
	Arg Leu Met Thr Glu Arg Cys Lys Glu Arg Val Asn Ala Leu Ala Ile	
	100 105 110	
55	GCG GTG ATG AAC ATG TGG CCC GGA GTA CGC CTA CGT GTG ACT GAA GGC	384
	Ala Val Met Asn Met Trp Pro Gly Val Arg Leu Arg Val Thr Glu Gly	
	115 120 125	
60	TGG GAC GAG GAC GGC CAC CAC GCA CAG GAT TCA CTC CAC TAC GAA GGC	432
	Trp Asp Glu Asp Gly His His Ala Gln Asp Ser Leu His Tyr Glu Gly	
	130 135 140	

5	CGT Arg 145	GCC Ala	TTG Leu	GAC Asp	ATC Ile	ACC Thr 150	ACG Thr	TCT Ser	GAC Asp	CGT Arg	GAC Asp 155	CGT Arg	AAT Asn	AAG Lys	TAT Tyr	GGT Gly 160	480
	TTG Leu	TTG Leu	GCG Ala	CGC Arg	CTA Leu 165	GCT Ala	GTG Val	GAA Glu	GCC Ala	GGA Gly 170	TTC Phe	GAC Asp	TGG Trp	GTC Val	TAC Tyr 175	TAC Tyr	528
10	GAG Glu	TCC Ser	CGC Arg	AAC Asn 180	CAC His	ATC Ile	CAC His	GTA Val	TCG Ser 185	GTC Val	AAA Lys	GCT Ala	GAT Asp	AAC Asn 190	TCA Ser	CTG Leu	576
15	GCG Ala	GTC Val	CGA Arg 195	GCC Ala	GGA Gly	GGC Gly	TGC Cys 200	TTT Phe	CCG Pro	GGA Gly	AAT Asn	GCC Ala	ACG Thr 205	GTG Val	CGC Arg	TTG Leu	624
20	CGG Arg	AGC Ser 210	GGC Gly	GAA Glu	CGG Arg	AAG Lys	GGG Gly 215	CTG Leu	AGG Arg	GAA Glu	CTA Leu	CAT His 220	CGT Arg	GGT Gly	GAC Asp	TGG Trp	672
25	GTA Val 225	CTG Leu	GCC Ala	GCT Ala	GAT Asp 230	GCA Ala	GCG Ala	GGC Gly	CGA Arg	GTG Val	GTA Val 235	CCC Pro	ACG Thr	CCA Pro	GTG Val	CTG Leu 240	720
30	CTC Leu	TTC Phe	CTG Leu	GAC Asp 245	CGG Arg	GAT Asp	CTG Leu	CAG Gln	CGC Arg	CGC Arg 250	GCC Ala	TCG Ser	TTC Phe	GTG Val 255	GCT Ala	GTG Val	768
35	GAG Glu	ACC Thr	GAG Glu	CGG Arg 260	CCT Pro	CCG Pro	CGC Arg	AAA Lys 265	CTG Leu	TTG Leu	CTC Leu	ACA Thr	CCC Pro	TGG Trp 270	CAT His	CTG Leu	816
40	GTG Val	TTC Phe 275	GCT Ala	GCT Ala	CGC Arg	GGG Gly	CCA Pro 280	GCG Ala	CCT Pro	GCT Ala	CCA Pro	GGT Gly 285	GAC Asp	TTT Phe	GCA Ala	CCG Pro	864
45	GTG Val 290	TTC Phe	GCG Ala	CGC Arg	CGC Arg	TTA Leu	CGT Arg 295	GCT Ala	GGC Gly	GAC Asp	TCG Ser 300	GTG Val	CTG Leu	GCT Ala	CCC Pro	GGC Gly	912
50	GGG Gly 305	GAC Asp	GCG Ala	CTC Leu	CAG Gln 310	CCG Pro	GCG Ala	CGC Arg	GTA Val	GCC Ala 315	CGC Arg	GTG Val	GCG Ala	CGC Arg	GAG Glu 320	GAA Glu	960
55	GCC Ala	GTG Val	GGC Gly	GTG Val	TTC Phe 325	GCA Ala	CCG Pro	CTC Leu	ACT Thr 330	GCG Ala	CAC His 335	GGG Gly	ACG Thr	CTG Leu	CTG Leu 335	GTC Val	1008
60	AAC Asn	GAC Asp	GTC Val	CTC Leu 340	GCC Ala	TCC Ser	TGC Cys	TAC Tyr 345	GCG Ala	GTT Val	CTA Leu	GAG Glu	AGT Ser 350	CAC His	CAG Gln	TGG Trp	1056
65	GCC Ala	CAC His 355	CGC Arg	GCC Ala	TTC Phe	GCC Ala	CCT Pro 360	TTG Leu	CGG Arg 365	CTG Leu	CTG Leu	CAC His 370	GCG Ala 375	CTC Leu	GGG Gly	GCT Ala	1104
70	CTG Leu	CTC Leu 370	CCT Pro	GGG Gly	GGT Gly	GCA Ala 375	GTC Val 380	CAG Gln	CCG Pro	ACT Thr	GGC Gly 385	ATG Met 390	CAT His	TGG Trp	TAC Tyr	TCT Ser	1152

CGC	CTC	CTT	TAC	CGC	TTG	GCC	GAG	GAG	TTA	ATG	GGC	TG	1190
Arg	Leu	Leu	Tyr	Arg	Leu	Ala	Glu	Glu	Leu	Met	Gly		
385					390					395			

5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1281 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

15

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1233

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

25	ATG	TCT	CCC	GCC	TGG	CTC	CGG	CCC	CGA	CTG	CGG	TTC	TGT	CTG	TTC	CTG	48
	Met	Ser	Pro	Ala	Trp	Leu	Arg	Pro	Arg	Leu	Arg	Phe	Cys	Leu	Phe	Leu	
	1				5					10					15		
30	CTG	CTG	CTG	CTT	CTG	GTG	CCG	GCG	GCG	CGG	GGC	TGC	GGG	CCG	GGC	CGG	96
	Leu	Leu	Leu	Leu	Leu	Val	Pro	Ala	Ala	Arg	Gly	Cys	Gly	Pro	Gly	Arg	
				20					25					30			
35	GTG	GTG	GGC	AGC	CGC	CGG	AGG	CCG	CCT	CGC	AAG	CTC	GTG	CCT	CTT	GCC	144
	Val	Val	Gly	Ser	Arg	Arg	Arg	Pro	Pro	Arg	Lys	Leu	Val	Pro	Leu	Ala	
			35					40					45				
40	TAC	AAG	CAG	TTC	AGC	CCC	AAC	GTG	CCG	GAG	AAG	ACC	CTG	GGC	GCC	AGC	192
	Tyr	Lys	Gln	Phe	Ser	Pro	Asn	Val	Pro	Glu	Lys	Thr	Leu	Gly	Ala	Ser	
		50					55					60					
45	GGG	CGC	TAC	GAA	GGC	AAG	ATC	GCG	CGC	AGC	TCT	GAG	CGC	TTC	AAA	GAG	240
	Gly	Arg	Tyr	Glu	Gly	Lys	Ile	Ala	Arg	Ser	Ser	Glu	Arg	Phe	Lys	Glu	
	65				70					75						80	
50	CTC	ACC	CCC	AAC	TAC	AAT	CCC	GAC	ATC	ATC	TTC	AAG	GAC	GAG	GAG	AAC	288
	Leu	Thr	Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	
				85					90					95			
55	ACG	GGT	GCC	GAC	CGC	CTC	ATG	ACC	CAG	CGC	TGC	AAG	GAC	CGT	CTG	AAC	336
	Thr	Gly	Ala	Asp	Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Arg	Leu	Asn	
			100					105						110			
60	TCA	CTG	GCC	ATC	TCT	GTC	ATG	AAC	CAG	TGG	CCT	GGT	GTG	AAA	CTG	CGG	384
	Ser	Leu	Ala	Ile	Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Lys	Leu	Arg	
			115					120					125				
55	GTG	ACC	GAA	GGC	CGG	GAT	GAA	GAT	GGC	CAT	CAC	TCA	GAG	GAG	TCT	TTA	432
	Val	Thr	Glu	Gly	Arg	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	
		130					135					140					
60	CAC	TAT	GAG	GGC	CGC	GCG	GTG	GAT	ATC	ACC	ACC	TCA	GAC	CGT	GAC	CGA	480
	His	Tyr	Glu	Gly	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	

	145		150		155		160		
5	AAT AAG TAT GGA CTG CTG GCG CGC TTA GCA GTG GAG GCC GGC TTC GAC	528							
	Asn Lys Tyr Gly Leu Leu Ala Arg Leu Ala Val Glu Ala Gly Phe Asp								
10	TGG GTG TAT TAC GAG TCC AAG GCC CAC GTG CAT TGC TCT GTC AAG TCT	576							
	Trp Val Tyr Tyr Glu Ser Lys Ala His Val His Cys Ser Val Lys Ser								
15	GAG CAT TCG GCC GCT GCC AAG ACA GGT GGC TGC TTT CCT GCC GGA GCC	624							
	Glu His Ser Ala Ala Ala Lys Thr Gly Gly Cys Phe Pro Ala Gly Ala								
20	CAG GTG CGC CTA GAG AAC GGG GAG CGT GTG GCC CTG TCA GCT GTA AAG	672							
	Gln Val Arg Leu Glu Asn Gly Glu Arg Val Ala Leu Ser Ala Val Lys								
25	CCA GGA GAC CGG GTG CTG GCC ATG GGG GAG GAT GGG ACC CCC ACC TTC	720							
	Pro Gly Asp Arg Val Leu Ala Met Gly Glu Asp Gly Thr Pro Thr Phe								
30	AGT GAT GTG CTT ATT TTC CTG GAC CGC GAG CCA AAC CGG CTG AGA GCT	768							
	Ser Asp Val Leu Ile Phe Leu Asp Arg Glu Pro Asn Arg Leu Arg Ala								
35	TTC CAG GTC ATC GAG ACT CAG GAT CCT CCG CGT CGG CTG GCG CTC ACG	816							
	Phe Gln Val Ile Glu Thr Gln Asp Pro Pro Arg Arg Leu Ala Leu Thr								
40	CCT GCC CAC CTG CTC TTC ATT GCG GAC AAT CAT ACA GAA CCA GCA GCC	864							
	Pro Ala His Leu Leu Phe Ile Ala Asp Asn His Thr Glu Pro Ala Ala								
45	CAC TTC CGG GCC ACA TTT GCC AGC CAT GTG CAA CCA GGC CAA TAT GTG	912							
	His Phe Arg Ala Thr Phe Ala Ser His Val Gln Pro Gly Gln Tyr Val								
50	CTG GTA TCA GGG GTA CCA GGC CTC CAG CCT GCT CGG GTG GCA GCT GTC	960							
	Leu Val Ser Gly Val Pro Gly Leu Gln Pro Ala Arg Val Ala Ala Val								
55	TCC ACC CAC GTG GCC CTT GGG TCC TAT GCT CCT CTC ACA AGG CAT GGG	1008							
	Ser Thr His Val Ala Leu Gly Ser Tyr Ala Pro Leu Thr Arg His Gly								
60	ACA CTT GTG GTG GAG GAT GTG GTG GCC TCC TGC TTT GCA GCT GTG GCT	1056							
	Thr Leu Val Val Glu Asp Val Val Ala Ser Cys Phe Ala Ala Val Ala								
65	GAC CAC CAT CTG GCT CAG TTG GCC TTC TGG CCC CTG CGA CTG TTT CCC	1104							
	Asp His His Leu Ala Gln Leu Ala Phe Trp Pro Leu Arg Leu Phe Pro								
70	AGT TTG GCA TGG GGC AGC TGG ACC CCA AGT GAG GGT GTT CAC TCC TAC	1152							
	Ser Leu Ala Trp Gly Ser Trp Thr Pro Ser Glu Gly Val His Ser Tyr								
75	CCT CAG ATG CTC TAC CGC CTG GGG CGT CTC TTG CTA GAA GAG AGC ACC	1200							
	Pro Gln Met Leu Tyr Arg Leu Gly Arg Leu Leu Leu Glu Glu Ser Thr								

TTC CAT CCA CTG GGC ATG TCT GGG GCA GGA AGC TGAAGGGACT CTAACCACTG 1253
Phe His Pro Leu Gly Met Ser Gly Ala Gly Ser
405 410

CCCTCCTGGA ACTGCTGTGC GTGGATCC 1281

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1313 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION: 1..1314

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATG	CTG	CTG	CTG	CTG	GCC	AGA	TGT	TTT	CTG	GTG	ATC	CTT	GCT	TCC	TCG	48
Met	Leu	Leu	Leu	Leu	Ala	Arg	Cys	Phe	Leu	Val	Ile	Leu	Ala	Ser	Ser	
1				5				10						15		
CTG	CTG	GTG	TGC	CCC	GGG	CTG	GCC	TGT	GGG	CCC	GGC	AGG	GGG	TTT	GGA	96
Leu	Leu	Val	Cys	Pro	Gly	Leu	Ala	Cys	Gly	Pro	Gly	Arg	Gly	Phe	Gly	
			20					25					30			
AAG	AGG	CGG	CAC	CCC	AAA	AAG	CTG	ACC	CCT	TTA	GCC	TAC	AAG	CAG	TTT	144
Lys	Arg	Arg	His	Pro	Lys	Lys	Leu	Thr	Pro	Leu	Ala	Tyr	Lys	Gln	Phe	
		35					40					45				
ATT	CCC	AAC	GTA	GCC	GAG	AAG	ACC	CTA	GGG	GCC	AGC	GGC	AGA	TAT	GAA	192
Ile	Pro	Asn	Val	Ala	Glu	Lys	Thr	Leu	Gly	Ala	Ser	Gly	Arg	Tyr	Glu	
	50				55				60							
GGG	AAG	ATC	ACA	AGA	AAC	TCC	GAA	CGA	TTT	AAG	GAA	CTC	ACC	CCC	AAT	240
Gly	Lys	Ile	Thr	Arg	Asn	Ser	Glu	Arg	Phe	Lys	Glu	Leu	Thr	Pro	Asn	
65				70				75						80		
TAC	AAC	CCC	GAC	ATC	ATA	TTT	AAG	GAT	GAG	GAA	AAC	ACG	GGA	GCA	GAC	288
Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Thr	Gly	Ala	Asp	
			85					90					95			
CGG	CTG	ATG	ACT	CAG	AGG	TGC	AAA	GAC	AAG	TTA	AAT	GCC	TTG	GCC	ATC	336
Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Lys	Leu	Asn	Ala	Leu	Ala	Ile	
			100				105						110			
TCT	GTG	ATG	AAC	CAG	TGG	CCT	GGA	GTG	AGG	CTG	CGA	GTG	ACC	GAG	GGC	384
Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Arg	Leu	Arg	Val	Thr	Glu	Gly	
		115					120					125				
TGG	GAT	GAG	GAC	GGC	CAT	CAT	TCA	GAG	GAG	TCT	CTA	CAC	TAT	GAG	GGT	432
Trp	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	His	Tyr	Glu	Gly	
	130						135					140				

5	CGA	GCA	GTG	GAC	ATC	ACC	ACG	TCC	GAC	CGG	GAC	CGC	AGC	AAG	TAC	GGC	480
	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Ser	Lys	Tyr	Gly	145
10	ATG	CTG	GCT	CGC	CTG	GCT	GTG	GAA	GCA	GGT	TTC	GAC	TGG	GTC	TAC	TAT	528
	Met	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr	165
15	GAA	TCC	AAA	GCT	CAC	ATC	CAC	TGT	TCT	GTG	AAA	GCA	GAG	AAC	TCC	GTG	576
	Glu	Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn	Ser	Val	180
20	GCG	GCC	AAA	TCC	GGC	GGC	TGT	TTC	CCG	GGA	TCC	GCC	ACC	GTG	CAC	CTG	624
	Ala	Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Thr	Val	His	Leu	195
25	GAG	CAG	GGC	GGC	ACC	AAG	CTG	GTG	AAG	GAC	TTA	CGT	CCC	GGA	GAC	CGC	672
	Glu	Gln	Gly	Gly	Thr	Lys	Leu	Val	Lys	Asp	Leu	Arg	Pro	Gly	Asp	Arg	210
30	GTG	CTG	GCG	GCT	GAC	GAC	CAG	GGC	CGG	CTG	CTG	TAC	AGC	GAC	TTC	CTC	720
	Val	Leu	Ala	Ala	Asp	Asp	Gln	Gly	Arg	Leu	Leu	Tyr	Ser	Asp	Phe	Leu	225
35	ACC	TTC	CTG	GAC	CGC	GAC	GAA	GGC	GCC	AAG	AAG	GTC	TTC	TAC	GTG	ATC	768
	Thr	Phe	Leu	Asp	Arg	Asp	Glu	Gly	Ala	Lys	Lys	Val	Phe	Tyr	Val	Ile	245
40	GAG	ACG	CTG	GAG	CCG	CGC	GAG	CGC	CTG	CTG	CTC	ACC	GCC	GCG	CAC	CTG	816
	Glu	Thr	Leu	Glu	Pro	Arg	Glu	Arg	Leu	Leu	Leu	Thr	Ala	Ala	His	Leu	260
45	CTC	TTC	GTG	GCG	CCG	CAC	AAC	GAC	TCG	GGG	CCC	ACG	CCC	GGG	CCA	AGC	864
	Leu	Phe	Val	Ala	Pro	His	Asn	Asp	Ser	Gly	Pro	Thr	Pro	Gly	Pro	Ser	275
50	GCG	CTC	TTT	GCC	AGC	CGC	GTG	CGC	CCC	GGG	CAG	CGC	GTG	TAC	GTG	GTG	912
	Ala	Leu	Phe	Ala	Ser	Arg	Val	Arg	Pro	Gly	Gln	Arg	Val	Tyr	Val	Val	290
55	GCT	GAA	CGC	GGC	GGG	GAC	CGC	CGG	CTG	CTG	CCC	GCC	GCG	GTG	CAC	AGC	960
	Ala	Glu	Arg	Gly	Gly	Asp	Arg	Arg	Leu	Leu	Pro	Ala	Ala	Val	His	Ser	305
60	GTG	ACG	CTG	CGA	GAG	GAG	GAG	GCG	GGC	GCG	TAC	GCG	CCG	CTC	ACG	GCG	1008
	Val	Thr	Leu	Arg	Glu	Glu	Glu	Ala	Gly	Ala	Tyr	Ala	Pro	Leu	Thr	Ala	325
65	CAC	GGC	ACC	ATT	CTC	ATC	AAC	CGG	GTG	CTC	GCC	TCG	TGC	TAC	GCT	GTC	1056
	His	Gly	Thr	Ile	Leu	Ile	Asn	Arg	Val	Leu	Ala	Ser	Cys	Tyr	Ala	Val	340
70	ATC	GAG	GAG	CAC	AGC	TGG	GCA	CAC	CGG	GCC	TTC	GCG	CCT	TTC	CGC	CTG	1104
	Ile	Glu	Glu	His	Ser	Trp	Ala	His	Arg	Ala	Phe	Ala	Pro	Phe	Arg	Leu	355
75	GCG	CAC	GCG	CTG	CTG	GCC	GCG	CTG	GCA	CCC	GCC	CGC	ACG	GAC	GGC	GGG	1152
	Ala	His	Ala	Leu	Leu	Ala	Ala	Leu	Ala	Pro	Ala	Arg	Thr	Asp	Gly	Gly	370

	GGC GGG GGC AGC ATC CCT GCA GCG CAA TCT GCA ACG GAA GCG AGG GGC	1200
	Gly Gly Gly Ser Ile Pro Ala Ala Gln Ser Ala Thr Glu Ala Arg Gly	
	385 390 395 400	
5	GCG GAG CCG ACT GCG GGC ATC CAC TGG TAC TCG CAG CTG CTC TAC CAC	1248
	Ala Glu Pro Thr Ala Gly Ile His Trp Tyr Ser Gln Leu Leu Tyr His	
	405 410 415	
10	ATT GGC ACC TGG CTG TTG GAC AGC GAG ACC ATG CAT CCC TTG GGA ATG	1296
	Ile Gly Thr Trp Leu Leu Asp Ser Glu Thr Met His Pro Leu Gly Met	
	420 425 430	
	GCG GTC AAG TCC AGC TG	1313
15	Ala Val Lys Ser Ser	
	435	
(2) INFORMATION FOR SEQ ID NO:5:		
20	(i) SEQUENCE CHARACTERISTICS:	
	(A) LENGTH: 1256 base pairs	
	(B) TYPE: nucleic acid	
	(C) STRANDEDNESS: both	
	(D) TOPOLOGY: linear	
25	(ii) MOLECULE TYPE: cDNA	
	(ix) FEATURE:	
30	(A) NAME/KEY: CDS	
	(B) LOCATION: 1..1257	
35	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:	
	ATG CGG CTT TTG ACG AGA GTG CTG CTG GTG TCT CTT CTC ACT CTG TCC	48
	Met Arg Leu Leu Thr Arg Val Leu Leu Val Ser Leu Leu Thr Leu Ser	
	1 5 10 15	
40	TTG GTG GTG TCC GGA CTG GCC TGC GGT CCT GGC AGA GGC TAC GGC AGA	96
	Leu Val Val Ser Gly Leu Ala Cys Gly Pro Gly Arg Gly Tyr Gly Arg	
	20 25 30	
45	AGA AGA CAT CCG AAG AAG CTG ACA CCT CTC GCC TAC AAG CAG TTC ATA	144
	Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys Gln Phe Ile	
	35 40 45	
	CCT AAT GTC GCG GAG AAG ACC TTA GGG GCC AGC GGC AGA TAC GAG GGC	192
50	Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Arg Tyr Glu Gly	
	50 55 60	
	AAG ATA ACG CGC AAT TCG GAG AGA TTT AAA GAA CTT ACT CCA AAT TAC	240
	Lys Ile Thr Arg Asn Ser Glu Arg Phe Lys Glu Leu Thr Pro Asn Tyr	
	65 70 75 80	
55	AAT CCC GAC ATT ATC TTT AAG GAT GAG GAG AAC ACG GGA GCG GAC AGG	288
	Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Thr Gly Ala Asp Arg	
	85 90 95	
60	CTC ATG ACA CAG AGA TGC AAA GAC AAG CTG AAC TCG CTG GCC ATC TCT	336
	Leu Met Thr Gln Arg Cys Lys Asp Lys Leu Asn Ser Leu Ala Ile Ser	

5

10

15

20

25

30

35

40

45

50

22

	GCG CAT TTG GCC TTC GCG CCC GCC AGG CTC TAT TAT TAC GTG TCA TCA	1104
	Ala His Leu Ala Phe Ala Pro Ala Arg Leu Tyr Tyr Tyr Val Ser Ser	
	355 360 365	
5	TTC CTG TCC CCC AAA ACT CCA GCA GTC GGT CCA ATG CGA CTT TAC AAC	1152
	Phe Leu Ser Pro Lys Thr Pro Ala Val Gly Pro Met Arg Leu Tyr Asn	
	370 375 380	
10	AGG AGG GGG TCC ACT GGT ACT CCA GGC TCC TGT CAT CAA ATG GGA ACG	1200
	Arg Arg Gly Ser Thr Gly Thr Pro Gly Ser Cys His Gln Met Gly Thr	
	385 390 395 400	
15	TGG CTT TTG GAC AGC AAC ATG CTT CAT CCT TTG GGG ATG TCA GTA AAC	1248
	Trp Leu Leu Asp Ser Asn Met Leu His Pro Leu Gly Met Ser Val Asn	
	405 410 415	
20	TCA AGC TG	1256
	Ser Ser	

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1425 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1425

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

	ATG CTG CTG CTG GCG AGA TGT CTG CTG CTA GTC CTC GTC TCC TCG CTG	48
	Met Leu Leu Leu Ala Arg Cys Leu Leu Leu Val Leu Val Ser Ser Leu	
	1 5 10 15	
45	CTG GTA TGC TCG GGA CTG GCG TGC GGA CCG GGC AGG GGG TTC GGG AAG	96
	Leu Val Cys Ser Gly Leu Ala Cys Gly Pro Gly Arg Gly Phe Gly Lys	
	20 25 30	
50	AGG AGG CAC CCC AAA AAG CTG ACC CCT TTA GCC TAC AAG CAG TTT ATC	144
	Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys Gln Phe Ile	
	35 40 45	
55	CCC AAT GTG GCC GAG AAG ACC CTA GGC GCC AGC GGA AGG TAT GAA GGG	192
	Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Arg Tyr Glu Gly	
	50 55 60	
60	AAG ATC TCC AGA AAC TCC GAG CGA TTT AAG GAA CTC ACC CCC AAT TAC	240
	Lys Ile Ser Arg Asn Ser Glu Arg Phe Lys Glu Leu Thr Pro Asn Tyr	
	65 70 75 80	
	AAC CCC GAC ATC ATA TTT AAG GAT GAA GAA AAC ACC GGA GCG GAC AGG	288

5
10
15
20
25
30
35
40
45
50
55
60

	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Thr	Gly	Ala	Asp	Arg	
				85						90					95		
5	CTG	ATG	ACT	CAG	AGG	TGT	AAG	GAC	AAG	TTG	AAC	GCT	TTG	GCC	ATC	TCG	336
	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Lys	Leu	Asn	Ala	Leu	Ala	Ile	Ser	
				100					105					110			
10	GTG	ATG	AAC	CAG	TGG	CCA	GGA	GTG	AAA	CTG	CGG	GTG	ACC	GAG	GGC	TGG	384
	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Lys	Leu	Arg	Val	Thr	Glu	Gly	Trp	
			115					120					125				
15	GAC	GAA	GAT	GGC	CAC	CAC	TCA	GAG	GAG	TCT	CTG	CAC	TAC	GAG	GGC	CGC	432
	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	His	Tyr	Glu	Gly	Arg	
		130					135					140					
20	GCA	GTG	GAC	ATC	ACC	ACG	TCT	GAC	CGC	GAC	CGC	AGC	AAG	TAC	GGC	ATG	480
	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Ser	Lys	Tyr	Gly	Met	
		145				150				155						160	
25	CTG	GCC	CGC	CTG	GCG	GTG	GAG	GCC	GGC	TTC	GAC	TGG	GTG	TAC	TAC	GAG	528
	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr	Glu	
				165						170					175		
30	TCC	AAG	GCA	CAT	ATC	CAC	TGC	TCG	GTG	AAA	GCA	GAG	AAC	TCG	GTG	GCG	576
	Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn	Ser	Val	Ala	
				180					185					190			
35	GCC	AAA	TCG	GGA	GGC	TGC	TTC	CCG	GGC	TCG	GCC	ACG	GTG	CAC	CTG	GAG	624
	Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Thr	Val	His	Leu	Glu	
			195					200					205				
40	CAG	GGC	GGC	ACC	AAG	CTG	GTG	AAG	GAC	CTG	AGC	CCC	GGG	GAC	CGC	GTG	672
	Gln	Gly	Gly	Thr	Lys	Leu	Val	Lys	Asp	Leu	Ser	Pro	Gly	Asp	Arg	Val	
		210				215						220					
45	CTG	GCG	GCG	GAC	GAC	CAG	GGC	CGG	CTG	CTC	TAC	AGC	GAC	TTC	CTC	ACT	720
	Leu	Ala	Ala	Asp	Asp	Gln	Gly	Arg	Leu	Leu	Tyr	Ser	Asp	Phe	Leu	Thr	
		225				230				235						240	
50	TTC	CTG	GAC	CGC	GAC	GAC	GGC	GCC	AAG	AAG	GTC	TTC	TAC	GTG	ATC	GAG	768
	Phe	Leu	Asp	Arg	Asp	Asp	Gly	Ala	Lys	Lys	Val	Phe	Tyr	Val	Ile	Glu	
				245						250					255		
55	ACG	CGG	GAG	CCG	CGC	GAG	CGC	CTG	CTG	CTC	ACC	GCC	GCG	CAC	CTG	CTC	816
	Thr	Arg	Glu	Pro	Arg	Glu	Arg	Leu	Leu	Leu	Thr	Ala	Ala	His	Leu	Leu	
				260				265						270			
60	TTT	GTG	GCG	CCG	CAC	AAC	GAC	TCG	GCC	ACC	GGG	GAG	CCC	GAG	GCG	TCC	864
	Phe	Val	Ala	Pro	His	Asn	Asp	Ser	Ala	Thr	Gly	Glu	Pro	Glu	Ala	Ser	
			275					280					285				
65	TCG	GGC	TCG	GGG	CCG	CCT	TCC	GGG	GGC	GCA	CTG	GGG	CCT	CGG	GCG	CTG	912
	Ser	Gly	Ser	Gly	Pro	Pro	Ser	Gly	Gly	Ala	Leu	Gly	Pro	Arg	Ala	Leu	
		290				295						300					
70	TTC	GCC	AGC	CGC	GTG	CGC	CCG	GGC	CAG	CGC	GTG	TAC	GTG	GTG	GCC	GAG	960
	Phe	Ala	Ser	Arg	Val	Arg	Pro	Gly	Gln	Arg	Val	Tyr	Val	Val	Ala	Glu	
		305				310					315					320	
75	CGT	GAC	GGG	GAC	CGC	CGG	CTC	CTG	CCC	GCC	GCT	GTG	CAC	AGC	GTG	ACC	1008
	Arg	Asp	Gly	Asp	Arg	Arg	Leu	Leu	Pro	Ala	Ala	Val	His	Ser	Val	Thr	

	325	330	335			
5	CTA AGC GAG GAG GCC GCG GGC GCC TAC GCG CCG CTC ACG GCC CAG GGC Leu Ser Glu Glu Ala Ala Gly Ala Tyr Ala Pro Leu Thr Ala Gln Gly	340	345	350	1056	
10	ACC ATT CTC ATC AAC CGG GTG CTG GCC TCG TGC TAC GCG GTC ATC GAG Thr Ile Leu Ile Asn Arg Val Leu Ala Ser Cys Tyr Ala Val Ile Glu	355	360	365	1104	
15	GAG CAC AGC TGG GCG CAC CGG GCC TTC GCG CCC TTC CGC CTG GCG CAC Glu His Ser Trp Ala His Arg Ala Phe Ala Pro Phe Arg Leu Ala His	370	375	380	1152	
20	GCG CTC CTG GCT GCA CTG GCG CCC GCG CGC ACG GAC CGC GGC GGG GAC Ala Leu Leu Ala Ala Leu Ala Pro Ala Arg Thr Asp Arg Gly Gly Asp	385	390	395	400	1200
25	AGC GGC GGC GGG GAC CGC GGG GGC GGC GGC GGC AGA GTA GCC CTA ACC Ser Gly Gly Gly Asp Arg Gly Gly Gly Gly Gly Arg Val Ala Leu Thr	405	410	415	1248	
30	GCT CCA GGT GCT GCC GAC GCT CCG GGT GCG GGG GCC ACC GCG GGC ATC Ala Pro Gly Ala Ala Asp Ala Pro Gly Ala Gly Ala Thr Ala Gly Ile	420	425	430	1296	
35	CAC TGG TAC TCG CAG CTG CTC TAC CAA ATA GGC ACC TGG CTC CTG GAC His Trp Tyr Ser Gln Leu Leu Tyr Gln Ile Gly Thr Trp Leu Leu Asp	435	440	445	1344	
40	AGC GAG GCC CTG CAC CCG CTG GGC ATG GCG GTC AAG TCC AGC NNN AGC Ser Glu Ala Leu His Pro Leu Gly Met Ala Val Lys Ser Ser Xaa Ser	450	455	460	1392	
45	CGG GGG GCC GGG GGA GGG GCG CGG GAG GGG GCC Arg Gly Ala Gly Gly Gly Ala Arg Glu Gly Ala	465	470	475	1425	

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1622 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 51..1283

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CATCAGCCCA CCAGGAGACC TCGCCCGCCG CTCCCCGGG CTCCCCGGCC ATG TCT	56
Met Ser	
1	

		CCC	GCC	CGG	CTC	CGG	CCC	CGA	CTG	CAC	TTC	TGC	CTG	GTC	CTG	TTG	CTG	104
		Pro	Ala	Arg	Leu	Arg	Pro	Arg	Leu	His	Phe	Cys	Leu	Val	Leu	Leu	Leu	
				5					10					15				
5		CTG	CTG	GTG	GTG	CCC	GCG	GCA	TGG	GGC	TGC	GGG	CCG	GGT	CGG	GTG	GTG	152
		Leu	Leu	Val	Val	Pro	Ala	Ala	Trp	Gly	Cys	Gly	Pro	Gly	Arg	Val	Val	
			20					25					30					
10		GGC	AGC	CGC	CGG	CGA	CCG	CCA	CGC	AAA	CTC	GTG	CCG	CTC	GCC	TAC	AAG	200
		Gly	Ser	Arg	Arg	Arg	Pro	Pro	Arg	Lys	Leu	Val	Pro	Leu	Ala	Tyr	Lys	
			35				40				45						50	
15		CAG	TTC	AGC	CCC	AAT	GTG	CCC	GAG	AAG	ACC	CTG	GGC	GCC	AGC	GGA	CGC	248
		Gln	Phe	Ser	Pro	Asn	Val	Pro	Glu	Lys	Thr	Leu	Gly	Ala	Ser	Gly	Arg	
						55					60					65		
20		TAT	GAA	GGC	AAG	ATC	GCT	CGC	AGC	TCC	GAG	CGC	TTC	AAG	GAG	CTC	ACC	296
		Tyr	Glu	Gly	Lys	Ile	Ala	Arg	Ser	Ser	Glu	Arg	Phe	Lys	Glu	Leu	Thr	
					70					75					80			
25		CCC	AAT	TAC	AAT	CCA	GAC	ATC	ATC	TTC	AAG	GAC	GAG	GAG	AAC	ACA	GGC	344
		Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Thr	Gly	
				85				90						95				
30		GCC	GAC	CGC	CTC	ATG	ACC	CAG	CGC	TGC	AAG	GAC	CGC	CTG	AAC	TCG	CTG	392
		Ala	Asp	Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Arg	Leu	Asn	Ser	Leu	
			100					105					110					
35		GCT	ATC	TCG	GTG	ATG	AAC	CAG	TGG	CCC	GGT	GTG	AAG	CTG	CGG	GTG	ACC	440
		Ala	Ile	Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Lys	Leu	Arg	Val	Thr	
			115				120					125					130	
40		GAG	GGC	TGG	GAC	GAG	GAC	GGC	CAC	CAC	TCA	GAG	GAG	TCC	CTG	CAT	TAT	488
		Glu	Gly	Trp	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	His	Tyr	
					135						140					145		
45		GAG	GGC	CGC	GCG	GTG	GAC	ATC	ACC	ACA	TCA	GAC	CGC	GAC	CGC	AAT	AAG	536
		Glu	Gly	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Asn	Lys	
					150					155					160			
50		TAT	GGA	CTG	CTG	GCG	CGC	TTG	GCA	GTG	GAG	GCC	GGC	TTT	GAC	TGG	GTG	584
		Tyr	Gly	Leu	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	
				165				170						175				
55		TAT	TAC	GAG	TCA	AAG	GCC	CAC	GTG	CAT	TGC	TCC	GTC	AAG	TCC	GAG	CAC	632
		Tyr	Tyr	Glu	Ser	Lys	Ala	His	Val	His	Cys	Ser	Val	Lys	Ser	Glu	His	
			180					185					190					
60		TCG	GCC	GCA	GCC	AAG	ACG	GGC	GGC	TGC	TTC	CCT	GCC	GGA	GCC	CAG	GTA	680
		Ser	Ala	Ala	Ala	Lys	Thr	Gly	Gly	Cys	Phe	Pro	Ala	Gly	Ala	Gln	Val	
			195				200					205					210	
65		CGC	CTG	GAG	AGT	GGG	GCG	CGT	GTG	GCC	TTG	TCA	GCC	GTG	AGG	CCG	GGA	728
		Arg	Leu	Glu	Ser	Gly	Ala	Arg	Val	Ala	Leu	Ser	Ala	Val	Arg	Pro	Gly	
						215					220					225		
70		GAC	CGT	GTG	CTG	GCC	ATG	GGG	GAG	GAT	GGG	AGC	CCC	ACC	TTC	AGC	GAT	776
		Asp	Arg	Val	Leu	Ala	Met	Gly	Glu	Asp	Gly	Ser	Pro	Thr	Phe	Ser	Asp	
					230						235					240		

	GTG CTC ATT TTC CTG GAC CGC GAG CCC CAC AGG CTG AGA GCC TTC CAG	824
	Val Leu Ile Phe Leu Asp Arg Glu Pro His Arg Leu Arg Ala Phe Gln	
	245 250 255	
5	GTC ATC GAG ACT CAG GAC CCC CCA CGC CGC CTG GCA CTC ACA CCC GCT	872
	Val Ile Glu Thr Gln Asp Pro Pro Arg Arg Leu Ala Leu Thr Pro Ala	
	260 265 270	
10	CAC CTG CTC TTT ACG GCT GAC AAT CAC ACG GAG CCG GCA GCC CGC TTC	920
	His Leu Leu Phe Thr Ala Asp Asn His Thr Glu Pro Ala Ala Arg Phe	
	275 280 285 290	
15	CGG GCC ACA TTT GCC AGC CAC GTG CAG CCT GGC CAG TAC GTG CTG GTG	968
	Arg Ala Thr Phe Ala Ser His Val Gln Pro Gly Gln Tyr Val Leu Val	
	295 300 305	
20	GCT GGG GTG CCA GGC CTG CAG CCT GCC CGC GTG GCA GCT GTC TCT ACA	1016
	Ala Gly Val Pro Gly Leu Gln Pro Ala Arg Val Ala Ala Val Ser Thr	
	310 315 320	
25	CAC GTG GCC CTC GGG GCC TAC GCC CCG CTC ACA AAG CAT GGG ACA CTG	1064
	His Val Ala Leu Gly Ala Tyr Ala Pro Leu Thr Lys His Gly Thr Leu	
	325 330 335	
30	GTG GTG GAG GAT GTG GTG GCA TCC TGC TTC GCG GCC GTG GCT GAC CAC	1112
	Val Val Glu Asp Val Val Ala Ser Cys Phe Ala Ala Val Ala Asp His	
	340 345 350	
35	CAC CTG GCT CAG TTG GCC TTC TGG CCC CTG AGA CTC TTT CAC AGC TTG	1160
	His Leu Ala Gln Leu Ala Phe Trp Pro Leu Arg Leu Phe His Ser Leu	
	355 360 365 370	
40	GCA TGG GGC AGC TGG ACC CCG GGG GAG GGT GTG CAT TGG TAC CCC CAG	1208
	Ala Trp Gly Ser Trp Thr Pro Gly Glu Gly Val His Trp Tyr Pro Gln	
	375 380 385	
45	CTG CTC TAC CGC CTG GGG CGT CTC CTG CTA GAA GAG GGC AGC TTC CAC	1256
	Leu Leu Tyr Arg Leu Gly Arg Leu Leu Leu Glu Glu Gly Ser Phe His	
	390 395 400	
50	CCA CTG GGC ATG TCC GGG GCA GGG AGC TGAAAGGACT CCACCGCTGC	1303
	Pro Leu Gly Met Ser Gly Ala Gly Ser	
	405 410	
55	CCTCCTGGAA CTGCTGTACT GGGTCCAGAA GCCTCTCAGC CAGGAGGGAG CTGGCCCTGG	1363
	AAGGGACCTG AGCTGGGGGA CACTGGCTCC TGCCATCTCC TCTGCCATGA AGATACACCA	1423
	TTGAGACTTG ACTGGGCAAC ACCAGCGTCC CCCACCCGCG TCGTGGTGTA GTCATAGAGC	1483
	TGCAAGCTGA GCTGGCGAGG GGATGGTTGT TGACCCCTCT CTCCTAGAGA CCTTGAGGCT	1543
	GGCACGGCGA CTCCCAACTC AGCCTGCTCT CACTACGAGT TTTCATACTC TGCCTCCCCC	1603
	ATTGGGAGGG CCCATTCCC	1622

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1191 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: both

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1191

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

15	ATG GCT CTC CTG ACC AAT CTA CTG CCC TTG TGC TGC TTG GCA CTT CTG	48
	Met Ala Leu Leu Thr Asn Leu Leu Pro Leu Cys Cys Leu Ala Leu Leu	
	1 5 10 15	
20	GCG CTG CCA GCC CAG AGC TGC GGG CCG GGC CGG GGG CCG GTT GGC CGG	96
	Ala Leu Pro Ala Gln Ser Cys Gly Pro Gly Arg Gly Pro Val Gly Arg	
	20 25 30	
25	CGC CGC TAT GCG CGC AAG CAG CTC GTG CCG CTA CTC TAC AAG CAA TTT	144
	Arg Arg Tyr Ala Arg Lys Gln Leu Val Pro Leu Leu Tyr Lys Gln Phe	
	35 40 45	
30	GTG CCC GGC GTG CCA GAG CGG ACC CTG GGC GCC AGT GGG CCA GCG GAG	192
	Val Pro Gly Val Pro Glu Arg Thr Leu Gly Ala Ser Gly Pro Ala Glu	
	50 55 60	
35	GGG AGG GTG GCA AGG GGC TCC GAG CGC TTC CGG GAC CTC GTG CCC AAC	240
	Gly Arg Val Ala Arg Gly Ser Glu Arg Phe Arg Asp Leu Val Pro Asn	
	65 70 75 80	
40	TAC AAC CCC GAC ATC ATC TTC AAG GAT GAG GAG AAC AGT GGA GCC GAC	288
	Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Ser Gly Ala Asp	
	85 90 95	
45	CGC CTG ATG ACC GAG CGT TGC AAG GAG AGG GTG AAC GCT TTG GCC ATT	336
	Arg Leu Met Thr Glu Arg Cys Lys Glu Arg Val Asn Ala Leu Ala Ile	
	100 105 110	
50	GCC GTG ATG AAC ATG TGG CCC GGA GTG CGC CTA CGA GTG ACT GAG GGC	384
	Ala Val Met Asn Met Trp Pro Gly Val Arg Leu Arg Val Thr Glu Gly	
	115 120 125	
55	TGG GAC GAG GAC GGC CAC CAC GCT CAG GAT TCA CTC CAC TAC GAA GGC	432
	Trp Asp Glu Asp Gly His His Ala Gln Asp Ser Leu His Tyr Glu Gly	
	130 135 140	
60	CGT GCT TTG GAC ATC ACT ACG TCT GAC CGC GAC CGC AAC AAG TAT GGG	480
	Arg Ala Leu Asp Ile Thr Thr Ser Asp Arg Asp Arg Asn Lys Tyr Gly	
	145 150 155 160	
65	TTG CTG GCG CGC CTC GCA GTG GAA GCC GGC TTC GAC TGG GTC TAC TAC	528
	Leu Leu Ala Arg Leu Ala Val Glu Ala Gly Phe Asp Trp Val Tyr Tyr	
	165 170 175	
70	GAG TCC CGC AAC CAC GTC CAC GTG TCG GTC AAA GCT GAT AAC TCA CTG	576
	Glu Ser Arg Asn His Val His Val Ser Val Lys Ala Asp Asn Ser Leu	
	180 185 190	

	GCG GTC CGG GCG GGC GGC TGC TTT CCG GGA AAT GCA ACT GTG CGC CTG	624
	Ala Val Arg Ala Gly Gly Cys Phe Pro Gly Asn Ala Thr Val Arg Leu	
	195 200 205	
5	TGG AGC GGC GAG CGG AAA GGG CTG CGG GAA CTG CAC CGC GGA GAC TGG	672
	Trp Ser Gly Glu Arg Lys Gly Leu Arg Glu Leu His Arg Gly Asp Trp	
	210 215 220	
10	GTT TTG GCG GCC GAT GCG TCA GGC CGG GTG GTG CCC ACG CCG GTG CTG	720
	Val Leu Ala Ala Asp Ala Ser Gly Arg Val Val Pro Thr Pro Val Leu	
	225 230 235 240	
15	CTC TTC CTG GAC CGG GAC TTG CAG CGC CGG GCT TCA TTT GTG GCT GTG	768
	Leu Phe Leu Asp Arg Asp Leu Gln Arg Arg Ala Ser Phe Val Ala Val	
	245 250 255	
20	GAG ACC GAG TGG CCT CCA CGC AAA CTG TTG CTC ACG CCC TGG CAC CTG	816
	Glu Thr Glu Trp Pro Pro Arg Lys Leu Leu Leu Thr Pro Trp His Leu	
	260 265 270	
25	GTG TTT GCC GCT CGA GGG CCG GCG CCC GCG CCA GGC GAC TTT GCA CCG	864
	Val Phe Ala Ala Arg Gly Pro Ala Pro Ala Pro Gly Asp Phe Ala Pro	
	275 280 285	
30	GTG TTC GCG CGC CGG CTA CGC GCT GGG GAC TCG GTG CTG GCG CCC GGC	912
	Val Phe Ala Arg Arg Leu Arg Ala Gly Asp Ser Val Leu Ala Pro Gly	
	290 295 300	
35	GGG GAT GCG CTT CGG CCA GCG CGC GTG GCC CGT GTG GCG CGG GAG GAA	960
	Gly Asp Ala Leu Arg Pro Ala Arg Val Ala Arg Val Ala Arg Glu Glu	
	305 310 315 320	
40	GCC GTG GGC GTG TTC GCG CCG CTC ACC GCG CAC GGG ACG CTG CTG GTG	1008
	Ala Val Gly Val Phe Ala Pro Leu Thr Ala His Gly Thr Leu Leu Val	
	325 330 335	
45	AAC GAT GTC CTG GCC TCT TGC TAC GCG GTT CTG GAG AGT CAC CAG TGG	1056
	Asn Asp Val Leu Ala Ser Cys Tyr Ala Val Leu Glu Ser His Gln Trp	
	340 345 350	
50	GCG CAC CGC GCT TTT GCC CCC TTG AGA CTG CTG CAC GCG CTA GGG GCG	1104
	Ala His Arg Ala Phe Ala Pro Leu Arg Leu Leu His Ala Leu Gly Ala	
	355 360 365	
55	CTG CTC CCC GGC GGG GCC GTC CAG CCG ACT GGC ATG CAT TGG TAC TCT	1152
	Leu Leu Pro Gly Gly Ala Val Gln Pro Thr Gly Met His Trp Tyr Ser	
	370 375 380	
60	CGG CTC CTC TAC CGC TTA GCG GAG GAG CTA CTG GGC TG	1191
	Arg Leu Leu Tyr Arg Leu Ala Glu Glu Leu Leu Gly	
	385 390 395	

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1251 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: linear

```
(ix) FEATURE:
      (A) NAME/KEY: CDS
      (B) LOCATION: 1..1248
```

ATG GAC GTA AGG CTG CAT CTG AAG CAA TTT GCT TTA CTG TGT TTT ATC 48
Met Asp Val Arg Leu His Leu Lys Gln Phe Ala Leu Leu Cys Phe Ile
1 5 10 15

AGC TTG CTT CTG ACG CCT TGT GGA TTA GCC TGT GGT CCT GGT AGA GGT 96
Ser Leu Leu Leu Thr Pro Cys Gly Leu Ala Cys Gly Pro Gly Arg Gly
20 25 30

TAT GGA AAA CGA AGA CAC CCA AAG AAA TTA ACC CCG TTG GCT TAC AAG 144
Tyr Gly Lys Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys
35 40 45

CAA TTC ATC CCC AAC GTT GCT GAG AAA ACG CTT GGA GCC AGC GGC AAA 192
Gln Phe Ile Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Lys
50 55 60

TAC GAA GGC AAA ATC ACA AGG AAT TCA GAG AGA TTT AAA GAG CTG ATT 240
Tyr Glu Gly Lys Ile Thr Arg Asn Ser Glu Arg Phe Lys Glu Leu Ile
65 70 75 80

CCG AAT TAT AAT CCC GAT ATC ATC TTT AAG GAC GAG GAA AAC ACA AAC 288
Pro Asn Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Thr Asn
85 90 95

GCT GAC AGG CTG ATG ACC AAG CGC TGT AAG GAC AAG TTA AAT TCG TTG 336
Ala Asp Arg Leu Met Thr Lys Arg Cys Lys Asp Lys Leu Asn Ser Leu
100 105 110

GCC ATA TCC GTC ATG AAC CAC TGG CCC GGC GTG AAA CTG CGC GTC ACT 384
Ala Ile Ser Val Met Asn His Trp Pro Gly Val Lys Leu Arg Val Thr
115 120 125

GAA GGC TGG GAT GAG GAT GGT CAC CAT TTA GAA GAA TCT TTG CAC TAT 432
Glu Gly Trp Asp Glu Asp Gly His His Leu Glu Glu Ser Leu His Tyr
130 135 140

GAG GGA CGG GCA GTG GAC ATC ACT ACC TCA GAC AGG GAT AAA AGC AAG 480
Glu Gly Arg Ala Val Asp Ile Thr Thr Ser Asp Arg Asp Lys Ser Lys
145 150 155 160

TAT GGG ATG CTA TCC AGG CTT GCA GTG GAG GCA GGA TTC GAC TGG GTC 528
Tyr Gly Met Leu Ser Arg Leu Ala Val Glu Ala Gly Phe Asp Trp Val
165 170 175

TAT TAT GAA TCT AAA GCC CAC ATA CAC TGC TCT GTC AAA GCA GAA AAT 576
Tyr Tyr Glu Ser Lys Ala His Ile His Cys Ser Val Lys Ala Glu Asn
180 185 190

TCA GTG GCT GCT AAA TCA GGA GGA TGT TTT CCT GGG TCT GGG ACG GTG 624
Ser Val Ala Ala Lys Ser Gly Gly Cys Phe Pro Gly Ser Gly Thr Val

	195	200	205	
5	ACA CTT GGT GAT GGG ACG AGG AAA CCC ATC AAA GAT CTT AAA GTG GGC Thr Leu Gly Asp Gly Thr Arg Lys Pro Ile Lys Asp Leu Lys Val Gly 210 215 220			672
10	GAC CGG GTT TTG GCT GCA GAC GAG AAG GGA AAT GTC TTA ATA AGC GAC Asp Arg Val Leu Ala Ala Asp Glu Lys Gly Asn Val Leu Ile Ser Asp 225 230 235 240			720
	TTT ATT ATG TTT ATA GAC CAC GAT CCG ACA ACG AGA AGG CAA TTC ATC Phe Ile Met Phe Ile Asp His Asp Pro Thr Thr Arg Arg Gln Phe Ile 245 250 255			768
15	GTC ATC GAG ACG TCA GAA CCT TTC ACC AAG CTC ACC CTC ACT GCC GCG Val Ile Glu Thr Ser Glu Pro Phe Thr Lys Leu Thr Leu Thr Ala Ala 260 265 270			816
20	CAC CTA GTT TTC GTT GGA AAC TCT TCA GCA GCT TCG GGT ATA ACA GCA His Leu Val Phe Val Gly Asn Ser Ser Ala Ala Ser Gly Ile Thr Ala 275 280 285			864
25	ACA TTT GCC AGC AAC GTG AAG CCT GGA GAT ACA GTT TTA GTG TGG GAA Thr Phe Ala Ser Asn Val Lys Pro Gly Asp Thr Val Leu Val Trp Glu 290 295 300			912
30	GAC ACA TGC GAG AGC CTC AAG AGC GTT ACA GTG AAA AGG ATT TAC ACT Asp Thr Cys Glu Ser Leu Lys Ser Val Thr Val Lys Arg Ile Tyr Thr 305 310 315 320			960
	GAG GAG CAC GAG GGC TCT TTT GCG CCA GTC ACC GCG CAC GGA ACC ATA Glu Glu His Glu Gly Ser Phe Ala Pro Val Thr Ala His Gly Thr Ile 325 330 335			1008
35	ATA GTG GAT CAG GTG TTG GCA TCG TGC TAC GCG GTC ATT GAG AAC CAC Ile Val Asp Gln Val Leu Ala Ser Cys Tyr Ala Val Ile Glu Asn His 340 345 350			1056
40	AAA TGG GCA CAT TGG GCT TTT GCG CCG GTC AGG TTG TGT CAC AAG CTG Lys Trp Ala His Trp Ala Phe Ala Pro Val Arg Leu Cys His Lys Leu 355 360 365			1104
45	ATG ACG TGG CTT TTT CCG GCT CGT GAA TCA AAC GTC AAT TTT CAG GAG Met Thr Trp Leu Phe Pro Ala Arg Glu Ser Asn Val Asn Phe Gln Glu 370 375 380			1152
50	GAT GGT ATC CAC TGG TAC TCA AAT ATG CTG TTT CAC ATC GGC TCT TGG Asp Gly Ile His Trp Tyr Ser Asn Met Leu Phe His Ile Gly Ser Trp 385 390 395 400			1200
	CTG CTG GAC AGA GAC TCT TTC CAT CCA CTC GGG ATT TTA CAC TTA AGT Leu Leu Asp Arg Asp Ser Phe His Pro Leu Gly Ile Leu His Leu Ser 405 410 415			1248
55	TGA			1251

(2) INFORMATION FOR SEQ ID NO:10:

- 60 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 425 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

5	Met	Val	Glu	Met	Leu	Leu	Leu	Thr	Arg	Ile	Leu	Leu	Val	Gly	Phe	Ile
10	1				5					10					15	
	Cys	Ala	Leu	Leu	Val	Ser	Ser	Gly	Leu	Thr	Cys	Gly	Pro	Gly	Arg	Gly
				20					25					30		
15	Ile	Gly	Lys	Arg	Arg	His	Pro	Lys	Lys	Leu	Thr	Pro	Leu	Ala	Tyr	Lys
			35					40					45			
	Gln	Phe	Ile	Pro	Asn	Val	Ala	Glu	Lys	Thr	Leu	Gly	Ala	Ser	Gly	Arg
		50					55					60				
20	Tyr	Glu	Gly	Lys	Ile	Thr	Arg	Asn	Ser	Glu	Arg	Phe	Lys	Glu	Leu	Thr
	65					70					75					80
	Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Thr	Gly
25					85					90					95	
	Ala	Asp	Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Lys	Leu	Asn	Ala	Leu
				100					105					110		
30	Ala	Ile	Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Lys	Leu	Arg	Val	Thr
			115					120					125			
	Glu	Gly	Trp	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	His	Tyr
		130					135					140				
35	Glu	Gly	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Ser	Lys
	145					150					155					160
	Tyr	Gly	Met	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val
40					165					170					175	
	Tyr	Tyr	Glu	Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn
			180						185					190		
45	Ser	Val	Ala	Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Thr	Val
			195					200					205			
	His	Leu	Glu	His	Gly	Gly	Thr	Lys	Leu	Val	Lys	Asp	Leu	Ser	Pro	Gly
		210					215					220				
50	Asp	Arg	Val	Leu	Ala	Ala	Asp	Ala	Asp	Gly	Arg	Leu	Leu	Tyr	Ser	Asp
	225					230					235					240
	Phe	Leu	Thr	Phe	Leu	Asp	Arg	Met	Asp	Ser	Ser	Arg	Lys	Leu	Phe	Tyr
55					245					250					255	
	Val	Ile	Glu	Thr	Arg	Gln	Pro	Arg	Ala	Arg	Leu	Leu	Leu	Thr	Ala	Ala
				260					265					270		
60	His	Leu	Leu	Phe	Val	Ala	Pro	Gln	His	Asn	Gln	Ser	Glu	Ala	Thr	Gly
			275					280					285			

Ser Thr Ser Gly Gln Ala Leu Phe Ala Ser Asn Val Lys Pro Gly Gln
290 295 300

5 Arg Val Tyr Val Leu Gly Glu Gly Gly Gln Gln Leu Leu Pro Ala Ser
305 310 315 320

Val His Ser Val Ser Leu Arg Glu Glu Ala Ser Gly Ala Tyr Ala Pro
325 330 335

10 Leu Thr Ala Gln Gly Thr Ile Leu Ile Asn Arg Val Leu Ala Ser Cys
340 345 350

15 Tyr Ala Val Ile Glu Glu His Ser Trp Ala His Trp Ala Phe Ala Pro
355 360 365

Phe Arg Leu Ala Gln Gly Leu Leu Ala Ala Leu Cys Pro Asp Gly Ala
370 375 380

20 Ile Pro Thr Ala Ala Thr Thr Thr Thr Gly Ile His Trp Tyr Ser Arg
385 390 395 400

Leu Leu Tyr Arg Ile Gly Ser Trp Val Leu Asp Gly Asp Ala Leu His
405 410 415

25 Pro Leu Gly Met Val Ala Pro Ala Ser
420 425

30 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met Ala Leu Pro Ala Ser Leu Leu Pro Leu Cys Cys Leu Ala Leu Leu
1 5 10 15

45 Ala Leu Ser Ala Gln Ser Cys Gly Pro Gly Arg Gly Pro Val Gly Arg
20 25 30

50 Arg Arg Tyr Val Arg Lys Gln Leu Val Pro Leu Leu Tyr Lys Gln Phe
35 40 45

Val Pro Ser Met Pro Glu Arg Thr Leu Gly Ala Ser Gly Pro Ala Glu
50 55 60

55 Gly Arg Val Thr Arg Gly Ser Glu Arg Phe Arg Asp Leu Val Pro Asn
65 70 75 80

Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Ser Gly Ala Asp
85 90 95

60 Arg Leu Met Thr Glu Arg Cys Lys Glu Arg Val Asn Ala Leu Ala Ile

	100		105		110
	Ala Val Met Asn Met Trp Pro Gly Val Arg Leu Arg Val Thr Glu Gly				
	115		120		125
5	Trp Asp Glu Asp Gly His His Ala Gln Asp Ser Leu His Tyr Glu Gly				
	130		135		140
10	Arg Ala Leu Asp Ile Thr Thr Ser Asp Arg Asp Arg Asn Lys Tyr Gly				
	145		150		155
	Leu Leu Ala Arg Leu Ala Val Glu Ala Gly Phe Asp Trp Val Tyr Tyr				
		165		170	175
15	Glu Ser Arg Asn His Ile His Val Ser Val Lys Ala Asp Asn Ser Leu				
		180		185	190
	Ala Val Arg Ala Gly Gly Cys Phe Pro Gly Asn Ala Thr Val Arg Leu				
		195		200	205
20	Arg Ser Gly Glu Arg Lys Gly Leu Arg Glu Leu His Arg Gly Asp Trp				
		210		215	220
25	Val Leu Ala Ala Asp Ala Ala Gly Arg Val Val Pro Thr Pro Val Leu				
		225		230	235
	Leu Phe Leu Asp Arg Asp Leu Gln Arg Arg Ala Ser Phe Val Ala Val				
		245		250	255
30	Glu Thr Glu Arg Pro Pro Arg Lys Leu Leu Leu Thr Pro Trp His Leu				
		260		265	270
	Val Phe Ala Ala Arg Gly Pro Ala Pro Ala Pro Gly Asp Phe Ala Pro				
		275		280	285
35	Val Phe Ala Arg Arg Leu Arg Ala Gly Asp Ser Val Leu Ala Pro Gly				
		290		295	300
40	Gly Asp Ala Leu Gln Pro Ala Arg Val Ala Arg Val Ala Arg Glu Glu				
		305		310	315
	Ala Val Gly Val Phe Ala Pro Leu Thr Ala His Gly Thr Leu Leu Val				
		325		330	335
45	Asn Asp Val Leu Ala Ser Cys Tyr Ala Val Leu Glu Ser His Gln Trp				
		340		345	350
	Ala His Arg Ala Phe Ala Pro Leu Arg Leu Leu His Ala Leu Gly Ala				
		355		360	365
50	Leu Leu Pro Gly Gly Ala Val Gln Pro Thr Gly Met His Trp Tyr Ser				
		370		375	380
55	Arg Leu Leu Tyr Arg Leu Ala Glu Glu Leu Met Gly				
		385		390	395

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 411 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

10	Met	Ser	Pro	Ala	Trp	Leu	Arg	Pro	Arg	Leu	Arg	Phe	Cys	Leu	Phe	Leu
	1				5					10					15	
	Leu	Leu	Leu	Leu	Leu	Val	Pro	Ala	Ala	Arg	Gly	Cys	Gly	Pro	Gly	Arg
				20					25					30		
15	Val	Val	Gly	Ser	Arg	Arg	Arg	Pro	Pro	Arg	Lys	Leu	Val	Pro	Leu	Ala
			35					40					45			
	Tyr	Lys	Gln	Phe	Ser	Pro	Asn	Val	Pro	Glu	Lys	Thr	Leu	Gly	Ala	Ser
20		50					55					60				
	Gly	Arg	Tyr	Glu	Gly	Lys	Ile	Ala	Arg	Ser	Ser	Glu	Arg	Phe	Lys	Glu
	65					70				75						80
25	Leu	Thr	Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn
					85				90						95	
	Thr	Gly	Ala	Asp	Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Arg	Leu	Asn
			100						105					110		
30	Ser	Leu	Ala	Ile	Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Lys	Leu	Arg
			115					120					125			
	Val	Thr	Glu	Gly	Arg	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu
35		130					135					140				
	His	Tyr	Glu	Gly	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg
	145					150					155					160
40	Asn	Lys	Tyr	Gly	Leu	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp
				165					170						175	
	Trp	Val	Tyr	Tyr	Glu	Ser	Lys	Ala	His	Val	His	Cys	Ser	Val	Lys	Ser
			180						185					190		
45	Glu	His	Ser	Ala	Ala	Ala	Lys	Thr	Gly	Gly	Cys	Phe	Pro	Ala	Gly	Ala
			195					200					205			
	Gln	Val	Arg	Leu	Glu	Asn	Gly	Glu	Arg	Val	Ala	Leu	Ser	Ala	Val	Lys
50		210					215					220				
	Pro	Gly	Asp	Arg	Val	Leu	Ala	Met	Gly	Glu	Asp	Gly	Thr	Pro	Thr	Phe
	225					230					235					240
55	Ser	Asp	Val	Leu	Ile	Phe	Leu	Asp	Arg	Glu	Pro	Asn	Arg	Leu	Arg	Ala
				245						250					255	
	Phe	Gln	Val	Ile	Glu	Thr	Gln	Asp	Pro	Pro	Arg	Arg	Leu	Ala	Leu	Thr
			260						265					270		
60	Pro	Ala	His	Leu	Leu	Phe	Ile	Ala	Asp	Asn	His	Thr	Glu	Pro	Ala	Ala

275 280 285
 His Phe Arg Ala Thr Phe Ala Ser His Val Gln Pro Gly Gln Tyr Val
 290 295 300
 5 Leu Val Ser Gly Val Pro Gly Leu Gln Pro Ala Arg Val Ala Ala Val
 305 310 315 320
 10 Ser Thr His Val Ala Leu Gly Ser Tyr Ala Pro Leu Thr Arg His Gly
 325 330 335
 Thr Leu Val Val Glu Asp Val Val Ala Ser Cys Phe Ala Ala Val Ala
 340 345 350
 15 Asp His His Leu Ala Gln Leu Ala Phe Trp Pro Leu Arg Leu Phe Pro
 355 360 365
 Ser Leu Ala Trp Gly Ser Trp Thr Pro Ser Glu Gly Val His Ser Tyr
 370 375 380
 20 Pro Gln Met Leu Tyr Arg Leu Gly Arg Leu Leu Leu Glu Glu Ser Thr
 385 390 395 400
 25 Phe His Pro Leu Gly Met Ser Gly Ala Gly Ser
 405 410

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met Leu Leu Leu Leu Ala Arg Cys Phe Leu Val Ile Leu Ala Ser Ser
 1 5 10 15
 45 Leu Leu Val Cys Pro Gly Leu Ala Cys Gly Pro Gly Arg Gly Phe Gly
 20 25 30
 Lys Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys Gln Phe
 35 40 45
 50 Ile Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Arg Tyr Glu
 50 55 60
 55 Gly Lys Ile Thr Arg Asn Ser Glu Arg Phe Lys Glu Leu Thr Pro Asn
 65 70 75 80
 Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Thr Gly Ala Asp
 85 90 95
 60 Arg Leu Met Thr Gln Arg Cys Lys Asp Lys Leu Asn Ala Leu Ala Ile
 100 105 110

	Ser	Val	Met	Asn	Gln	Trp	Pro	Gly	Val	Arg	Leu	Arg	Val	Thr	Glu	Gly	
			115					120					125				
5	Trp	Asp	Glu	Asp	Gly	His	His	Ser	Glu	Glu	Ser	Leu	His	Tyr	Glu	Gly	
		130					135					140					
	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Ser	Lys	Tyr	Gly	
	145					150					155					160	
10	Met	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr	
					165					170					175		
	Glu	Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn	Ser	Val	
15				180					185					190			
	Ala	Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Thr	Val	His	Leu	
			195					200						205			
20	Glu	Gln	Gly	Gly	Thr	Lys	Leu	Val	Lys	Asp	Leu	Arg	Pro	Gly	Asp	Arg	
		210					215					220					
	Val	Leu	Ala	Ala	Asp	Asp	Gln	Gly	Arg	Leu	Leu	Tyr	Ser	Asp	Phe	Leu	
25		225				230					235					240	
	Thr	Phe	Leu	Asp	Arg	Asp	Glu	Gly	Ala	Lys	Lys	Val	Phe	Tyr	Val	Ile	
				245						250					255		
30	Glu	Thr	Leu	Glu	Pro	Arg	Glu	Arg	Leu	Leu	Leu	Thr	Ala	Ala	His	Leu	
			260						265					270			
	Leu	Phe	Val	Ala	Pro	His	Asn	Asp	Ser	Gly	Pro	Thr	Pro	Gly	Pro	Ser	
			275					280					285				
35	Ala	Leu	Phe	Ala	Ser	Arg	Val	Arg	Pro	Gly	Gln	Arg	Val	Tyr	Val	Val	
		290					295					300					
	Ala	Glu	Arg	Gly	Gly	Asp	Arg	Arg	Leu	Leu	Pro	Ala	Ala	Val	His	Ser	
40		305				310					315					320	
	Val	Thr	Leu	Arg	Glu	Glu	Glu	Ala	Gly	Ala	Tyr	Ala	Pro	Leu	Thr	Ala	
				325					330						335		
	His	Gly	Thr	Ile	Leu	Ile	Asn	Arg	Val	Leu	Ala	Ser	Cys	Tyr	Ala	Val	
45				340					345					350			
	Ile	Glu	Glu	His	Ser	Trp	Ala	His	Arg	Ala	Phe	Ala	Pro	Phe	Arg	Leu	
		355						360					365				
50	Ala	His	Ala	Leu	Leu	Ala	Ala	Leu	Ala	Pro	Ala	Arg	Thr	Asp	Gly	Gly	
		370					375					380					
	Gly	Gly	Gly	Ser	Ile	Pro	Ala	Ala	Gln	Ser	Ala	Thr	Glu	Ala	Arg	Gly	
	385					390					395					400	
55	Ala	Glu	Pro	Thr	Ala	Gly	Ile	His	Trp	Tyr	Ser	Gln	Leu	Leu	Tyr	His	
				405					410						415		
	Ile	Gly	Thr	Trp	Leu	Leu	Asp	Ser	Glu	Thr	Met	His	Pro	Leu	Gly	Met	
60				420					425					430			

Ala Val Lys Ser Ser
435

5 (2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 418 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met	Arg	Leu	Leu	Thr	Arg	Val	Leu	Leu	Val	Ser	Leu	Leu	Thr	Leu	Ser
1				5					10					15	
Leu	Val	Val	Ser	Gly	Leu	Ala	Cys	Gly	Pro	Gly	Arg	Gly	Tyr	Gly	Arg
			20					25					30		
Arg	Arg	His	Pro	Lys	Lys	Leu	Thr	Pro	Leu	Ala	Tyr	Lys	Gln	Phe	Ile
		35					40					45			
Pro	Asn	Val	Ala	Glu	Lys	Thr	Leu	Gly	Ala	Ser	Gly	Arg	Tyr	Glu	Gly
	50					55					60				
Lys	Ile	Thr	Arg	Asn	Ser	Glu	Arg	Phe	Lys	Glu	Leu	Thr	Pro	Asn	Tyr
65				70					75					80	
Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Thr	Gly	Ala	Asp	Arg
			85					90						95	
Leu	Met	Thr	Gln	Arg	Cys	Lys	Asp	Lys	Leu	Asn	Ser	Leu	Ala	Ile	Ser
			100					105					110		
Val	Met	Asn	His	Trp	Pro	Gly	Val	Lys	Leu	Arg	Val	Thr	Glu	Gly	Trp
		115					120					125			
Asp	Glu	Asp	Gly	His	His	Phe	Glu	Glu	Ser	Leu	His	Tyr	Glu	Gly	Arg
	130					135					140				
Ala	Val	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Lys	Ser	Lys	Tyr	Gly	Thr
145					150					155				160	
Leu	Ser	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr	Glu
			165						170					175	
Ser	Lys	Ala	His	Ile	His	Cys	Ser	Val	Lys	Ala	Glu	Asn	Ser	Val	Ala
			180					185					190		
Ala	Lys	Ser	Gly	Gly	Cys	Phe	Pro	Gly	Ser	Ala	Leu	Val	Ser	Leu	Gln
		195					200					205			
Asp	Gly	Gly	Gln	Lys	Ala	Val	Lys	Asp	Leu	Asn	Pro	Gly	Asp	Lys	Val
	210					215					220				
Leu	Ala	Ala	Asp	Ser	Ala	Gly	Asn	Leu	Val	Phe	Ser	Asp	Phe	Ile	Met
225					230					235				240	

5 Thr Gln Glu Pro Val Glu Lys Ile Thr Leu Thr Ala Ala His Leu Leu
260 265 270

10 Tyr Ala Ser Ser Val Arg Ala Gly Gln Lys Val Met Val Val Asp Asp
290 295 300

Gln Arg Gly Ser Phe Ala Pro Val Thr Ala His Gly Thr Ile Val Val
325 330 335

Ala His Leu Ala Phe Ala Pro Ala Arg Leu Tyr Tyr Tyr Val Ser Ser
355 360 365

Phe Leu Ser Pro Lys Thr Pro Ala Val Gly Pro Met Arg Leu Tyr Asn
370 375 380

Arg Arg Gly Ser Thr Gly Thr Pro Gly Ser Cys His Gln Met Gly Thr
385 390 395 400

Trp Leu Leu Asp Ser Asn Met Leu His Pro Leu Gly Met Ser Val Asn
405 410 415

Ser Ser

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 475 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Met Leu Leu Leu Ala Arg Cys Leu Leu Leu Val Leu Val Ser Ser Leu
1 5 10 15

Leu Val Cys Ser Gly Leu Ala Cys Gly Pro Gly Arg Gly Phe Gly Lys
20 25 30

Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys Gln Phe Ile
35 40 45

Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Arg Tyr Glu Gly

	50		55		60	
	Lys 65	Ile Ser Arg Asn Ser 70	Glu Arg Phe Lys Glu 75	Leu Thr Pro Asn Tyr 80		
5	Asn Pro Asp Ile 85	Ile Phe Lys Asp Glu 90	Glu Asn Thr Gly Ala Asp 95	Arg		
10	Leu Met Thr 100	Gln Arg Cys Lys Asp Lys 105	Leu Asn Ala Leu Ala 110	Ile Ser		
	Val Met Asn 115	Gln Trp Pro Gly Val 120	Lys Leu Arg Val Thr 125	Glu Gly Trp		
15	Asp Glu Asp 130	Gly His His Ser 135	Glu Glu Ser Leu His 140	Tyr Glu Gly Arg		
20	Ala Val Asp 145	Ile Thr Thr Ser Asp Arg 150	Asp Arg Ser Lys Tyr Gly 160	Met		
	Leu Ala Arg 165	Leu Ala Val Glu Ala Gly 170	Phe Asp Trp Val Tyr Tyr 175	Glu		
25	Ser Lys Ala 180	His Ile His Cys Ser 185	Val Lys Ala Glu Asn 190	Ser Val Ala		
	Ala Lys Ser 195	Gly Gly Cys Phe Pro 200	Gly Ser Ala Thr Val 205	His Leu Glu		
30	Gln Gly 210	Gly Thr Lys Leu Val 215	Lys Asp Leu Ser Pro 220	Gly Asp Arg Val		
	Leu Ala Ala 225	Asp Asp Gln Gly Arg Leu 230	Leu Tyr Ser Asp Phe Leu 240	Thr		
35	Phe Leu Asp 245	Arg Asp Asp Gly Ala Lys 250	Lys Val Phe Tyr Val 255	Ile Glu		
40	Thr Arg Glu 260	Pro Arg Glu Arg Leu 265	Leu Thr Ala Ala His 270	Leu Leu		
	Phe Val Ala 275	Pro His Asn Asp Ser 280	Ala Thr Gly Glu Pro 285	Glu Ala Ser		
45	Ser Gly 290	Ser Gly Pro Pro Ser 295	Gly Gly Ala Leu Gly 300	Pro Arg Ala Leu		
50	Phe Ala Ser 305	Arg Val Arg Pro Gly Gln Arg 310	Val Tyr Val Val Ala 320	Glu		
	Arg Asp Gly 325	Asp Arg Arg Leu Leu Pro 330	Ala Ala Val His Ser 335	Val Thr		
55	Leu Ser Glu 340	Glu Ala Ala Gly Ala Tyr 345	Ala Pro Leu Thr Ala 350	Gln Gly		
	Thr Ile Leu 355	Ile Asn Arg Val Leu 360	Ala Ser Cys Tyr Ala 365	Val Ile Glu		
60	Glu His 370	Ser Trp Ala His Arg 375	Ala Phe Ala Pro Phe 380	Arg Leu Ala His		

Ala Leu Leu Ala Ala Leu Ala Pro Ala Arg Thr Asp Arg Gly Gly Asp
385 390 395 400

5 Ser Gly Gly Gly Asp Arg Gly Gly Gly Gly Gly Arg Val Ala Leu Thr
405 410 415

Ala Pro Gly Ala Ala Asp Ala Pro Gly Ala Gly Ala Thr Ala Gly Ile
420 425 430

10 His Trp Tyr Ser Gln Leu Leu Tyr Gln Ile Gly Thr Trp Leu Leu Asp
435 440 445

15 Ser Glu Ala Leu His Pro Leu Gly Met Ala Val Lys Ser Ser Xaa Ser
450 455 460

Arg Gly Ala Gly Gly Gly Ala Arg Glu Gly Ala
465 470 475

20 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 411 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Met Ser Pro Ala Arg Leu Arg Pro Arg Leu His Phe Cys Leu Val Leu
1 5 10 15

Leu Leu Leu Leu Val Val Pro Ala Ala Trp Gly Cys Gly Pro Gly Arg
20 25 30

Val Val Gly Ser Arg Arg Arg Pro Pro Arg Lys Leu Val Pro Leu Ala
35 40 45

Tyr Lys Gln Phe Ser Pro Asn Val Pro Glu Lys Thr Leu Gly Ala Ser
50 55 60

Gly Arg Tyr Glu Gly Lys Ile Ala Arg Ser Ser Glu Arg Phe Lys Glu
65 70 75 80

45 Leu Thr Pro Asn Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn
85 90 95

50 Thr Gly Ala Asp Arg Leu Met Thr Gln Arg Cys Lys Asp Arg Leu Asn
100 105 110

Ser Leu Ala Ile Ser Val Met Asn Gln Trp Pro Gly Val Lys Leu Arg
115 120 125

55 Val Thr Glu Gly Trp Asp Glu Asp Gly His His Ser Glu Glu Ser Leu
130 135 140

His Tyr Glu Gly Arg Ala Val Asp Ile Thr Thr Ser Asp Arg Asp Arg
145 150 155 160

60 Asn Lys Tyr Gly Leu Leu Ala Arg Leu Ala Val Glu Ala Gly Phe Asp

				165					170					175			
	Trp	Val	Tyr	Tyr	Glu	Ser	Lys	Ala	His	Val	His	Cys	Ser	Val	Lys	Ser	
				180					185					190			
5	Glu	His	Ser	Ala	Ala	Ala	Lys	Thr	Gly	Gly	Cys	Phe	Pro	Ala	Gly	Ala	
			195					200					205				
10	Gln	Val	Arg	Leu	Glu	Ser	Gly	Ala	Arg	Val	Ala	Leu	Ser	Ala	Val	Arg	
		210					215					220					
	Pro	Gly	Asp	Arg	Val	Leu	Ala	Met	Gly	Glu	Asp	Gly	Ser	Pro	Thr	Phe	
	225					230					235					240	
15	Ser	Asp	Val	Leu	Ile	Phe	Leu	Asp	Arg	Glu	Pro	His	Arg	Leu	Arg	Ala	
				245						250					255		
	Phe	Gln	Val	Ile	Glu	Thr	Gln	Asp	Pro	Pro	Arg	Arg	Leu	Ala	Leu	Thr	
			260						265					270			
20	Pro	Ala	His	Leu	Leu	Phe	Thr	Ala	Asp	Asn	His	Thr	Glu	Pro	Ala	Ala	
			275					280					285				
25	Arg	Phe	Arg	Ala	Thr	Phe	Ala	Ser	His	Val	Gln	Pro	Gly	Gln	Tyr	Val	
		290					295					300					
	Leu	Val	Ala	Gly	Val	Pro	Gly	Leu	Gln	Pro	Ala	Arg	Val	Ala	Ala	Val	
	305					310					315					320	
30	Ser	Thr	His	Val	Ala	Leu	Gly	Ala	Tyr	Ala	Pro	Leu	Thr	Lys	His	Gly	
				325					330						335		
	Thr	Leu	Val	Val	Glu	Asp	Val	Val	Ala	Ser	Cys	Phe	Ala	Ala	Val	Ala	
			340						345					350			
35	Asp	His	His	Leu	Ala	Gln	Leu	Ala	Phe	Trp	Pro	Leu	Arg	Leu	Phe	His	
			355					360					365				
	Ser	Leu	Ala	Trp	Gly	Ser	Trp	Thr	Pro	Gly	Glu	Gly	Val	His	Trp	Tyr	
	370					375						380					
	Pro	Gln	Leu	Leu	Tyr	Arg	Leu	Gly	Arg	Leu	Leu	Leu	Glu	Glu	Gly	Ser	
	385					390					395					400	
45	Phe	His	Pro	Leu	Gly	Met	Ser	Gly	Ala	Gly	Ser						
				405						410							

(2) INFORMATION FOR SEQ ID NO:17:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 396 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

60	Met	Ala	Leu	Leu	Thr	Asn	Leu	Leu	Pro	Leu	Cys	Cys	Leu	Ala	Leu	Leu
	1				5				10					15		

	Ala	Leu	Pro	Ala	Gln	Ser	Cys	Gly	Pro	Gly	Arg	Gly	Pro	Val	Gly	Arg
				20					25					30		
5	Arg	Arg	Tyr	Ala	Arg	Lys	Gln	Leu	Val	Pro	Leu	Leu	Tyr	Lys	Gln	Phe
			35					40					45			
	Val	Pro	Gly	Val	Pro	Glu	Arg	Thr	Leu	Gly	Ala	Ser	Gly	Pro	Ala	Glu
		50					55					60				
10	Gly	Arg	Val	Ala	Arg	Gly	Ser	Glu	Arg	Phe	Arg	Asp	Leu	Val	Pro	Asn
	65					70				75						80
	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys	Asp	Glu	Glu	Asn	Ser	Gly	Ala	Asp
					85					90					95	
15	Arg	Leu	Met	Thr	Glu	Arg	Cys	Lys	Glu	Arg	Val	Asn	Ala	Leu	Ala	Ile
				100					105					110		
	Ala	Val	Met	Asn	Met	Trp	Pro	Gly	Val	Arg	Leu	Arg	Val	Thr	Glu	Gly
20			115					120					125			
	Trp	Asp	Glu	Asp	Gly	His	His	Ala	Gln	Asp	Ser	Leu	His	Tyr	Glu	Gly
	130						135					140				
25	Arg	Ala	Leu	Asp	Ile	Thr	Thr	Ser	Asp	Arg	Asp	Arg	Asn	Lys	Tyr	Gly
	145				150						155					160
	Leu	Leu	Ala	Arg	Leu	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr
					165				170						175	
30	Glu	Ser	Arg	Asn	His	Val	His	Val	Ser	Val	Lys	Ala	Asp	Asn	Ser	Leu
				180					185					190		
	Ala	Val	Arg	Ala	Gly	Gly	Cys	Phe	Pro	Gly	Asn	Ala	Thr	Val	Arg	Leu
35			195					200					205			
	Trp	Ser	Gly	Glu	Arg	Lys	Gly	Leu	Arg	Glu	Leu	His	Arg	Gly	Asp	Trp
	210					215						220				
40	Val	Leu	Ala	Ala	Asp	Ala	Ser	Gly	Arg	Val	Val	Pro	Thr	Pro	Val	Leu
	225				230					235						240
	Leu	Phe	Leu	Asp	Arg	Asp	Leu	Gln	Arg	Arg	Ala	Ser	Phe	Val	Ala	Val
				245					250						255	
45	Glu	Thr	Glu	Trp	Pro	Pro	Arg	Lys	Leu	Leu	Leu	Thr	Pro	Trp	His	Leu
				260					265					270		
	Val	Phe	Ala	Ala	Arg	Gly	Pro	Ala	Pro	Ala	Pro	Gly	Asp	Phe	Ala	Pro
50			275					280					285			
	Val	Phe	Ala	Arg	Arg	Leu	Arg	Ala	Gly	Asp	Ser	Val	Leu	Ala	Pro	Gly
		290					295					300				
55	Gly	Asp	Ala	Leu	Arg	Pro	Ala	Arg	Val	Ala	Arg	Val	Ala	Arg	Glu	Glu
	305					310					315					320
	Ala	Val	Gly	Val	Phe	Ala	Pro	Leu	Thr	Ala	His	Gly	Thr	Leu	Leu	Val
					325					330					335	
60	Asn	Asp	Val	Leu	Ala	Ser	Cys	Tyr	Ala	Val	Leu	Glu	Ser	His	Gln	Trp

340 345 350

Ala His Arg Ala Phe Ala Pro Leu Arg Leu Leu His Ala Leu Gly Ala
355 360 365

5 Leu Leu Pro Gly Gly Ala Val Gln Pro Thr Gly Met His Trp Tyr Ser
370 375 380

10 Arg Leu Leu Tyr Arg Leu Ala Glu Glu Leu Leu Gly
385 390 395

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 416 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Asp Val Arg Leu His Leu Lys Gln Phe Ala Leu Leu Cys Phe Ile
1 5 10 15

Ser Leu Leu Leu Thr Pro Cys Gly Leu Ala Cys Gly Pro Gly Arg Gly
20 25 30

Tyr Gly Lys Arg Arg His Pro Lys Lys Leu Thr Pro Leu Ala Tyr Lys
35 40 45

Gln Phe Ile Pro Asn Val Ala Glu Lys Thr Leu Gly Ala Ser Gly Lys
50 55 60

Tyr Glu Gly Lys Ile Thr Arg Asn Ser Glu Arg Phe Lys Glu Leu Ile
65 70 75 80

Pro Asn Tyr Asn Pro Asp Ile Ile Phe Lys Asp Glu Glu Asn Thr Asn
85 90 95

Ala Asp Arg Leu Met Thr Lys Arg Cys Lys Asp Lys Leu Asn Ser Leu
100 105 110

Ala Ile Ser Val Met Asn His Trp Pro Gly Val Lys Leu Arg Val Thr
115 120 125

Glu Gly Trp Asp Glu Asp Gly His His Leu Glu Glu Ser Leu His Tyr
130 135 140

Glu Gly Arg Ala Val Asp Ile Thr Thr Ser Asp Arg Asp Lys Ser Lys
145 150 155 160

Tyr Gly Met Leu Ser Arg Leu Ala Val Glu Ala Gly Phe Asp Trp Val
165 170 175

Tyr Tyr Glu Ser Lys Ala His Ile His Cys Ser Val Lys Ala Glu Asn
180 185 190

Ser Val Ala Ala Lys Ser Gly Gly Cys Phe Pro Gly Ser Gly Thr Val

	195	200	205
5	Thr Leu Gly Asp Gly Thr Arg Lys Pro Ile Lys Asp Leu Lys Val Gly 210 215 220		
	Asp Arg Val Leu Ala Ala Asp Glu Lys Gly Asn Val Leu Ile Ser Asp 225 230 235 240		
10	Phe Ile Met Phe Ile Asp His Asp Pro Thr Thr Arg Arg Gln Phe Ile 245 250 255		
	Val Ile Glu Thr Ser Glu Pro Phe Thr Lys Leu Thr Leu Thr Ala Ala 260 265 270		
15	His Leu Val Phe Val Gly Asn Ser Ser Ala Ala Ser Gly Ile Thr Ala 275 280 285		
	Thr Phe Ala Ser Asn Val Lys Pro Gly Asp Thr Val Leu Val Trp Glu 290 295 300		
20	Asp Thr Cys Glu Ser Leu Lys Ser Val Thr Val Lys Arg Ile Tyr Thr 305 310 315 320		
25	Glu Glu His Glu Gly Ser Phe Ala Pro Val Thr Ala His Gly Thr Ile 325 330 335		
	Ile Val Asp Gln Val Leu Ala Ser Cys Tyr Ala Val Ile Glu Asn His 340 345 350		
30	Lys Trp Ala His Trp Ala Phe Ala Pro Val Arg Leu Cys His Lys Leu 355 360 365		
	Met Thr Trp Leu Phe Pro Ala Arg Glu Ser Asn Val Asn Phe Gln Glu 370 375 380		
35	Asp Gly Ile His Trp Tyr Ser Asn Met Leu Phe His Ile Gly Ser Trp 385 390 395 400		
40	Leu Leu Asp Arg Asp Ser Phe His Pro Leu Gly Ile Leu His Leu Ser 405 410 415		

(2) INFORMATION FOR SEQ ID NO:19:

45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1416 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

55 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1413

60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ATG GAT AAC CAC AGC TCA GTG CCT TGG GCC AGT GCC GCC AGT GTC ACC

	Met	Asp	Asn	His	Ser	Ser	Val	Pro	Trp	Ala	Ser	Ala	Ala	Ser	Val	Thr	
	1				5					10					15		
5	TGT	CTC	TCC	CTG	GGA	TGC	CAA	ATG	CCA	CAG	TTC	CAG	TTC	CAG	TTC	CAG	96
	Cys	Leu	Ser	Leu	Gly	Cys	Gln	Met	Pro	Gln	Phe	Gln	Phe	Gln	Phe	Gln	
				20					25					30			
10	CTC	CAA	ATC	CGC	AGC	GAG	CTC	CAT	CTC	CGC	AAG	CCC	GCA	AGA	AGA	ACG	144
	Leu	Gln	Ile	Arg	Ser	Glu	Leu	His	Leu	Arg	Lys	Pro	Ala	Arg	Arg	Thr	
			35					40					45				
15	CAA	ACG	ATG	CGC	CAC	ATT	GCG	CAT	ACG	CAG	CGT	TGC	CTC	AGC	AGG	CTG	192
	Gln	Thr	Met	Arg	His	Ile	Ala	His	Thr	Gln	Arg	Cys	Leu	Ser	Arg	Leu	
		50					55					60					
20	ACC	TCT	CTG	GTG	GCC	CTG	CTG	CTG	ATC	GTC	TTG	CCG	ATG	GTC	TTT	AGC	240
	Thr	Ser	Leu	Val	Ala	Leu	Leu	Leu	Ile	Val	Leu	Pro	Met	Val	Phe	Ser	
	65					70				75						80	
25	CCG	GCT	CAC	AGC	TGC	GGT	CCT	GGC	CGA	GGA	TTG	GGT	CGT	CAT	AGG	GCG	288
	Pro	Ala	His	Ser	Cys	Gly	Pro	Gly	Arg	Gly	Leu	Gly	Arg	His	Arg	Ala	
					85					90					95		
30	CGC	AAC	CTG	TAT	CCG	CTG	GTC	CTC	AAG	CAG	ACA	ATT	CCC	AAT	CTA	TCC	336
	Arg	Asn	Leu	Tyr	Pro	Leu	Val	Leu	Lys	Gln	Thr	Ile	Pro	Asn	Leu	Ser	
				100					105					110			
35	GAG	TAC	ACG	AAC	AGC	GCC	TCC	GGA	CCT	CTG	GAG	GGT	GTG	ATC	CGT	CGG	384
	Glu	Tyr	Thr	Asn	Ser	Ala	Ser	Gly	Pro	Leu	Glu	Gly	Val	Ile	Arg	Arg	
			115					120					125				
40	GAT	TCG	CCC	AAA	TTC	AAG	GAC	CTC	GTG	CCC	AAC	TAC	AAC	AGG	GAC	ATC	432
	Asp	Ser	Pro	Lys	Phe	Lys	Asp	Leu	Val	Pro	Asn	Tyr	Asn	Arg	Asp	Ile	
		130					135					140					
45	CTT	TTC	CGT	GAC	GAG	GAA	GGC	ACC	GGA	GCG	GAT	GGC	TTG	ATG	AGC	AAG	480
	Leu	Phe	Arg	Asp	Glu	Glu	Gly	Thr	Gly	Ala	Asp	Gly	Leu	Met	Ser	Lys	
					150				155							160	
50	CGC	TGC	AAG	GAG	AAG	CTA	AAC	GTG	CTG	GCC	TAC	TCG	GTG	ATG	AAC	GAA	528
	Arg	Cys	Lys	Glu	Lys	Leu	Asn	Val	Leu	Ala	Tyr	Ser	Val	Met	Asn	Glu	
					165				170						175		
55	TGG	CCC	GGC	ATC	CGG	CTG	CTG	GTC	ACC	GAG	AGC	TGG	GAC	GAG	GAC	TAC	576
	Trp	Pro	Gly	Ile	Arg	Leu	Leu	Val	Thr	Glu	Ser	Trp	Asp	Glu	Asp	Tyr	
				180				185						190			
60	CAT	CAC	GGC	CAG	GAG	TCG	CTC	CAC	TAC	GAG	GGC	CGA	GCG	GTG	ACC	ATT	624
	His	His	Gly	Gln	Glu	Ser	Leu	His	Tyr	Glu	Gly	Arg	Ala	Val	Thr	Ile	
			195				200						205				
65	GCC	ACC	TCC	GAT	CGC	GAC	CAG	TCC	AAA	TAC	GGC	ATG	CTC	GCT	CGC	CTG	672
	Ala	Thr	Ser	Asp	Arg	Asp	Gln	Ser	Lys	Tyr	Gly	Met	Leu	Ala	Arg	Leu	
		210					215					220					
70	GCC	GTC	GAG	GCT	GGA	TTC	GAT	TGG	GTC	TCC	TAC	GTC	AGC	AGG	CGC	CAC	720
	Ala	Val	Glu	Ala	Gly	Phe	Asp	Trp	Val	Ser	Tyr	Val	Ser	Arg	Arg	His	
		225				230					235					240	
75	ATC	TAC	TGC	TCC	GTC	AAG	TCA	GAT	TCG	TCG	ATC	AGT	TCC	CAC	GTG	CAC	768
	Ile	Tyr	Cys	Ser	Val	Lys	Ser	Asp	Ser	Ser	Ile	Ser	Ser	His	Val	His	

	245	250	255	
5	GGC TGC TTC ACG CCG GAG AGC ACA GCG CTG CTG GAG AGT GGA GTC CGG Gly Cys Phe Thr Pro Glu Ser Thr Ala Leu Leu Glu Ser Gly Val Arg 260 265 270	816		
10	AAG CCG CTC GGC GAG CTC TCT ATC GGA GAT CGT GTT TTG AGC ATG ACC Lys Pro Leu Gly Glu Leu Ser Ile Gly Asp Arg Val Leu Ser Met Thr 275 280 285	864		
15	GCC AAC GGA CAG GCC GTC TAC AGC GAA GTG ATC CTC TTC ATG GAC CGC Ala Asn Gly Gln Ala Val Tyr Ser Glu Val Ile Leu Phe Met Asp Arg 290 295 300	912		
20	AAC CTC GAG CAG ATG CAA AAC TTT GTG CAG CTG CAC ACG GAC GGT GGA Asn Leu Glu Gln Met Gln Asn Phe Val Gln Leu His Thr Asp Gly Gly 305 310 315 320	960		
25	GCA GTG CTC ACG GTG ACG CCG GCT CAC CTG GTT AGC GTT TGG CAG CCG Ala Val Leu Thr Val Thr Pro Ala His Leu Val Ser Val Trp Gln Pro 325 330 335	1008		
30	GAG AGC CAG AAG CTC ACG TTT GTG TTT GCG CAT CGC ATC GAG GAG AAG Glu Ser Gln Lys Leu Thr Phe Val Phe Ala His Arg Ile Glu Glu Lys 340 345 350	1056		
35	AAC CAG GTG CTC GTA CGG GAT GTG GAG ACG GGC GAG CTG AGG CCC CAG Asn Gln Val Leu Val Arg Asp Val Glu Thr Gly Glu Leu Arg Pro Gln 355 360 365	1104		
40	CGA GTG GTC AAG TTG GGC AGT GTG CGC AGT AAG GGC GTG GTC GCG CCG Arg Val Val Lys Leu Gly Ser Val Arg Ser Lys Gly Val Val Ala Pro 370 375 380	1152		
45	CTG ACC CGC GAG GGC ACC ATT GTG GTC AAC TCG GTG GCC GCC AGT TGC Leu Thr Arg Glu Gly Thr Ile Val Val Asn Ser Val Ala Ala Ser Cys 385 390 395 400	1200		
50	TAT GCG GTG ATC AAC AGT CAG TCG CTG GCC CAC TGG GGA CTG GCT CCC Tyr Ala Val Ile Asn Ser Gln Ser Leu Ala His Trp Gly Leu Ala Pro 405 410 415	1248		
55	ATG CGC CTG CTG TCC ACG CTG GAG GCG TGG CTG CCC GCC AAG GAG CAG Met Arg Leu Leu Ser Thr Leu Glu Ala Trp Leu Pro Ala Lys Glu Gln 420 425 430	1296		
60	TTG CAC AGT TCG CCG AAG GTG GTG AGC TCG GCG CAG CAG CAG AAT GGC Leu His Ser Ser Pro Lys Val Val Ser Ser Ala Gln Gln Gln Asn Gly 435 440 445	1344		
65	ATC CAT TGG TAT GCC AAT GCG CTC TAC AAG GTC AAG GAC TAC GTG CTG Ile His Trp Tyr Ala Asn Ala Leu Tyr Lys Val Lys Asp Tyr Val Leu 450 455 460	1392		
70	CCG CAG AGC TGG CGC CAC GAT TGA Pro Gln Ser Trp Arg His Asp 465 470	1416		

60 (2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 471 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

10 Met Asp Asn His Ser Ser Val Pro Trp Ala Ser Ala Ala Ser Val Thr
 1 5 10 15
 Cys Leu Ser Leu Gly Cys Gln Met Pro Gln Phe Gln Phe Gln Phe Gln
 20 25 30
 15 Leu Gln Ile Arg Ser Glu Leu His Leu Arg Lys Pro Ala Arg Arg Thr
 35 40 45
 20 Gln Thr Met Arg His Ile Ala His Thr Gln Arg Cys Leu Ser Arg Leu
 50 55 60
 Thr Ser Leu Val Ala Leu Leu Ile Val Leu Pro Met Val Phe Ser
 65 70 75 80
 25 Pro Ala His Ser Cys Gly Pro Gly Arg Gly Leu Gly Arg His Arg Ala
 85 90 95
 Arg Asn Leu Tyr Pro Leu Val Leu Lys Gln Thr Ile Pro Asn Leu Ser
 100 105 110
 30 Glu Tyr Thr Asn Ser Ala Ser Gly Pro Leu Glu Gly Val Ile Arg Arg
 115 120 125
 Asp Ser Pro Lys Phe Lys Asp Leu Val Pro Asn Tyr Asn Arg Asp Ile
 130 135 140
 35 Leu Phe Arg Asp Glu Glu Gly Thr Gly Ala Asp Gly Leu Met Ser Lys
 145 150 155 160
 40 Arg Cys Lys Glu Lys Leu Asn Val Leu Ala Tyr Ser Val Met Asn Glu
 165 170 175
 Trp Pro Gly Ile Arg Leu Leu Val Thr Glu Ser Trp Asp Glu Asp Tyr
 180 185 190
 45 His His Gly Gln Glu Ser Leu His Tyr Glu Gly Arg Ala Val Thr Ile
 195 200 205
 Ala Thr Ser Asp Arg Asp Gln Ser Lys Tyr Gly Met Leu Ala Arg Leu
 210 215 220
 50 Ala Val Glu Ala Gly Phe Asp Trp Val Ser Tyr Val Ser Arg Arg His
 225 230 235 240
 55 Ile Tyr Cys Ser Val Lys Ser Asp Ser Ser Ile Ser Ser His Val His
 245 250 255
 Gly Cys Phe Thr Pro Glu Ser Thr Ala Leu Leu Glu Ser Gly Val Arg
 260 265 270
 60 Lys Pro Leu Gly Glu Leu Ser Ile Gly Asp Arg Val Leu Ser Met Thr

275 280 285

Ala Asn Gly Gln Ala Val Tyr Ser Glu Val Ile Leu Phe Met Asp Arg
290 295 300

5 Asn Leu Glu Gln Met Gln Asn Phe Val Gln Leu His Thr Asp Gly Gly
305 310 315 320

Ala Val Leu Thr Val Thr Pro Ala His Leu Val Ser Val Trp Gln Pro
325 330 335

10 Glu Ser Gln Lys Leu Thr Phe Val Phe Ala His Arg Ile Glu Glu Lys
340 345 350

15 Asn Gln Val Leu Val Arg Asp Val Glu Thr Gly Glu Leu Arg Pro Gln
355 360 365

Arg Val Val Lys Leu Gly Ser Val Arg Ser Lys Gly Val Val Ala Pro
370 375 380

20 Leu Thr Arg Glu Gly Thr Ile Val Val Asn Ser Val Ala Ala Ser Cys
385 390 395 400

25 Tyr Ala Val Ile Asn Ser Gln Ser Leu Ala His Trp Gly Leu Ala Pro
405 410 415

Met Arg Leu Leu Ser Thr Leu Glu Ala Trp Leu Pro Ala Lys Glu Gln
420 425 430

30 Leu His Ser Ser Pro Lys Val Val Ser Ser Ala Gln Gln Gln Asn Gly
435 440 445

Ile His Trp Tyr Ala Asn Ala Leu Tyr Lys Val Lys Asp Tyr Val Leu
450 455 460

35 Pro Gln Ser Trp Arg His Asp
465 470

40 (2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 221 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

55 Cys Gly Pro Gly Arg Gly Xaa Gly Xaa Arg Arg His Pro Lys Lys Leu
1 5 10 15

Thr Pro Leu Ala Tyr Lys Gln Phe Ile Pro Asn Val Ala Glu Lys Thr
20 25 30

60 Leu Gly Ala Ser Gly Arg Tyr Glu Gly Lys Ile Xaa Arg Asn Ser Glu
35 40 45

	Arg	Phe	Lys	Glu	Leu	Thr	Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile	Phe	Lys
	50						55					60				
5	Asp	Glu	Glu	Asn	Thr	Gly	Ala	Asp	Arg	Leu	Met	Thr	Gln	Arg	Cys	Lys
	65					70					75				80	
	Asp	Lys	Leu	Asn	Xaa	Leu	Ala	Ile	Ser	Val	Met	Asn	Xaa	Trp	Pro	Gly
				85						90					95	
10	Val	Xaa	Leu	Arg	Val	Thr	Glu	Gly	Trp	Asp	Glu	Asp	Gly	His	His	Xaa
				100					105					110		
	Glu	Glu	Ser	Leu	His	Tyr	Glu	Gly	Arg	Ala	Val	Asp	Ile	Thr	Thr	Ser
15			115					120					125			
	Asp	Arg	Asp	Xaa	Ser	Lys	Tyr	Gly	Xaa	Leu	Xaa	Arg	Leu	Ala	Val	Glu
	130						135					140				
20	Ala	Gly	Phe	Asp	Trp	Val	Tyr	Tyr	Glu	Ser	Lys	Ala	His	Ile	His	Cys
	145					150					155					160
	Ser	Val	Lys	Ala	Glu	Asn	Ser	Val	Ala	Ala	Lys	Ser	Gly	Gly	Cys	Phe
				165					170						175	
25	Pro	Gly	Ser	Ala	Xaa	Val	Xaa	Leu	Xaa	Xaa	Gly	Gly	Xaa	Lys	Xaa	Val
				180					185					190		
	Lys	Asp	Leu	Xaa	Pro	Gly	Asp	Xaa	Val	Leu	Ala	Ala	Asp	Xaa	Xaa	Gly
30			195					200					205			
	Xaa	Leu	Xaa	Xaa	Ser	Asp	Phe	Xaa	Xaa	Phe	Xaa	Asp	Arg			
	210						215					220				

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 167 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Cys	Gly	Pro	Gly	Arg	Gly	Xaa	Xaa	Xaa	Arg	Arg	Xaa	Xaa	Xaa	Pro	Lys
1				5					10					15	
Xaa	Leu	Xaa	Pro	Leu	Xaa	Tyr	Lys	Gln	Phe	Xaa	Pro	Xaa	Xaa	Xaa	Glu
			20					25					30		
Xaa	Thr	Leu	Gly	Ala	Ser	Gly	Xaa	Xaa	Glu	Gly	Xaa	Xaa	Xaa	Arg	Xaa
		35				40						45			
Ser	Glu	Arg	Phe	Xaa	Xaa	Leu	Thr	Pro	Asn	Tyr	Asn	Pro	Asp	Ile	Ile
50						55					60				


[illegible]

We claim:

1. A method for preventing degradation in functional performance of motor or sensory nerves in an animal comprising administering to the animal a therapeutic amount of a *hedgehog* or *ptc* therapeutic.
2. A method for preventing dysfunction of motor or sensory nerve cells comprising contacting the cells with an effective amount of a *hedgehog* or *ptc* therapeutic.
3. A method for treating or preventing peripheral neuroathy comprising administering to an animal a protective amount of a *hedgehog* or *ptc* therapeutic.
4. A method for protecting peripheral nerve cells under conditions which otherwise result in peripheral neuropathy, comprising administering to a patient in need thereof a therapeutically effective amount of a *hedgehog* or *ptc* therapeutic.
5. A method for the treating or preventing diabetic neuropathy comprising administering to a patient in need thereof a therapeutically effective amount of a *hedgehog* or *ptc* therapeutic.
6. A method for the treating or preventing virally-induced peripheral neuropathy comprising administering to a patient in need thereof a therapeutically effective amount of a *hedgehog* or *ptc* therapeutic.
7. The method of any of claims 1-6, wherein the *hedgehog* therapeutic is a polypeptide which includes a hedgehog amino acid sequence which is identical or homologous to an amino acid sequence of any one of SEQ ID Nos. 10-18.
8. The method of claim 7, wherein the hedgehog amino acid sequence is sufficient for specific binding of the polypeptide to a *patched* protein.
9. The method of claim 7, wherein the hedgehog amino acid sequence is at least 80 percent identical to an amino acid sequence of any one of SEQ ID Nos. 10-18.
10. The method of claim 7, wherein the hedgehog amino acid sequence is encodable by a nucleic acid which hybridizes under stringent conditions to any one of SEQ ID Nos. 1-9.
11. The method of claim 7, wherein the hedgehog amino acid sequence is of a vertebrate hedgehog protein.
12. The method of claim 11, wherein the vertebrate hedgehog protein is *Dhh*.
13. The method of claim 7, wherein the polypeptide includes at least a 50 amino acid extracellular portion of a vertebrate hedgehog protein.

14. The method of claim 7, wherein the polypeptide includes at least a 150 amino acid extracellular portion of a vertebrate hedgehog protein.
15. The method of claim 7, wherein the polypeptide includes at least an extracellular portion of a vertebrate hedgehog protein corresponding to residues 24-194 of SEQ ID No:15.
- 5 16. The method of claim 7, wherein the hedgehog polypeptide is modified with one or more lipophilic moieties.
17. The method of claim 16, wherein the hedgehog polypeptide is modified with one or more sterol moieties.
18. The method of claim 17, wherein the sterol moiety is cholesterol.
- 10 19. The method of claim 16, wherein the hedgehog polypeptide is modified with one or more fatty acid moieties.
20. The method of claim 19, wherein each fatty acid moiety is independently selected from the group consisting of myristoyl, palmitoyl, stearoyl, and arachidoyl.
- 15 21. The method of claim 16, wherein the hedgehog polypeptide is modified with one or more aromatic hydrocarbons.
22. The method of claim 21, wherein each aromatic hydrocarbon is ondependently selected from the group consisting of benzene, perylene, phenanthrene, anthracene, naphthalene, pyrene, chrysene, and naphthacene.
- 20 23. The method of claim 16, wherein the hedgehog polypeptide is modified one or more times with a C7 - C30 alkyl or cycloalkyl.
24. The method of of any of claims 1-6, wherein the *ptc* therapeutic is a small organic molecule.
25. The method of claim 24, wherein the binding of the *ptc* therapeutic to *patched* results in upregulation of patched and/or gli expression.
- 25 26. The method of any of claims 1-6, wherein the *ptc* therapeutic binds to *patched* and mimics *hedgehog*-mediated *patched* signal transduction.
27. The method of claim 26, wherein the *ptc* therapeutic is a small organic molecule.
28. The method of claim 26, wherein the binding of the *ptc* therapeutic to *patched* results in upregulation of patched and/or gli expression.

29. The method of any of claims 1-6, wherein the *ptc* therapeutic is a small organic molecule which interacts with neuronal cells to mimic *hedgehog*-mediated *patched* signal transduction.
30. The method of any of claims 1-6, wherein the *ptc* therapeutic mimics *hedgehog*-mediated *patched* signal transduction by altering the localization, protein-protein binding and/or enzymatic activity of an intracellular protein involved in a *patched* signal pathway.
31. The method of any of claims 1-6, wherein the *ptc* therapeutic alters the level of expression of a *hedgehog* protein, a *patched* protein or a protein involved in the intracellular signal transduction pathway of *patched*.
32. The method of claim 31, wherein the *ptc* therapeutic is an antisense construct which inhibits the expression of a protein which is involved in the signal transduction pathway of *patched* and the expression of which antagonizes *hedgehog*-mediated signals.
33. The method of claim 32, wherein the antisense construct is an oligonucleotide of about 20-30 nucleotides in length and having a GC content of at least 50 percent.
34. The method of claim 33, wherein the antisense oligonucleotide is selected from the group consisting of: 5'-GTCCTGGCGCCGCCGCCGCCGTCGCC;
5'-TTCCGATGACCGGCCTTTCGCGGTGA; and
5'-GTGCACGGAAAGGTGCAGGCCACACT
35. The method of claims 31, wherein the *ptc* therapeutic is a small organic molecule which binds to *patched* and regulates *patched*-dependent gene expression.
36. The method of claim 35, wherein the *ptc* therapeutic is an inhibitor of protein kinase A.
37. The method of claim 36, wherein the PKA inhibitor is a 5-isoquinolinesulfonamide
38. The method of claim 37, wherein the PKA inhibitor is represented in the general formula:



wherein,

R₁ and R₂ each can independently represent hydrogen, and as valence and stability permit a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, 5
-(CH₂)_m-R₈, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₈, -(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₈, or

R₁ and R₂ taken together with N form a heterocycle (substituted or unsubstituted);

10 R₃ is absent or represents one or more substitutions to the isoquinoline ring such as a lower alkyl, a lower alkenyl, a lower alkynyl, a carbonyl (such as a carboxyl, an ester, a formate, or a ketone), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an amino, an acylamino, an amido, a cyano, a nitro, an azido, a sulfate, a sulfonate, a sulfonamido, -(CH₂)_m-R₈, -(CH₂)_m-OH, -(CH₂)_m-O-lower alkyl, -(CH₂)_m-O-lower alkenyl, -(CH₂)_n-O-(CH₂)_m-R₈, 15
-(CH₂)_m-SH, -(CH₂)_m-S-lower alkyl, -(CH₂)_m-S-lower alkenyl, -(CH₂)_n-S-(CH₂)_m-R₈;

R₈ represents a substituted or unsubstituted aryl, aralkyl, cycloalkyl, cycloalkenyl, or heterocycle; and

n and m are independently for each occurrence zero or an integer in the range of 1 to 6.

39. The method of claim 36, wherein the PKA inhibitor is cyclic AMP analog.

20 40. The method of claim 36, wherein the PKA inhibitor is selected from the group consisting of N-[2-((p-bromocinnamyl)amino)ethyl]-5-isoquinolinesulfonamide, 1-(5-isoquinoline-sulfonyl)-2-methylpiperazine, KT5720, 8-bromo-cAMP, dibutyryl-cAMP and PKA Heat Stable Inhibitor isoform α.

41. The method of any of claims 4-6, wherein patient is being treated prophylactically.

25 42. A therapeutic preparation of a small molecule antagonist of *patched*, which *patched* antagonist is provided in a pharmaceutically acceptable carrier and in an amount sufficient to treat a peripheral neuropathy.

30 43. A method for protecting peripheral nerve cells under conditions which otherwise result in peripheral neuropathy, comprising administering to a patient a gene activation construct which recombines with a genomic *hedgehog* gene of the patient to provide a heterologous transcriptional regulatory sequence operatively linked to a coding sequence of the *hedgehog* gene.


44. The method of claim 4, 5, 6 or 43, which method is part of a protocol for the treatment of an acquired neuropathy.
45. The method of claim 44, wherein the neuropathy is due to viral infection, diabetes or inflammation.
46. The method of claim 44, wherein the neuropathy is due to contact with a toxic agent.
47. The method of claim 44, wherein the neuropathy is selected from the group consisting of diabetic neuropathy; immune-mediated neuropathy, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic polyneuropathy with antibodies to peripheral nerves, neuropathies associated with vasculitis or inflammation of the blood vessels in peripheral nerve, brachial or lumbosacral plexitis, and neuropathies associated with monoclonal gammopathies; neuropathies associated with tumors or neoplasms such as sensory neuropathy associated with lung cancer, neuropathy associated with multiple myeloma, neuropathy associated with waldenstrom's macroglobulemia, chronic lymphocytic leukemia, or B-cell lymphoma; neuropathy associated with amyloidosis; neuropathies caused by infections; neuropathies caused by nutritional imbalance; neuropathy in kidney disease; hypothyroid neuropathy; neuropathy caused by alcohol and toxins; neuropathies caused by drugs; neuropathy resulting from local irradiation; neuropathies caused by trauma or compression; and idiopathic neuropathies
48. The method of claim 4, 5, 6 or 43, which method is part of a protocol for the treatment of a hereditary neuropathy.
49. The method of claim 48, wherein the neuropathy is selected from the group consisting of Charcot-Marie Tooth Disease (CMT); Familial Amyloidotic Neuropathy and Hereditary Porphyria.
50. The method of claim 4, 5, 6 or 43, which method is part of a protocol for slowing neurodegenerative events associated with age-related neuropathology.
51. The method of claim 7, wherein the hedgehog polypeptide is a fusion protein.

5

10

The present application is directed to the discovery that hedgehog gene products are able to protect peripheral nerve cells under conditions which otherwise result in peripheral neuropathy. Certain aspects of the invention are directed to preparations of hedgehog polypeptides, or other molecules which regulate *patched* or *smoothened* signalling, and their uses as protective agents against both acquired and hereditary neuropathies. As used herein, "peripheral neuropathy" refers to a disorder affecting a segment of the peripheral nervous system. For instance, the method of the present invention can be used as part of a treatment program in the management of neuropathies associated with systemic disease, e.g., viral infections, diabetes, inflammation; as well as genetically acquired (hereditary) neuropathies, e.g., Charcot-Marie-Tooth disease; and neuropathies caused by a toxic agent, e.g., a chemotherapeutic agent such as vincristine.

Figure 1



668007-000000

Figure 2




Figure 3





Figure 4



66001-6625460

Figure 5




Figure 6




Figure 7




Figure 8





Figure 9







Figure 10A

Figure 10B






Figure 11A

Figure 11B




Figure 12

Functional Deficit in Peripheral
Nerve of Dhh $-/-$ Mice




Figure 13A

Peripheral Nerve Morphology in the Normal Mouse







Figure 13B

Peripheral Nerve Morphology in the Dhh $-/-$ Mouse





Figure 14A


Immunocytochemistry of Dhh $-/-$ Mouse Nerve



Neurofilament
(axonal marker)




Laminin - low mag.
(ECM/connect. tiss. marker)




Laminin - high mag.

Figure 14B

Immunocytochemistry of Normal Mouse Nerve




Neurofilament
(axonal marker)



Laminin
(ECM/connective tissue marker)

Figure 15

Hh Effects on Perineurial Cell Proliferation



- Dissociate P4 sciatic nerve
- Plate cells overnight
- Add Hh's and BrdU
- Fix 18-24 hours later
- Score cells




Figure 16






Figure 17



Mean \pm s.e.m. ** p < 0.01, *** p < 0.001.

Figure 18



Page twenty of twenty-five

Figure 19




Figure 20




Figure 21




Figure 22





Figure 23



DECLARATION FOR PATENT APPLICATION

Docket Number: **BIV-052.02**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Methods and Composition for Treating or Preventing Peripheral Neuropathies

the specification of which (check one): (X) is attached hereto.
() was filed on _____ as United States Application Number
or PCT International Application Number _____, and
was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulation, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)			Priority Claimed
_____	_____	_____	() Yes () No
(Number)	(Country)	(Day/Month/Year Filed)	
_____	_____	_____	() Yes () No
(Number)	(Country)	(Day/Month/Year Filed)	

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States Provisional application(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>09/187,387</u>	<u>November 6, 1998</u>	_____
(Application Number)	(Filing Date)	(Status: patent, pending, abandoned)
_____	_____	_____
(Application Number)	(Filing Date)	(Status: patent, pending, abandoned)

I hereby appoint Beth E. Arnold, Reg. No. 35,430; Paula Campbell, Reg. No. 32,503, Charles H. Cella, Reg. No. 38,099; Isabelle M. Clauss, Reg. (see attached); Edward J. Kelly, Reg. No. 38,936; Donald W. Muirhead, Reg. No. 33,978; Chinh Pham, Reg. No. 39,329; Anne Saturnelli, Reg. No. 41,290; Diana Steel, Reg. No. 43,153, Wolfgang Stutius, Reg. No. 40,256; Kingsley Taft, Reg. No. 43,946; Matthew P. Vincent, Reg. No. 36,709; and Anita Varma, Reg. No. 43,221 as attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

Address all telephone calls to Matthew P. Vincent at telephone number (617) 832-1000.

Address all correspondence to: Patent Group
 Foley, Hoag & Eliot LLP
 One Post Office Square
 Boston, Ma. 02109-2170

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor (given name, family name): Alphonse Galdes

Inventor's signature	Date
<u>One Douglas Road, Lexington, MA 02173</u>	<u>US</u>
Residence	Citizenship

Post Office Address
.....

Full name of second joint inventor, if any (given name, family name): Nagesh Mahanthappa

Inventor's signature	Date
<u>240 Norfolk Street, Cambridge, MA 02139</u>	<u>US</u>
Residence	Citizenship

Post Office Address
.....